












## PROCESS FOR PREPARING SUBSTITUED IMIDAZOLE DERIVATIVES AND PHARMACEUTICAL PREPARATIONS CONTAINING THEM

**Patent number:** HU218460  
**Publication date:** 2000-08-28  
**Inventor:** CARINI DAVID JOHN (US); DUNCIA JOHN JONAS VYTAUTAS (US)  
**Applicant:**  
**Classification:**  
**- international:** C07D233/64; C07D233/66; C07D233/68; C07D403/06; C07D403/10; C07D403/12; C07D233/84; A61K31/4174; A61P9/00; C07D233/91; C07D405/10; C07D405/12  
**- european:**  
**Application number:** HU19870003174 19870710  
**Priority number(s):** US19860884920 19860711; US19870050341 19870522

### Also published as:

 EP0253310 (A2)  
 SU1694062 (A3)  
 LU88662 (A)  
 FI873071 (A)  
 EP0253310 (A3)  
 EP0253310 (B1)  
 PT85312 (B)  
 IE871852L (L)  
 IE69984 (B1)  
 FI96025C (C)  
 FI96025B (B)

less <<

Abstract not available for HU218460  
Abstract of corresponding document: **EP0253310**

Substituted imidazoles such as  
<CHEM>

are useful as angiotensin II blockers. These compounds have activity in treating hypertension and congestive heart failure.

---

Data supplied from the **esp@cenet** database - Worldwide

**THIS PAGE BLANK (USPTO)**

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

**0 253 310 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 26.10.94 (51) Int. Cl.<sup>5</sup>: **C07D 233/68**, C07D 403/14,

(21) Application number: **87109919.8**

C07D 403/10, C07D 405/10,

C07D 403/06, C07D 233/64,

C07D 233/61, C07D 233/60,

C07D 233/84, C07D 233/70,

C07D 233/90

(22) Date of filing: **09.07.87**

The file contains technical information submitted  
after the application was filed and not included in  
this specification

(54) **Angiotensin II receptor blocking Imidazoles.**

(30) Priority: **11.07.86 US 884920**  
**22.05.87 US 50341**

(43) Date of publication of application:  
**20.01.88 Bulletin 88/03**

(45) Publication of the grant of the patent:  
**26.10.94 Bulletin 94/43**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

(56) References cited:  
**EP-A- 0 028 834**  
**EP-A- 0 142 754**  
**DE-A- 2 946 020**

(73) Proprietor: **E.I. DU PONT DE NEMOURS AND  
COMPANY**  
**1007 Market Street**  
**Wilmington Delaware 19898 (US)**

(72) Inventor: **Carini, David John**  
**1921 Julian Road**  
**Wilmington Delaware 19803 (US)**  
Inventor: **Duncia, John Jonas Vytautas**  
**35-B Martine Court**  
**Newark Delaware 19711 (US)**

(74) Representative: **von Kreisler, Alek,**  
**Dipl.-Chem. et al**  
**Patentanwälte**  
**von Kreisler-Selting-Werner**  
**Postfach 10 22 41**  
**D-50462 Köln (DE)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

**EP 0 253 310 B1**

## Description

## BACKGROUND OF THE INVENTION

5 Field of the Invention

This invention relates to novel substituted imidazoles, and processes for their preparation, pharmaceutical compositions containing them and pharmaceutical methods using them.


The compounds of this invention inhibit the action of the hormone angiotensin II (All) and are useful  
 10 therefore in alleviating angiotensin induced hypertension. The enzyme renin acts on a blood plasma  $\alpha_2$ -globulin, angiotensinogen, to produce angiotensin I, which is then converted by angiotensin converting-enzyme to All. The latter substance is a powerful vasopressor agent which has been implicated as a causative agent for producing high blood pressure in various mammalian species, such as the rat, dog, and man. The compounds of this invention inhibit the action of All at its receptors on target cells and thus  
 15 prevent the increase in blood pressure produced by this hormone-receptor interaction. By administering a compound of this invention to a species of mammal with hypertension due to All, the blood pressure is reduced. The compounds of this invention are also useful for the treatment of congestive heart failure.

K. Matsumura, et al., in U.S. Patent 4,207,324 issued June 10, 1980 discloses 1,2-disubstituted-4-haloimidazole-5-acetic acid derivatives of the formula:

20

25

30




Wherein R¹ is hydrogen, nitro or amino; R² is phenyl, furyl or thienyl optionally substituted by halogen,  
 35 lower alkyl, lower alkoxy or di-lower alkylamino; R³ is hydrogen or lower alkyl and X is halogen; and their physiologically acceptable salts. These compounds have diuretic and hypotensive actions.

Furukawa, et al., in U.S. Patent 4,355,040 issued October 19, 1982 discloses hypotensive imidazole-5-acetic acid derivatives having the formula:

40

45


50



Wherein R¹ is lower alkyl, cycloalkyl, or phenyl optionally substituted; X¹, X², and X³ are each hydrogen,  
 55 halogen, nitro, amino, lower alkyl, lower alkoxy, benzyloxy, or hydroxy; Y is halogen and R² is hydrogen or lower alkyl; and salts thereof.

Furukawa, et al., in U.S. Patent 4,340,598, issued July 20, 1982, discloses hypotensive imidazole derivatives of the formula:






5

Wherein R<sup>1</sup> is lower alkyl or, phenyl C<sub>1-2</sub> alkyl optionally substituted with halogen or nitro; R<sup>2</sup> is lower alkyl, cycloalkyl or phenyl optionally substituted; one of R<sup>3</sup> and R<sup>4</sup> is -(CH<sub>2</sub>)<sub>n</sub>COR<sup>5</sup> where R<sup>5</sup> is amino, lower alkoxy or hydroxyl and n is 0, 1, 2 and the other of R<sup>3</sup> and R<sup>4</sup> is hydrogen or halogen; provided that R<sup>1</sup> is lower alkyl or phenethyl when R<sup>3</sup> is hydrogen, n = 1 and R<sup>5</sup> is lower alkoxy or hydroxyl; and salts thereof.

10

Furukawa et al., in European Patent Application 103,647 discloses 4-chloro-2-phenylimidazole-5-acetic acid derivatives useful for treating edema and hypertension of the formula:

15



20

25

Where R represents lower alkyl and salts thereof.

30

The metabolism and disposition of hypotensive agent 4-chloro-1-(4-methoxy-3-methylbenzyl)-2-phenylimidazole-5-acetic acid is disclosed by H. Torii in Takeda Kenkyushoho, 41, No 3/4, 180-191 (1982).

Frazer et al., in European Patent Application 125,033-A discloses 1-phenyl(alkyl)-2-(alkyl)thioimidazole derivatives which are inhibitors of dopamine-β-hydroxylase and are useful as antihypertensives, diuretics and cardiotonics.

35

European Patent Application 146,228 filed October 16, 1984 by S.S.L. Parhi discloses a process for the preparation of 1-substituted-5-hydroxymethyl-2-mercaptoimidazoles.

A number of references disclose 1-benzylimidazoles such as U.S. Patent 4,448,781 to Cross and Dickinson (issued May 15, 1984); U.S. Patent 4,226,878 to Itzuka et al. (issued October 7, 1980); U.S. Patent 3,772,315 to Regel et al. (issued November 13, 1973); U.S. Patent 4,379,927 to Vorbrüggen et al. (issued April 12, 1983); amongst others.

40

Pals et al., Circulation Research, 29, 673 (1971) describe the introduction of a sarcosine residue in position 1 and alanine in position 8 of the endogenous vasoconstrictor hormone All to yield an (octa)peptide that blocks the effects of All on the blood pressure of pithed rats. This analog, [Sar<sup>1</sup>, Ala<sup>8</sup>] All, initially called "P-113" and subsequently "Saralasin", was found to be one of the most potent competitive antagonists of the actions of All, although, like most of the so-called peptide-All-antagonists, it also possesses agonistic actions of its own. Saralasin has been demonstrated to lower arterial pressure in mammals and man when the (elevated) pressure is dependent on circulating All (Pals et al., Circulation Research, 29, 673 (1971); Streeten and Anderson, Handbook of Hypertension, Vol. 5, Clinical Pharmacology of Antihypertensive Drugs, A. E. Doyle (Editor), Elsevier Science Publishers B.V., p. 246 (1984)). However, due to its agonistic character, saralasin generally elicits pressor effects when the pressure is not sustained by All. Being a peptide, the pharmacological effects to saralasin are relatively short-lasting and are only manifest after parenteral administration, oral doses being ineffective. Although the therapeutic uses of peptide All-blockers, like saralasin, are severely limited due to their oral ineffectiveness and short duration of action, their major utility is as a pharmaceutical standard.

45


50

55

EP-A-0 028 834 discloses 1-benzyl-imidazole-5-acetic acid derivatives having an antagonistic effect on angiotensin II and hypotensive activity.

Summary Of The Invention


According to the present invention there are provided novel compounds of formula (I) which have angiotensin II-antagonizing properties and are useful as antihypertensives.




(I)

wherein


R<sup>1</sup> is -4-CO<sub>2</sub>H; -4-CO<sub>2</sub>R<sup>9</sup>;




-SO<sub>3</sub>H, -C(CF<sub>3</sub>)<sub>2</sub>OH;



-PO<sub>3</sub>H;




4-NHSO<sub>2</sub>CH<sub>3</sub> ; -4-NHSO<sub>2</sub>CF<sub>3</sub>; -CONHOR<sup>12</sup>; -SO<sub>2</sub>NH<sub>2</sub> ;



R<sup>2</sup> is H; Cl; Br; I; F; NO<sub>2</sub>; alkyl of 1 to 4 carbon atoms; acyloxy of 1 to 4 carbon atoms;

alkoxy of 1 to 4 carbon atoms;  $\text{CO}_2\text{H}$ ;  $\text{CO}_2\text{R}^9$ ;  $\text{NHSO}_2\text{CH}_3$ ;  $\text{NHSO}_2\text{CF}_3$ ;  $\text{CONHOR}^{12}$ ;  $\text{SO}_2\text{NH}_2$  ;

5



10

aryl; or furyl;

$\text{R}^3$  is H; Cl, Br, I or F; alkyl of 1 to 4 carbon atoms or alkoxy of 1 to 4 carbon atoms;

$\text{R}^4$  is CN,  $\text{NO}_2$  or  $\text{CO}_2\text{R}^{11}$ ;

$\text{R}^5$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms alkenyl or alkynyl of 2 to 4 carbon atoms;

15

$\text{R}^6$  is alkyl of 2 to 10 carbon atoms, alkenyl or alkynyl of 3 to 10 carbon atoms or the same groups substituted with F or  $\text{CO}_2\text{R}^{14}$ ; cycloalkyl of 3 to 8 carbon atoms, cycloalkylalkyl, of 4 to 10 carbon atoms; cycloalkylalkenyl or cycloalkylalkynyl of 5 to 10 carbon atoms;  $(\text{CH}_2)_s\text{Z}(\text{CH}_2)_m\text{R}^5$  optionally substituted with F or  $\text{CO}_2\text{R}^{14}$ ; benzyl or benzyl substituted on the phenyl ring with 1 or 2 halogens, alkoxy of 1 to 4 carbon atoms, alkyl of 1 to 4 carbon atoms or nitro;


20

$\text{R}^7$  is H, F, Cl, Br, I,  $\text{NO}_2$ ,  $\text{CF}_3$  or CN;

$\text{R}^8$  is H, CN, alkyl of 1 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, or the same groups substituted with F; phenylalkenyl wherein the aliphatic portion is 2 to 6 carbon atoms;  $-(\text{CH}_2)_m$ -imidazol-1-yl;  $-(\text{CH}_2)_m$ -1,2,3-triazolyl optionally substituted with one or two groups selected from  $\text{CO}_2\text{CH}_3$  or alkyl of 1 to 4 carbon atoms;  $-(\text{CH}_2)_m$ -tetrazolyl;  $-(\text{CH}_2)_n\text{OR}^{11}$ ;


25

30




$-(\text{CH}_2)_n\text{SR}^{15}$ ;


35




40



45




50




55


$-(\text{CH}_2)_n\text{NR}^{11}\text{SO}_2\text{R}^{10}$ ;



5




10



15

R<sup>9</sup> is

20



R<sup>10</sup> is alkyl of 1 to 6 carbon atoms or perfluoroalkyl of 1 to 6 carbon atoms, 1-adamantyl, 1-naphthyl, 1-(1-naphthyl)ethyl, or (CH<sub>2</sub>)<sub>6</sub>C<sub>6</sub>H<sub>5</sub>;


25

R<sup>11</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;

R<sup>12</sup> is H, methyl or benzyl;

R<sup>13</sup> is -CO<sub>2</sub>H; -CO<sub>2</sub>R<sup>9</sup>; -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>R<sup>9</sup>;



30





35




40



45




50



55






10  $R^{14}$  is H, alkyl or perfluoroalkyl of 1 to 8 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;

$R^{15}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl, benzyl, acyl of 1 to 4 carbon atoms, phenacyl;

$R^{16}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms,  $(CH_2)_p C_6H_5$ ,  $OR^{17}$ , or  $NR^{18}R^{19}$ ;

15  $R^{17}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;


$R^{18}$  and  $R^{19}$  independently are H, alkyl of 1 to 4 carbon atoms, phenyl, benzyl,  $\alpha$ -methylbenzyl, or taken together form a ring of the formula



25 Q is  $NR^{20}$ , O or  $CH_2$ ;

$R^{20}$  is H, alkyl of 1-4 carbon atoms, or phenyl;


$R^{21}$  is alkyl of 1 to 6 carbon atoms,  $-NR^{22}R^{23}$ , or



35  $R^{22}$  and  $R^{23}$  independently are H, alkyl of 1 to 6 carbon atoms, benzyl, or are taken together as  $(CH_2)_u$  where u is 3-6;

$R^{24}$  is H,  $CH_3$  or  $-C_6H_5$ ;

$R^{25}$  is  $NR^{27}R^{28}$ ,  $OR^{28}$ ,  $NHCONH_2$ ,  $NHCSNH_2$ ,



45  $R^{26}$  is hydrogen, alkyl with from 1 to 6 carbon atoms, benzyl, or allyl;


$R^{27}$  and  $R^{28}$  are independently hydrogen, alkyl with from 1 to 5 carbon atoms, or phenyl;

$R^{29}$  and  $R^{30}$  are independently alkyl of 1-4 carbon atoms or taken together are  $-(CH_2)_q$ ;

$R^{31}$  is H, alkyl of 1 to 4 carbon atoms,  $-CH_2CH=CH_2$  or  $-CH_2C_6H_4R^{32}$ ;

$R^{32}$  is H,  $NO_2$ ,  $NH_2$ , OH or  $OCH_3$ ;


50 X is a carbon-carbon single bond,  $-CO-$ ,  $-O-$ ,  $-S-$ ,  $-NH-$ ,



$-OCH_2-$ ,  $-CH_2O-$ ,  $-SCH_2-$ ,  $-CH_2S-$ ,  $-NHC(R^{27})(R^{28})$ ,  $-NR^{23}SO_2-$ ,  $-SCH_2-$ ,  $-CH_2S-$ ,  $-NHC-$   
 $(R^{27})(R^{28})$ ,  $-NR^{23}SO_2-$ ,  $-SO_2NR^{23}-$ ,  $-C(R^{27})(R^{28})NH-$ ,  $-CH=CH-$ ,  $-CF=CF-$ ,  $-CH=CF-$ ,  
 $-CF=CH-$ ,  $-CH_2CH_2-$ ,  $-CF_2CF_2-$ ,



5




Y is O or S;  
 Z is O,  $\text{NR}^{11}$ , or S;  
 m is 1 to 5;  
 15 n is 1 to 10;  
 p is 0 to 3;  
 q is 2 to 3;  
 r is 0 to 2;  
 s is 0 to 5;  
 20 t is 0 or 1;

and pharmaceutically acceptable salts of these compounds;

provided that:


- (1) the  $\text{R}^1$  group is not in the ortho position;  
 (2) when  $\text{R}^1$  is

25



30


35 X is a single bond, and  $\text{R}^{13}$  is  $\text{CO}_2\text{H}$ , or



40

then  $\text{R}^{13}$  must be in the ortho or meta position; or when  $\text{R}^1$  and X are as above and  $\text{R}^{13}$  is  $\text{NHSO}_2\text{CF}_3$  or  $\text{NHSO}_2\text{CH}_3$ ,  $\text{R}^{13}$  must be ortho;

45 (3) when  $\text{R}^1$  is




50

55


and X is other than a single bond, then  $\text{R}^{13}$  must be ortho except when  $\text{X} = \text{NR}^{23}\text{CO}$  and  $\text{R}^{13}$  is  $\text{NHSO}_2\text{CF}_3$  or  $\text{NHSO}_2\text{CH}_3$ , then  $\text{R}^{13}$  must be ortho or meta;

(4) when  $\text{R}^1$  is 4- $\text{CO}_2\text{H}$  or a salt thereof,  $\text{R}^6$  cannot be S-alkyl;


- (5) when R<sup>1</sup> is 4-CO<sub>2</sub>H or a salt thereof,  
the substituent on the 4-position of the imidazole cannot be CH<sub>2</sub>OH, CH<sub>2</sub>OCOCH<sub>3</sub>, or CH<sub>2</sub>CO<sub>2</sub>H;  
(6) when R<sup>1</sup> is




- X is -OCH<sub>2</sub>-, and R<sup>13</sup> is 2-CO<sub>2</sub>H, and R<sup>7</sup> is H then R<sup>6</sup> is not C<sub>2</sub>H<sub>5</sub>S;  
(7) when R<sup>1</sup> is



- and R<sup>6</sup> is n-hexyl then R<sup>7</sup> and R<sup>8</sup> are not both hydrogen;  
(8) when R<sup>1</sup> is




- R<sup>6</sup> is not methoxybenzyl;  
(9) the R<sup>6</sup> group is not



or CH<sub>2</sub>OH.

Preferred for their antihypertensive activity are novel compounds having the formula:




Wherein


R<sup>1</sup> is -CO<sub>2</sub>H; -NHSO<sub>2</sub>CF<sub>3</sub>;



5



10




15

R<sup>6</sup> is alkyl of 3 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, alkynyl of 3 to 10 carbon atoms, cycloalkyl of 3 to 8 carbon atoms, benzyl substituted on the phenyl ring with up to two groups selected from alkoxy of 1 to 4 carbon atoms, halogen, alkyl of 1 to 4 carbon atoms, and nitro;


20

R<sup>8</sup> is phenylalkenyl wherein the aliphatic portion is 2 to 4 carbon atoms,  $-(CH_2)_m$ -imidazol-1-yl,  $-(CH_2)_m$ -1,2,3-triazolyl optionally substituted with one or two groups selected from  $CO_2CH_3$  or alkyl of 1 to 4 carbon atoms,  $(CH_2)_m$ -tetrazolyl,  $-(CH_2)_nOR^{11}$ ;



25




30



35




40



45


R<sup>13</sup> is  $-CO_2H$ ,  $-CO_2R^9$ ,  $NHSO_2CF_3$ ; and

50




55

R<sup>16</sup> is H, alkyl of 1 to 5 carbon atoms,  $OR^{17}$ , or  $NR^{18}R^{19}$ ;  
X is carbon-carbon single bond,  $-CO-$ ,



-CH<sub>2</sub>CH<sub>2</sub>-,




5

-OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHCH<sub>2</sub>-, -CH<sub>2</sub>NH- or -CH=CH-; and pharmaceutically acceptable salts of these compounds.


More preferred are compounds of the preferred scope where:

- 10 R<sup>2</sup> is H, alkyl of 1 to 4 carbon atoms, halogen, or alkoxy of 1 to 4 carbon atoms;  
 R<sup>6</sup> is alkyl, alkenyl or alkynyl of 3 to 7 carbon atoms;  
 R<sup>7</sup> is H, Cl, Br, or CF<sub>3</sub>;  
 R<sup>8</sup> is -(CH<sub>2</sub>)<sub>m</sub>OR<sup>11</sup>;

15




20



-(CH<sub>2</sub>)<sub>m</sub>NHSO<sub>2</sub>R<sup>10</sup>;


25



30

- or -COR<sup>16</sup>;  
 R<sup>10</sup> is CF<sub>3</sub>, alkyl of 1 to 6 carbon atoms or phenyl;  
 R<sup>11</sup> is H, or alkyl of 1 to 4 carbon atoms;  
 R<sup>13</sup> is CO<sub>2</sub>H; CO<sub>2</sub>CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>; NHSO<sub>2</sub>CF<sub>3</sub>  
 and


35



40

- R<sup>14</sup> is H, or alkyl of 1 to 4 carbon atoms;  
 R<sup>15</sup> is H, alkyl of 1 to 4 carbon atoms, or acyl of 1 to 4 carbon atoms;  
 R<sup>16</sup> is H, alkyl of 1 to 5 carbon atoms; OR<sup>17</sup>; or

45



50

m is 1 to 5;

X = single bond, -O-, -CO-, -NHCO-, or -OCH<sub>2</sub>-; and pharmaceutically acceptable salts.

55

Specifically preferred for their antihypertensive activity are:

- 2-Butyl-4-chloro-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole.
- 2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole.
- 2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(methoxycarbonyl)aminomethyl]imidazole.

- 2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(propoxycarbonyl)aminomethyl]imidazole.
- 2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde
- 2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde
- 2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole
- 5 • 2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde
- 2-Propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole
- 2-Propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde
- 2-Butyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde
- 2-(1E-Butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-hydroxymethylimidazole
- 10 • 2-(1E-Butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde and pharmaceutically acceptable salts thereof.

Note that throughout the text when an alkyl substituent is mentioned, the normal alkyl structure is meant (i.e., butyl is *n*-butyl) unless otherwise specified.

Also within the scope of this invention are pharmaceutical compositions comprising a suitable pharmaceutical carrier and a compound of Formula (I), and methods of using the compounds of Formula (I) to treat hypertension and congestive heart failure. The compounds of this invention can also be used as diagnostic agents to test the renin angiotensin system.

It should be noted in the foregoing structural formula, when a radical can be a substituent in more than one previously defined radical, that first radical can be selected independently in each previously defined radical. For example, R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> can each be CONHOR<sup>12</sup>. R<sup>12</sup> need not be the same substituent in each of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> but can be selected independently for each of them.

### Synthesis

25 The novel compounds of Formula (I) may be prepared using the reactions and techniques described in this section. The reactions are performed in a solvent appropriate to the reagents and materials employed and suitable for the transformation being effected. It is understood by those skilled in the art of organic synthesis that the functionality present on the imidazole and other portions of the molecule must be consistent with the chemical transformations proposed. This will frequently necessitate judgment as to the order of synthetic steps, protecting groups required, deprotection conditions, and activation of a benzylic position to enable attachment to nitrogen on the imidazole nucleus. Throughout the following section, not all compounds of formula (I) falling into a given class may necessarily be prepared by all methods described for that class. Substituents on the starting materials may be incompatible with some of the reaction conditions required in some of the methods described. Such restrictions to the substituents which are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternative methods described must then be used.


40

45

50

55

### Scheme 1




Generally, compounds of Formula (3) can be prepared by direct alkylation onto imidazole (1) prepared as described in U.S. 4,355,040 and references cited therein, with an appropriately protected benzyl halide, tosylate or mesylate (2) in the presence of base, as shown in path a). Preferably, the metallic imidazolide salt is prepared by reacting imidazole (1) with a proton acceptor such as MH where M is lithium, sodium or potassium in a solvent such as dimethylformamide (DMF) or by reacting it with a metal alkoxide of formula MOR where R is methyl, ethyl, t-butyl or the like in an alcohol solvent such as ethanol or t-butanol, or a


dipolar aprotic solvent such as dimethylformamide. The imidazole salt is dissolved in an inert aprotic solvent such as DMF, and treated with an appropriate alkylating agent (2). Alternatively, imidazole (1) can be alkylated with a benzyl halide (2, where X = Br, Cl) in the presence of a base such as sodium carbonate, potassium carbonate, triethylamine or pyridine. The reaction is run in an inert solvent such as DMF or DMSO at 20 °C to the reflux temperature of the solvent for 1-10 hours.

For example, the 4-nitrobenzyl intermediate (3a, wherein  $R^1 = 4\text{-NO}_2$ ,  $R^2 = R^3 = \text{H}$ ) may be obtained by direct alkylation onto imidazole (1) with a 4-nitrobenzyl halide, tosylate or mesylate in the presence of base.





If  $R^7$  and  $R^8$  are different, mixtures of two regioisomer alkylation products (3b, and 3c) are obtained in which  $R^7$  and  $R^8$  are interchanged. When  $R^8$  is CHO the alkylation is such that the benzyl group becomes attached to the adjacent nitrogen preferentially. These isomers possess distinct physical and biological properties and can usually be separated and isolated by conventional separation techniques such as chromatography and/or crystallization.



3b



3c



In all series examined, the more rapidly eluted isomer of a given pair has greater biological potency than the less rapidly eluted isomer. The absolute structure of the compounds 3d and 3e has been confirmed by X-ray crystallographic analysis to establish the relationship between structure, physical properties and biological activity. Sulfonamide 3d is the more rapidly eluted isomer in its series, acid 3e is the less rapidly eluted isomer in its series.

Alternatively, any properly functionalized benzylamine derivative (4) may be converted to imine (6) by treatment with an acylamino ketone (5) in the presence of an inert solvent such as benzene, toluene, or the like, and a catalytic amount of p-toluenesulfonic acid or molecular sieves, N. Engel, and W. Steglich, *Liebigs Ann. Chem.*, 1916, (1978), or in the presence of alumina, F. Texier-Boulet, *Synthesis*, 679 (1985). The resulting imine (6) can be cyclized to the N-benzyl imidazole (3) with phosphorus pentachloride ( $\text{PCl}_5$ ), phosphorus oxychloride ( $\text{POCl}_3$ ) or triphenylphosphine ( $\text{PPh}_3$ ) in dichloroethane in the presence of a base such as triethylamine, N. Engel and W. Steglich, *Liebigs Ann. Chem.*, 1916, (1978).

Acylamino ketone (5) is readily obtainable from amino acids via the Dakin-West reaction, H.D. Dakin, R. West, *J. Biol. Chem.*, 78, 95 and 745 (1928), and various modifications thereof, W. Steglich, G. Höfle, *Angew. Chem. Int. Ed. Engl.*, 8, 981 (1969); G. Höfle, W. Steglich, H. Vorbrüggen, *Angew. Chem. Int. Ed. Engl.*, 17, 569 (1978); W. Steglich, G. Höfle, *Ber.*, 102, 883 (1969), or by selective reduction of acyl cyanides, A. Pfaltz, S. Anwar, *Tet. Lett.* 2977 (1984), or from  $\alpha$ -halo,  $\alpha$ -tosyl or  $\alpha$ -mesyl ketones via the appropriate substitution reactions that one skilled in the art will readily recognize.

The functionalized benzylamines (4) may be made from the corresponding benzyl halide, tosylate or mesylate (2) via displacement with a nitrogen nucleophile, a procedure familiar to one skilled in the art. This displacement may be achieved using azide ion, ammonia, or phthalimide anion, etc., in a neutral solvent such as dimethylformamide, dimethylsulfoxide etc., or under phase transfer conditions. The benzyl halide (2) may be made by a variety of benzylic halogenation methods familiar to one skilled in the art, for example benzylic bromination of toluene derivatives with N-bromosuccinimide in an inert solvent such as carbon tetrachloride in the presence of a radical initiator such as benzoyl peroxide at temperatures up to reflux conditions.

A wide variety of toluene derivatives may be made from simple electrophilic substitution reactions on an aromatic ring. This includes nitration, sulfonation, phosphorylation, Friedel-Crafts alkylation, Friedel-Crafts acylation, halogenation, and other similar reactions known to one skilled in the art, G. A. Olah, "Friedel-Crafts and Related Reactions," Vol. 1-5, Interscience, New York, (1965).

Another way to synthesize functionalized benzyl halides is via chloromethylation of the corresponding aromatic precursor. Thus, the appropriately substituted benzene ring may be chloromethylated with formaldehyde and hydrochloric acid (HCl) for example with or without an inert solvent such as chloroform, carbon tetrachloride, light petroleum ether or acetic acid. A Lewis acid such as zinc chloride ( $\text{ZnCl}_2$ ) or a mineral acid such as phosphoric acid may also be added as a catalyst or condensing agent, R. C. Fuson, C. H. McKeever, *Org. Reactions*, 1, 63 (1942).

Alternatively, N-benzylimidazoles (3) can also be prepared as shown in path b) by forming an  $\text{R}^6$  substituted amidine (7) from an appropriately substituted benzylamine (4) which is in turn reacted with an  $\alpha$ -haloketone,  $\alpha$ -hydroxyketone (8),  $\alpha$ -haloaldehyde, or  $\alpha$ -hydroxyaldehyde, F. Kunckell, *Ber.*, 34, 637 (1901).

As shown in path a), imidazole (1) may be alkylated by a variety of benzyl derivatives. These include compounds with latent acid functionalities such as o, m, and p-cyanobenzylhalides, mesylates or tosylates as shown in path c). Nitriles of formula (9) may be hydrolyzed to carboxylic acids of formula (10) by treatment with strong acid or alkali. Preferably, treatment with a 1:1 (v/v) mixture of concentrated aqueous hydrochloric acid/glacial acetic acid at reflux temperatures for 2-96 hours or by treatment with 1N sodium hydroxide in an alcohol solvent such as ethanol or ethylene glycol for 2-96 hours at temperatures from 20°C to reflux can be used. If another nitrile group is present it will also be hydrolyzed. The nitrile functionality can also be hydrolyzed in two steps by first stirring in sulfuric acid to form the amide followed by hydrolysis with sodium hydroxide or a mineral acid to give the carboxylic acid (10).

The nitriles (9) can be converted into the corresponding tetrazole derivative (11) by a variety of methods using hydrazoic acid. For example, the nitrile can be heated with sodium azide and ammonium chloride in DMF at temperatures between 30°C and reflux for 1-10 days, J. P. Hurwitz and A. J. Tomson, *J. Org. Chem.*, 26, 3392 (1961). Preferably, the tetrazole is prepared by the 1,3-dipolar cycloaddition of trialkyltin or triaryltin azides to the appropriately substituted nitrile as described in detail by Scheme 15.

The starting imidazole compounds (1) are readily available by any of a number of standard methods. For example, acylaminoketone (5) can be cyclized with ammonia or equivalents thereof, D. Davidson, et al., *J. Org. Chem.*, 2, 319 (1937) to the corresponding imidazole as shown in Scheme 1. The corresponding oxazole can also be converted to imidazole (1) by action of ammonia or amines in general, H. Brederick, et al., *Ber.*, 88, 1351 (1955); J. W. Cornforth and R. H. Cornforth, *J. Chem Soc.*, 96, (1947).

Several alternative routes to imidazoles (1) are illustrated in Scheme 2. As shown in Scheme 2 equation a), reaction of the appropriate  $\text{R}^6$  substituted imidate esters (12) with an appropriately substituted  $\alpha$ -hydroxy- or  $\alpha$ -haloketone or aldehyde (8) in ammonia leads to imidazoles of formula (1), P. Dziuron, and W. Schunack, *Archiv. Pharmaz.*, 307 and 470 (1974).

The starting imidazole compounds (1) wherein  $\text{R}^7$  and  $\text{R}^8$  are both hydrogen can be prepared as shown in equation b) by reaction of the appropriate  $\text{R}^6$ -substituted imidate ester (12) with  $\alpha$ -aminoacetaldehyde dimethyl acetal (13), M. R. Grimmett, *Adv. Heterocyclic Chem.*, 12, 103 (1970).

As shown in equation c), imidazole (15; wherein  $\text{R}^7$  = hydrogen and  $\text{R}^8$  =  $\text{CH}_2\text{OH}$ ) can be prepared by treatment of the imidate ester (12) with 1,3-dihydroxyacetone (14) in ammonia by the procedure described in *Archive der Pharmazie*, 307, 470 (1974). Halogenation of imidazole (15) or any imidazole wherein  $\text{R}^7$  or  $\text{R}^8$  is hydrogen is preferably accomplished by reaction with one to two equivalents of N-halosuccinimide in a polar solvent such as dioxane or 2-methoxyethanol at a temperature of 40-100°C for 1-10 hours. Reaction of the halogenated imidazole (16) with a benzylhalide (2) in the manner described in Scheme 1 affords the corresponding N-benzylimidazole (17); wherein  $\text{R}^7$  is halogen and  $\text{R}^8$  is  $\text{CH}_2\text{OH}$ . This procedure is described in U.S. Patent 4,355,040. Alternatively, imidazole (17) can be prepared by the procedure described in U.S. Patent 4,207,324.

Compounds of formula (17) can also be prepared by treatment of the starting imidazole compound (1) wherein  $\text{R}^7$  and  $\text{R}^8$  are both hydrogen, with the appropriate benzyl halide followed by functionalization of  $\text{R}^7$

and R<sup>8</sup> by treatment with formaldehyde as described in E. F. Godefroi, et al., *Recueil*, 91, 1383 (1972) followed by halogenation as was described above.

As shown in equation d) the imidazoles (1) can also be prepared by reaction of R<sup>6</sup> substituted amidines (18) with an  $\alpha$ -hydroxy- or  $\alpha$ -haloketone or aldehyde (8) as described by F. Kunckel, *Ber.*, 34, 637, (1901).

5 As shown in equation e), preparation of the nitroimidazoles (1, R<sup>7</sup> or R<sup>8</sup> = NO<sub>2</sub>) is preferably accomplished by heating the appropriate starting imidazole in a 3:1 mixture of conc. sulfuric acid/conc. nitric acid at 60-100 °C for 1-6 hours. Nitration of the imidazole (15) can be achieved by first converting the hydroxymethylimidazole to the corresponding chloromethylimidazole (22) employing thionyl chloride or oxalyl chloride. Nitration, as described above, followed by hydrolysis provides the nitroimidazoles (24).

10 Imidazoles (21) where R<sup>7</sup> and R<sup>8</sup> = CN can be prepared as shown in equation f) by reaction of R<sup>6</sup> substituted ortho esters, ortho acids or aldehydes (followed by oxidation of the aldehyde) with diaminomaleonitrile (20) by the procedure described by R. W. Begland et al., *J. Org. Chem.*, 39, 2341 (1974). Likewise, R<sup>6</sup> substituted imidate esters (12) also react with diaminomaleonitrile to give 4,5 dicyanoimidazoles (21). The nitrile groups can be further elaborated into other functional groups by  
15 methods familiar to one skilled in the art.

20

25

30

35

40

45

50

55



## Scheme 2

5

10

15

20

25

30


35

40



45

50


55







Scheme 3



As shown in Scheme 3, path a) for benzylimidazoles (17) where  $R^7 = Cl$  and  $R^8 = CH_2OH$ , the hydroxymethyl groups may be easily converted to the corresponding halide, mesylate or tosylate by a variety of methods familiar to one skilled in the art. Preferably, the alcohol (17) is converted to the chloride (25) with thionyl chloride in an inert solvent at temperatures of  $20^\circ C$  to the reflux temperature of the solvent.

Chloride (25) may be displaced by a variety of nucleophiles by nucleophilic displacement reaction procedures familiar to one skilled in the art. For example, excess sodium cyanide in DMSO may be used to form cyanomethyl derivatives (26) at temperatures of 20 °C to 100 °C.

Nitrile (26) may be hydrolyzed to acetic acid derivative (27), by a variety of methods. These methods include methods described previously for the hydrolysis of nitriles of formula (9). Examples of desired acids and bases for this hydrolysis include mineral acids such as sulfuric acid, hydrochloric acid, and mixtures of either of the above with 30-50% acetic acid (when solubility is a problem), and alkali metal hydroxides such as sodium hydroxide or potassium hydroxide. The hydrolysis reaction proceeds under heating at temperatures ranging from 50-160 °C for 2-48 hours. Carboxylic acid (27) may be esterified by a variety of methods without affecting other parts of the molecule. Preferably, (27) is refluxed in a hydrochloric acid/methanol solution for 2-48 hours to give ester (28).

Ester (28) may be hydrolyzed to carboxylic acid (27), for instance, after R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> have been elaborated. Various methods, acidic or basic, may be used. For example, compound (28) is stirred with 0.5N potassium hydroxide in methanol, or if base soluble, it is stirred in 1.0N sodium hydroxide for 1-48 h at 20 °C to reflux temperatures.

Hydroxymethyl derivative (17) may be acylated to give (29) by a variety of procedures. As shown in path b) acylation can be achieved with 1-3 equivalents of an acyl halide or an anhydride in a solvent such as diethyl ether, tetrahydrofuran, methylene chloride or the like in the presence of a base such as pyridine or triethylamine. Alternatively (17) may be acylated by reaction with a carboxylic acid and dicyclohexylcarbodiimide (DCC) in the presence of a catalytic amount of 4-(N,N-dimethylamino)pyridine (DMAP) via the procedure described by A. Hassner, *Tet. Lett.*, 46, 4475 (1978). Treatment of (17) with a solution of carboxylic acid anhydride in pyridine optionally with a catalytic amount of DMAP at temperatures of 20-100 °C for 2-48 hours is the preferred method.

The ether (30) can be prepared from the alcohol (17) as shown in path c) by methods such as treatment of (17) in a solvent such as dimethylformamide or dimethylsulfoxide with potassium *t*-butoxide, sodium hydride, or the like followed by treatment with R<sup>11</sup>L at 25 °C for 1-20 hours, where L is a halogen, tosylate or mesylate.

Alternatively, treatment of (17) with 1-5 equivalents of thionyl chloride in chloroform for 2-6 hours at 25 °C followed by treatment of the intermediate (25) with 1-3 equivalents of MOR<sup>11</sup>, where M is sodium or potassium, for 2-10 hours at 25 °C either in R<sup>11</sup>OH as solvent or in a polar solvent such as dimethylformamide or the like will also yield ether (30).

The ether (30) can also be prepared for example by heating (17) for 3-15 hours at 60-160 °C in R<sup>11</sup>OH containing an inorganic acid such as a hydrochloric acid or sulfuric acid.

Compound (17) can be dehalogenated to compound (31) preferably by catalytic hydrogenolysis (over an appropriate catalyst such as 10% palladium on carbon) in methanol at 25 °C for 1-6 hours or by treatment with zinc metal in acetic acid.

As shown in Scheme 3, the trifluoromethyl imidazoles (33) can be prepared from the corresponding iodoimidazoles (32) by treatment with trifluoromethyl copper, *J. Am. Chem. Soc.*, 108, 832 (1986).


N-arylimidazoles of formula I (compounds wherein r = o) can be prepared by the following methods, it being understood by one skilled in the art that certain manipulations, protecting and deprotecting steps, and other synthetic procedures disclosed above may be necessary to produce compounds with the desired combinations of R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>13</sup>.

As shown in Scheme 4, equation a) the reaction of aniline derivative (34) with imidate ester (12) to form the substituted amidine (35) provides material which can be cyclized with dihydroxyacetone to form structure (36). Subsequent elaboration into (I) provides the N-arylimidazole compounds of the invention.


Alternatively as shown by equation b) the Marckwald procedure, described by Marckwald et al., *Ber.*, 22, 568, 1353 (1889); *Ber.*, 25, 2354 (1892) can be used to form a 2-mercaptoimidazole (38) from aniline derivative (34) via isothiocyanate (37). Desulfurization of (38) with dilute nitric acid followed by anion formation at the 2-position of the imidazole (39) and reaction with R<sup>6</sup>X where X is Cl, Br, I, allows the formation of (40) which can be subsequently elaborated to I.

A variation of Marckwald's process as shown in equation c) using an  $\alpha$ -aminoketone (41) and isothiocyanate (37) can also be employed, see Norris and McKee, *J. Amer. Chem. Soc.*, 77, 1056 (1955) can also be employed. Intermediate (42) can be converted to (I) by known sequences. The general procedure of Carboni et al., *J. Amer. Chem. Soc.*, 89, 2626 (1967) (illustrated by equation d)) can also be used to prepare N-aryl substituted imidazoles from appropriate haloaromatic compounds (43; X = F, Cl, Br) and imidazoles (1):

## Scheme 4




Scheme 4 (continued)



In various synthetic routes  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  do not necessarily remain the same from the starting  
55 compound to the final products, but are often manipulated through known reactions in the intermediate  
steps as shown in Schemes 5-22. All of the transformations shown in Schemes 5-10 and 12 can also be  
carried out on the terminal aromatic ring (i.e., biphenyl ring).

Scheme 5



As shown in Scheme 5, compounds where  $R^1$  is a sulfonic acid group may be prepared by oxidation of the corresponding thiol (45). Thus, an N-benzylimidazole derivative bearing a thiol group may be converted into a sulfonic acid (46) by the action of hydrogen peroxide, peroxyacids such as metachloroperoxybenzoic acid, potassium permanganate or by a variety of other oxidizing agents, E. E. Reid, Organic Chemistry of Bivalent Sulfur, 1, Chemical Publishing Co., New York, 120-121 (1958).

Aromatic hydroxy or thiol groups are obtained from deprotection of the corresponding alkyl ether or thioethers. Thus, for example, a methyl ether or a methyl thioether derivative (44) of an N-benzylimidazole containing one or more aromatic rings may be converted into the free phenol or thiophenol (45) by the action of boron tribromide methyl sulfide, P. G. Willard and C. F. Fryhle, Tet. Lett., 21, 3731 (1980); trimethylsilyl iodide, M. E. Jung and M. A. Lyster, J. Org. Chem., 42, 3761 (1977); KSEt and derivatives thereof, G. I. Feutrell, R. N. Mirrington, Tet. Lett., 1327, (1970), and a variety of other reagents.

Alternatively, N-benzylimidazoles may be sulfonated by stirring with  $H_2SO_4$  at a variety of different concentrations or with other sulfonating agents such as chlorosulfonic acid or sulfur trioxide with or without complexing agents such as dioxane or pyridine at temperatures from 0 to 200 °C with or without solvent, K. LeRoi Nelson in Friedel-Crafts and Related Reactions, III part 2, G. A. Olah, ed., Interscience Publ., 1355 (1964).

The synthesis of compounds where  $R^1$  is a sulfate, phosphate or phosphonic acid are depicted in Scheme 6:

## Scheme 6

5

10

15

20

25

30


35

40


45

50

55



## Scheme 6 (continued)



N-Benzylimidazoles containing a phenolic hydroxyl group (47) may be readily converted into the corresponding sulfate (48) or phosphate (49). As shown in equation a), reaction of the phenol with a sulfur trioxide-amine complex will give the corresponding sulfate (48), E. E. Gilbert, *Sulfonation and Related Reactions*, Interscience, New York, chapter 6 (1965). Reaction of the phenol (47) with phosphorus pentachloride followed by hydrolysis will give the corresponding phosphate (49), G. M. Kosolapoff, *Organophosphorus Compounds*, John Wiley, New York, 235 (1950).




As shown in equation b) N-benzylimidazoles may be converted into the corresponding phosphonic acids by reaction with phosphorus trichloride ( $\text{PCl}_3$ ) and aluminum chloride ( $\text{AlCl}_3$ ) in an inert solvent for 0.5-96 hours from temperatures of  $25^\circ\text{C}$  to the reflux temperatures of the solvent. Appropriate workup followed by reaction with chlorine ( $\text{Cl}_2$ ) and subsequent hydrolysis of the tetrachloride (51) gives the phosphonic acid derivative (52), G. M. Kosolapoff in *Org. Reactions*, 6, R. Adams, editor, John Wiley and Sons, New York, 297 (1951). Another more direct route involves reaction of the N-benzylimidazole with  $\text{PSCl}_3$  and  $\text{AlCl}_3$  followed by hydrolysis, R. S. Edmunson in *Comprehensive Organic Chemistry*, Vol. 2, D. Barton and W. D. Ollis editors, Pergamon Press, New York, 1285 (1979).

Alternatively, equation c) illustrates that aryl phosphonic acids (52) may be formed from reaction of the corresponding diazonium salt (53) with  $\text{PCl}_3$  in the presence of  $\text{Cu(I)}$  followed by hydrolysis with water (ibid, p. 1286).

As shown in equation d), the aryl halides (55) may be photolyzed in the presence of phosphite esters to give phosphonate esters (56), R. Kluger, J. L. W. Chan, *J. Am. Chem. Soc.*, 95, 2362, (1973). These same aryl halides also react with phosphite esters in the presence of nickel or palladium salts to give phosphonate esters, P. Tavs, *Chem. Ber.*, 103, 2428 (1970), which can be subsequently converted to phosphonic acids (52) by procedures known to one skilled in the art.


N-Benzylimidazoles containing an aldehyde or ketone (57) may be reacted with a phosphorus trihalide followed by water hydrolysis to give  $\alpha$ -hydroxyphosphonic acid derivatives, G.M. Kosolapoff, *op. cit.*, 304, as shown in Scheme 7.

Scheme 7




Compounds where  $\text{R}^1$  is  $-\text{CONHOR}^{12}$  may be prepared as shown in Scheme 8, by the treatment of a carboxylic acid (10) with 1-4 equivalents of thionyl chloride for 1-10 hours. This reaction can be run without solvent or in a nonreactive solvent such as benzene or chloroform at temperatures of  $25$ - $65^\circ\text{C}$ . The intermediate acid chloride is then treated with 2-10 equivalents of the appropriate amine derivative,  $\text{H}_2\text{N}-\text{OR}^{12}$ , for 2-18 hours at temperatures of  $25$ - $80^\circ\text{C}$  in a polar aprotic solvent such as tetrahydrofuran or dimethylsulfoxide to give the hydroxamic acid (59).

Scheme 8



15 Alternatively, the carboxylic acid (10) can be converted to the hydroxamic acid (59) according to the procedure in *J. Med. Chem.*, 28, 1158 (1985) by employing dicyclohexylcarbodiimide, 1-hydroxybenzotriazole, and  $H_2NOR^{12}$  or according to the procedure described in *Synthesis*, 929 (1985) employing the Vilsmeier reagent and  $H_2NOR^{12}$ .

Scheme 9




Aniline intermediates (63) are disclosed in U.S. Patent No. 4,355,040 and may be obtained from the corresponding nitro compound precursor by reduction. A variety of reduction procedures may be used such

as iron/acetic acid, D. C. Owsley, J. J. Bloomfield, *Synthesis*, 118, (1977), stannous chloride, F. D. Bellamy, *Tet. Lett.*, 839, (1984) or careful hydro-genation over a metal catalyst such as palladium.

As shown in Scheme 9, aniline intermediates of N-benzylimidazoles may also be prepared from the corresponding carboxylic acid (10) or acid chloride via a Curtius rearrangement of an intermediate acyl azide (60). More modern methods include using diphenylphosphoryl azide as a source of azide, T. Shioiri, K. Ninomiya, S. Yamada, *J. Am. Chem. Soc.*, 94, 6203 (1972), and trapping the intermediate isocyanate (61) produced by the Curtius rearrangement with 2-trimethylsilylethanol and cleaving the resultant carbamate (62) with fluoride to liberate the amine (63), T. L. Capson and C. D. Poulter, *Tet. Lett.*, 25, 3515 (1984). Classical procedures familiar to one skilled in the art may also be employed.

Compounds where R<sup>1</sup> is -SO<sub>2</sub>NH<sub>2</sub> may be made as shown in Scheme 10:


Scheme 10



Sulfonamide compounds (65) may be made by reacting an arylsulfonyl chloride (64) with ammonia, or its equivalent. Unsubstituted arylsulfonamides are made by reaction with ammonia in aqueous solution or in an inert organic solvent, F. H. Bergheim and W. Braker, *J. Am. Chem. Soc.*, 66, 1459 (1944), or with dry powdered ammonium carbonate, E. H. Huntress and J. S. Autenrieth, *J. Am. Chem. Soc.*, 63, 3446 (1941); E. H. Huntress and F. H. Carten, *J. Am. Chem. Soc.*, 62, 511 (1940).

The sulfonyl chloride precursor may be prepared by chlorosulfonation with chlorosulfonic acid on the aromatic ring directly, E. H. Huntress and F. H. Carten, *ibid.*; E. E. Gilbert, *op. cit.*, 84, or by reacting the corresponding aromatic diazonium chloride salt (53) with sulfur dioxide in the presence of a copper catalyst, H. Meerwein, et al., *J. Prakt. Chem.*, [ii], 152, 251 (1939), or by reacting the aromatic sulfonic acid (46) with PCl<sub>5</sub> or POCl<sub>3</sub>, C. M. Suter, *The Organic Chemistry of Sulfur*, John Wiley, 459 (1948).

Linked ester compounds of formula (I) where R<sup>1</sup> is



can be made by procedures well known in penicillin and cephalosporin chemistry. The purpose is to provide materials which are more lipophilic and which will be useful orally by rapid transit from the gut into the bloodstream, and which will then cleave at a sufficiently rapid rate to provide therapeutically useful concentrations of the active carboxylic acid form. The following review articles and references cited therein discuss this concept and the chemistry involved in preparing such compounds V. J. Stella, et al., *Drugs*, 29, 455-473 (1985); H. Ferres, *Drugs of Today*, 19 (9), 499-538 (1983); A. A. Sirkula, *Ann. Repts. Med. Chem.*, 10, 306-315 (1975).

Experimental procedures which are applicable to the preparation of chemically stable linked esters are illustrated by equations a-e of Scheme 11.



Scheme 11



10


66

G. Francheschi et al., J. Antibiotics, 36, (7),  
938-941 (1983).




67

J. Budavin, U.S. Patent 4,440,942




68

B. Daehne et al., G.B. Patent 1,290,787



69


Perres, Chem. Ind., 435-440 (1980)



70


Clayton et al., Antimicrob. Agents Chemotherapy,  
5, (6), 670-671 (1974)

In equations a-e: R=




Compounds of Formula I where R<sup>1</sup> is -C(CF<sub>3</sub>)<sub>2</sub>OH may be prepared as shown in Scheme 12.

Scheme 12



Hexafluoroisopropanol compounds (72) may be prepared by treatment of arylsilane (71) with 1-5 equivalents of hexafluoroacetone in a solvent such as methylene chloride at temperatures ranging from about -50° to 25°C for a period of 2-10 hours. The requisite arylsilane (71) can be prepared using methods known to one skilled in the art such as the procedures described in Chapter 10 of Butterworth's "Silicon in Organic Chemistry".

Scheme 13



As shown in Scheme 13, compound (73) in which X = -NHCO and R<sup>13</sup> = -COOH may be easily prepared, for example, by reacting aniline precursor (63) with a phthalic anhydride derivative in an appropriate solvent such as benzene, chloroform, ethyl acetate, etc. Often the carboxylic acid product will precipitate from solution with the reactants remaining behind, M.L. Sherrill, F.L. Schaeffer, E.P. Shoyer, J. Am. Chem. Soc., 50, 474 (1928).

When  $R^{13} = \text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{CF}_3$  or tetrazolyl (or a variety of other carboxylic acid equivalents), compound (73) may be obtained by reacting aniline (63) with the requisite acid chloride by either a Schotten-Baumann procedure, or simply stirring in a solvent such as methylene chloride in the presence of a base such as sodium bicarbonate, pyridine, or triethylamine.

5 Likewise, aniline (63) may be coupled with an appropriate carboxylic acid via a variety of amide or peptide bond forming reactions such as DCC coupling, azide coupling, mixed anhydride synthesis, or any other coupling procedure familiar to one skilled in the art.

Aniline derivatives (63) will undergo reductive amination with aldehydes and ketones to form secondary amines (74). Thus the aniline is first stirred with the carbonyl compound in the presence of a dehydration catalyst such as molecular sieves or p-toluenesulfonic acid. Afterwards the resultant imine is reduced to the amine with a borohydride reducing agent such as sodium cyanoborohydride or sodium borohydride. Standard catalytic hydrogenation reagents such as hydrogen and palladium/carbon can also be employed.

Alternatively, aniline (63) may be monoalkylated by reaction with ethyl formate followed by reduction with, for example, lithium aluminum hydride to produce the N-methyl derivative (74). Anilines (74) may in turn be reacted with carboxylic acid anhydrides and acid chlorides or carboxylic acids by any of the coupling procedures described previously to yield (73) where  $X = -\text{N}(\text{CH}_3)\text{CO}-$ .

Aniline (63) or (74) or other intermediate anilines where the amino group may be located on another aromatic ring for example, also react with other anhydrides to make amide-carboxylic acid derivatives of formula (75). Thus, for example, maleic anhydride, 2,3-naphthalenedicarboxylic acid anhydride, and diphenic anhydride are reacted in a similar fashion to phthalic anhydride with aniline (63) or (74) to yield carboxylic acids (76), (77), and (78), respectively.

Phthalimide derivatives of aniline (63) may be made by a variety of methods, preferably by stirring aniline (63) with phthalic anhydride in acetic acid at a temperature between 20 °C and reflux, G. Wanag, A. Veinbergs, *Ber.*, 75, 1558 (1942), or by stirring (63) with phthaloyl chloride, a base such as triethylamine, and an inert solvent. Aniline (63) may be converted into its trifluoromethanesulfonamide derivative or its trifluoroacetamido derivative preferably by reacting it with triflic anhydride or trifluoroacetic anhydride and a base such as triethylamine in an inert solvent such as methylene chloride at -78 °C followed by warming to room temperature.

Compounds of structure (I) where X is a carbon-carbon linkage which are depicted as (80) can be made as shown in Scheme 14.

35

40

45

50

55

## Scheme 14

5

10

15

20

25

30


35

40

45


50

55





Scheme 14 (Cont'd.)



Equation a) illustrates that the biphenyl compounds (80) can be prepared by alkylation of imidazole (1) with the appropriate halomethylbiphenyl compound (79) by the general procedure described in Scheme 1.

The requisite halomethylbiphenyl intermediates (79) are prepared by Ullman Coupling of (81) and (82) as described in "Organic Reactions", 2, 6 (1944) to provide intermediates (83), which are in turn halogenated. Halogenation can be accomplished by refluxing (83) in an inert solvent such as carbon tetrachloride for 1-6 hours in the presence of a N-halosuccinimide and an initiator such as azobisisobutyronitrile (equation b).

As shown in equation c), derivatives of intermediate (83) in which  $R^{13}$  is at the 2' position (83a) can also be prepared by the method described in J. Org. Chem., 41, 1320 (1976), that is Diels-Alder addition of a 1,3-butadiene to a styrene (84) followed by aromatization of intermediate (85).

Alternatively, the substituted biphenyl precursors (83; where  $R^{13} = COOH$ ) and their esters (89) can be prepared as illustrated in equation d), which involves oxazoline compounds as key intermediates, A. I. Meyers and E. D. Mihelich, J. Am. Chem. Soc., 97, 7383 (1975).

The substituted biphenyl tetrazoles (83; where




can be prepared from the nitrile precursors ( $R^{13} = CN$ ) by the methods described in Scheme 1, equation c) and Scheme 15, equation c).

However, a preferred method for preparing tetrazoles is described in Scheme 15, equations a) and b). Compounds (90) may be prepared by the 1,3-dipolar cycloaddition of trialkyltin or triphenyltin azides to the appropriately substituted nitrile (83) as in equation a). Alkyl is defined as normal alkyl of 1-6 carbon atoms and cyclohexyl. An example of this technique is described by S. Kozima, et al., J. Organometallic Chemistry, 337 (1971). The required trialkyl or triaryl tin azides are made from the requisite commercial trialkyl or triaryl tin chloride and sodium azide. The trialkyl or triaryl tin group is removed via acidic or basic hydrolysis and the tetrazole can be protected with the trityl group by reaction with trityl chloride and triethylamine to give (91). Bromination as previously described herein with N-bromosuccinimide and

dibenzoylperoxide affords compound (92). Alkylation of (1) with the appropriately substituted benzyl halide using conditions previously described followed by deprotection of the trityl group via hydrolysis affords (80;  $R^{13}$  = tetrazole). Other protecting groups such as p-nitrobenzyl and 1-ethoxyethyl can be used instead of the trityl group to protect the tetrazole moiety. These groups as well as the trityl group can be introduced and removed by procedures described in Greene, Protective Groups in Organic Synthesis, Wiley-Inter-science, (1980).

## Scheme 15



10




45

50

55


Scheme 15 (continued)

Compounds of structure 93-95 where X is an -O-, -S-, or




linkage can be prepared as shown in Scheme 16 by alkylation of imidazole (1) with the appropriate benzyl halide (96).

Scheme 16



## Scheme 16 (continued)




The halomethyldiphenyl ether (109) employed as an alkylating agent in the present invention is prepared as shown in equation b). An Ullman ether condensation of the phenol (97) and a halobenzoic acid as described in *Russian Chemical Reviews*, 43, 679 (1974) provides the intermediate acid (101). The conversion of (101) into (109) is accomplished by esterification with diazomethane to afford (105) followed by halogenation employing the procedure used in the preparation of (79). The diphenylsulfide (110) and the diphenylamine (111) can be prepared from the appropriate thiophenol (98) or aniline (99) by this procedure.

The tertiary diphenylamine (112) can be prepared from the secondary aniline (100) by the above procedure. Alternatively (107) can be alkylated by one of the following procedures: 1) direct alkylation of (107) with R<sup>26</sup>L where L is a leaving group such as a halogen or tosylate employing phase-transfer conditions and ultrasound as described in *Tetrahedron Letters*, 24, 5907 (1983), 2) treatment of (107) with 1-1.5 equivalents of an appropriate aldehyde and 0.5-5.0 equivalents of sodium cyanoborohydride in a solvent such as methanol at 25°C at a pH of 3-6 for 1-24 hours, or 3) reductive amination of (107) employing an appropriate carboxylic acid and sodium borohydride as described in *J. Am. Chem. Soc.*, 96, 7812 (1974).

The tertiary amine (108) is then halogenated by the procedure previously described to give (112).


## Scheme 17




Compounds of structure (73) where X is -CO- are prepared as shown in Scheme 17 by alkylation of imidazole (1) with the requisite benzoylbenzyl halides. For example, esters (113) where  $R^{13}$  is 2-CO<sub>2</sub>CH<sub>3</sub> are prepared by alkylation of imidazole (1) with carbomethoxybenzoyl benzyl halide (114). Ester (113) may be hydrolyzed to the corresponding carboxylic acid (116) by a variety of methods including hydrolysis with a base such as sodium hydroxide or potassium hydroxide in an alcoholic aqueous solvent such as methanol/H<sub>2</sub>O at a temperature from 20 °C to the reflux temperature of the solvent.

Carboalkoxybenzoylbenzyl halides (114) are prepared by benzylic halogenation of the corresponding toluoylbenzene precursor by a variety of methods previously described herein. For example, methyl 2-(4-methylbenzoyl)benzoate (115) can be refluxed for 2-48 hours with N-bromosuccinimide, benzoyl peroxide and carbon tetrachloride to effect benzylic bromination.

Scheme 18



As shown in Scheme 18 the toluoyl ketones (73; where X=CO) may be further transformed into a variety of ketone derivatives including compounds where X is



5

Reaction of ketone (73a) with a hydroxylamine or an appropriately substituted hydrazine will give the requisite oximes (117) and hydrazones (118). Reaction with alcohols in the presence of an acidic catalyst with removal of water will give ketals (119). Reduction, with lithium aluminum hydride, a metal borohydride, zinc/acetic acid or catalytic hydrogenation will give the corresponding alcohol (120) or fully reduced methylene compound (121). These alcohols may be acylated by a variety of anhydrides or acid halides in the presence of a base with or without solvent to give the corresponding esters (122). The alcohols (120) may be converted into their corresponding ethers (123) by reaction of the metal alkoxide with an alkyl halide, mesylate or tosylate in the appropriate solvent or by treatment with a mineral acid in an alcoholic solvent, or by reaction of the alcohol with diazomethane G. Hilgetag and A. Martini, "Preparative Organic Chemistry", John Wiley, New York, 355-368 (1972).

15

Compounds of formula (I) where X is -OCH<sub>2</sub>-, -SCH<sub>2</sub>-, and -NHCH<sub>2</sub>- are prepared as shown in Scheme 19.

20

25

30

35

40


45

50

55



Scheme 19



As illustrated in Scheme 19, equation a, hydrolysis of benzyl ether (124) or methyl ether (125) affords hydroxy compound (126) which can be alkylated with the appropriate benzyl halide to give (127). In the case of the methyl ethers (125), the hydrolysis step can be effected by heating the ether at temperatures of 50°-150°C for 1-10 hours in 20-60% hydrobromic acid, or heating at 50°-90°C in acetonitrile with 1-5 equivalents of trimethylsilyl iodide for 10-50 hours followed by treatment with water. Hydrolysis can also be carried out by treatment with 1-2 equivalents of boron tribromide in methylene chloride at 10°-30°C for 1-10 hours followed by treatment with water, or by treatment with an acid such as aluminum chloride and 3-30 equivalents of a sulfur-containing compound such as thiophenol, ethanedithiol, or dimethyl disulfide in methylene chloride at 0-30°C for 1-20 hours followed by treatment with water. For compound (124),

hydrolysis can be accomplished by refluxing in trifluoroacetic acid for 0.2-1 hours or by catalytic hydrogenolysis in the presence of a suitable catalyst such as 10% palladium on carbon. Deprotonation of (126) with a base, such as sodium methoxide, sodium hydride or the like in a solvent such as dimethylformamide or dimethylsulfoxide at room temperature followed by alkylation with an appropriate benzyl halide at 25 °C for 2-20 hours affords ethers of formula (127), as shown in equation a.

The sulfide (129) can be prepared from the thiophenol (45) by the procedure described above to prepare the ether (127) from the phenol (126). The thiophenol (45) can be prepared for example by treatment of the benzylsulfide (128) with sodium in liquid ammonia.


The amine (130) can be prepared as shown in equation c, from the aniline (63), itself available from reduction of the corresponding p-nitro compound (3a) which has previously been described. The reductive amination can be carried out by the same procedure as described in Scheme 13 for the preparation of compound (74).

Compounds of Formula (1) where the X linkage is -CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, and



are prepared as shown in Scheme 20.

Scheme 20



The cis or trans stilbene (132) can be obtained by employing a Wittig reaction between the aldehyde (57) and the phosphorane (131).


The stilbene (132) can readily be converted to the saturated derivative (133) for example by catalytic hydrogenation employing a heterogeneous catalyst such as palladium/carbon or platinum/carbon or alter-

natively with a homogeneous catalyst such as trisphenylphosphine rhodium chloride. The reduction is performed in a solvent such as benzene, tetrahydrofuran or ethanol at 25°C under 1-3 atmospheres of hydrogen for 1-24 hours.

The cyclopropane (134) can be prepared by treating the stilbene (132) with the Simmons-Smith reagent as described in *J. Am. Chem. Soc.*, **81**, 4256 (1959), or by treating (132) with methylene diiodide and copper powder as described in *J. Am. Chem. Soc.*, **101**, 2139 (1979), or by treatment with the iron-containing methylene-transfer reagent described in *J. Am. Chem. Soc.*, **101**, 6473 (1979).

The preparation of compounds of formula (I) where X is -CF<sub>2</sub>CH<sub>2</sub>-, -CF=CH-, -CH=CF-, -CF=CF- and -CF<sub>2</sub>CF<sub>2</sub>- are depicted in Scheme 21.

Scheme 21




Vinylene fluorides (137) and (140) can be prepared by reaction of SF<sub>4</sub> or Et<sub>2</sub>NSF<sub>3</sub> (DAST) with the appropriate ketone (135) or (138) in which Ar bears a methyl group convertible to a benzylic halide suitable for attachment to an imidazole nitrogen, and Ar' bears a cyano, nitro, ester, or other suitable group which

can be subsequently converted to  $\text{CO}_2\text{H}$ ,  $\text{NHSO}_2\text{CF}_3$ , etc. The initially formed difluoroethylene (136) and (139) can be formed in a non-polar solvent such as methylene chloride and subsequently converted to the vinylene fluoride by means of alumina, or converted directly into the unsaturated fluoride by running the reaction in a polar solvent such as tetrahydrofuran, diglyme or N-methylpyrrolidone in the presence of mineral acid. [Equations a and b]. Experimental details of such procedures are found in D.R. Strobach and G.A. Boswell, *J. Org. Chem.*, **36**, 818 (1971); G.A. Boswell, U.S. Patents 3,413,321 (1968) and 4,212,515 (1980).

As shown in equation c an appropriate benzoin (141) may be similarly converted to the corresponding 1,2-difluorostilbene (143). Likewise as shown in equation d an appropriate benzil (144) can be converted to a tetrafluorodiarylethylene (145) using DAST or  $\text{SF}_4$ . Experimental details are described in M.E. Christy, et al., *J. Med. Chem.*, **20**, (3), 421-430, (1977).


Compounds of formula 1 where X =



-CH<sub>2</sub>O-, -CH<sub>2</sub>S-, -CH<sub>2</sub>NH-, can be made as shown in Scheme 22.

Scheme 22

5



As previously described, acid (10) can be made by alkylating the appropriate imidazole with methyl 4-chloromethylbenzoate in the presence of a base such as potassium carbonate in a polar solvent such as dimethylformamide followed by hydrolysis of the resulting ester. Compound (10) can be converted to (148) by reaction with the requisite amine (146) (R<sup>13</sup> may need to be protected and subsequently deprotected) and dicyclohexyl carbodiimide (DCC) in methylene chloride [J. R. Beek, et al., J. Am. Chem. Soc., 90, 4706 (1968)] or by reaction with tosyl chloride in pyridine [J. H. Brewster and C. J. Ciotti, Jr., J. Am. Chem. Soc., 77, 6214 (1955)]. Yet another process involves conversion of carboxylic acid (10) to its acid chloride with, for example, thionyl chloride followed by reaction with the amine in aqueous base (Schotten-Baumann conditions) or in an organic solvent in the presence of an acid scavenger such as NaHCO<sub>3</sub>, pyridine or triethylamine, or by other procedures known to form an amide bond between an aromatic acid and an amine.

The compounds where  $X = -CH_2O-$ ,  $-CH_2S-$ , and  $-CH_2NH_2-$  can be made as shown in pathway b. The ester (149) is reduced with a reducing agent such as lithium aluminum hydride in an inert solvent to form the alcohol (150) which can then be reacted with tosyl chloride in pyridine to form tosylate (151), which is in turn reacted in the presence of base with a corresponding phenol (152) thiophenol (153), or aniline (146; where  $R^{23} = H$ ) to form compounds (154), (155) or (156). Again this may require that  $R^{13}$  be protected with a suitable protecting group, however modifications necessary because of specific functional groups are understood to be incorporated by one skilled in the art of organic synthesis.

Alternatively, the alcohol (150) can be converted to the corresponding halide with  $SOCl_2$ ,  $(COCl)_2$ , etc, and the resulting halide can then be reacted with a phenol, thiophenol or aniline in the presence of base to form the desired compound, where  $X$  is  $-CH_2O-$ ,  $-CH_2S-$ ,  $-CH_2NH-$  respectively.

Scheme 23

15

20

25

30


35

40

45

50

55



Compounds of Formula (I) where  $X = -SO_2NR^{23}-$  and  $-NR^{23}SO_2-$  may be prepared as shown in Scheme 23. As shown in equation a, sulfonylchloride derivative (157) can be reacted with aniline derivative (158) in a

solvent in the presence of an acid scavenger such as sodium bicarbonate, triethylamine or pyridine or under Schotten-Baumann like conditions to give (159). Sulfonylchloride derivative (157) can be obtained by sulfonation of the corresponding benzyl derivative as described earlier, followed by reaction with  $\text{PCl}_5$  or  $\text{POCl}_3$ . Likewise, aniline (74) may be reacted in the same manner as described above with sulfonylchloride derivative (160) to give (161).

Scheme 24 shows the preparation of furan analogs of the biphenyl compounds (80). Thus,  $\alpha$ -ketoester (162), W. Wierenga and H. I. Skulnick, *J. Org. Chem.*, **44**, 310 (1979), or the corresponding nitrile ( $\text{E} = \text{CN}$ ) can be easily alkylated via standard procedures already mentioned by an alkyl bromide derivative to give (163). The alkene moiety of (163) can be subsequently cleaved by oxidation, for example, with osmium tetroxide, Fieser and Fieser, V.1, p. 812 (Lemieux-Johnson oxidation) to yield dicarbonyl-containing compound (164). Cyclization in mineral acids, acidic ion-exchange resin,  $\text{POCl}_3$ /pyridine, or trifluoroacetic anhydride with a catalytic amount of trifluoroacetic acid yields furan (165;  $\text{Z} = \text{O}$ ). Reaction of (164) with  $\text{P}_4\text{S}_{10}$ , for example, will yield the corresponding thiophene (165;  $\text{Z} = \text{S}$ ). Reaction of (164) with an amine in refluxing benzene, with azeotropic removal of water or by using molecular sieves to absorb the water will yield the corresponding pyrrole (165;  $\text{Z} = \text{NR}^{11}$ ). Compounds (166) may be prepared from (165) by standard procedures already described.

20

25

30

35


40

45

50

55

Scheme 24




Compounds wherein a methylene group is inserted between the terminal aromatic ring and the acidic functionality may be prepared as shown in Scheme 25, equation a). Thus reduction of ester (167) with, for example, lithium aluminum hydride, gives alcohol (168). Conversion of (168) to the chloride (169) via thionyl chloride followed by reaction with cyanide anion as previously described yields nitrile (170). Compound (170) may be hydrolyzed to carboxylic acid (171) by methods already described or reacted with a



hydrazoic acid equivalent to produce tetrazole (172).

Compounds wherein  $R^{13}$  is a trifluoromethylsulfonyl hydrazide acidic functional group were prepared by the procedure described in equation b). That is, conversion of ester (167) to the hydrazide (173) by standard hydrazinolysis followed by reaction with triflic anhydride affords hydrazides (174).

Scheme 25



The syntheses of compounds wherein R<sup>13</sup> is substituted and unsubstituted 1,2,3-triazoles are described in Scheme 26. Thus reduction of ester (175) with a reducing agent such as lithium aluminum hydride or diisobutylaluminum hydride gives alcohol (176). Oxidation with MnO<sub>2</sub> or pyridinium chlorochromate converts (176) into aldehyde (177). Nitroethylene derivative (178) is prepared by condensation of aldehyde (177) with nitromethane in the presence of a catalyst, R. M. Letcher and M. P. Sammes, *J. Chem. Ed.*, **62**, 262 (1985). Reaction of (178) with sodium azide produces the 1,2,3-triazole (179), (N. S. Zefirov, et al., *J. Chem. Soc. Chem. Comm.*, 1001 (1971)) which may be transformed via procedures already described into product (180).

Aldehyde (177) can also be converted into substituted 1,2,3-triazoles (183) via the sulfone (181), G. Beck, D. Günther, *Chem. Ber.*, **106**, 2758 (1973), followed by reaction with sodium azide to give the 1,2,3-triazole (182). Subsequent standard manipulations lead to 1,2,3-triazoles (183) where E = CN and CO<sub>2</sub>R<sup>11</sup>. The nitrotriazole (183; E = NO<sub>2</sub>) may be synthesized from the unprotected triazole (179; P = H) via nitration, R. Hüttel, et al., *Chem. Ber.*, **88**, 1586 (1955), C. L. Habraken and P. Cohen-Fernandes *J. Chem. Soc.*, 37 (1972), or from bromonitroethylene derivative (184), G. Kh. Khisamutdinov, et al., *Zh. Org. Khim.*, **11**, 2445 (1975), by reaction with sodium azide.

A variety of protecting groups may be used in the manipulation of the above triazoles, amongst which is the trityl group. This group may be easily attached by reaction of the triazole with triphenylmethyl bromide or chloride in an inert solvent such as methylene chloride in the presence of an acid scavenger such as triethyl amine. The trityl group may be later removed by stirring or refluxing in an acidic medium such as trifluoroacetic acid/water, HCl in methylene chloride, or acetic acid/water. The trityl group may also be hydrogenolyzed using a noble metal catalyst such as palladium and hydrogen.

25

30

35


40

45

50

55

Scheme 26




P=protecting group

The synthesis of trifluoromethyl-1,2,4-triazoles (190) is depicted in Scheme 27. Acid chloride (186) is converted to amide (187) using standard procedures familiar to one skilled in the art. A preferred protecting group is the 2-propionitrile group (P=CH<sub>2</sub>CH<sub>2</sub>CN). Thus (187; P=CH<sub>2</sub>CH<sub>2</sub>CN) can be synthesized from (186) and β-aminopropionitrile under Schotten-Baumann like conditions, using aqueous base in an organic solvent to help solubilize (186) and (187). Amide (187) is converted to amidrazone (188) by reaction with

PCl<sub>5</sub> or phosgene to make an iminoyl chloride which then in turn is reacted with excess hydrazine. Amidrazone (188) is cyclized to the trifluoromethyl-1,2,4-triazole (189) with trifluoroacetic anhydride and then converted to 190 via bromination, alkylation and deprotection as previously described.

Scheme 27



45 Pertinent R<sup>6</sup> groups may be variously introduced by many procedures including those described in Scheme 28 which describes imidazole construction.

The R<sup>6</sup> groups so introduced may stand unchanged or may be further elaborated if appropriately functionalized, according to methods familiar to those skilled in the art such as are illustrated in Scheme 28.

## Scheme 28

5

10

15


20

25

30

35

40




The 2-alkenylimidazoles (201) can be prepared by bromination of the 2-alkylimidazoles (199) followed by elimination of hydrogen bromide. The bromization is preferably accomplished by UV-irradiation for 1-4 hours of imadoyole (199) and N-bromosuccinimide, in an inert solvent, such as carbon tetrachloride at 25°C. Treatment of the intermediate bromide (200) with a base, such as DBU, triethylamine, or potassium t-butoxide, affords the trans 2-alkenylimidazoles (201). Cis alkenyl derivatives (203) are prepared from the trans alkenyl compounds by treatment with osmium tetroxide and sodium periodate to afford aldehydes (202) followed by Wittig reaction.

50

55

## Scheme 29



R = alkyl, cycloalkyl

45


Alternatively, R<sup>6</sup> groups may be introduced by metallation of a protected imidazole or protected 2-methylimidazole followed by addition of an appropriate electrophile as illustrated in Scheme 30, equations a) and b). The products (alcohols, esters, halides, aldehydes, alkyls) are suitable for further elaboration by methods familiar to those skilled in the art. Metallation of imidazoles is described in K.L. Kirk, *J. Org. Chem.*, **43**, 4381 (1978); R.J. Sundberg, *J. Het. Chem.*, **14**, 517 (1977); J.V. Hay et al., *J. Org. Chem.*, **38**, 4379 (1973); B. Iddon, *Heterocycles*, **23**, 417 (1985).

50

Condensation of 2-methylimidazole and appropriate electrophiles (equation b) with catalytic acid or base as described in A.R. Katritzky (Ed.), "Comprehensive Heterocyclic Chemistry", Vol. 5, p. 431, Pergamon Press, N.Y., 1984 affords products wherein R<sub>5</sub> is alkenyl which are suitable for further elaboration.


55

## Scheme 30




Various 2-substituted imidazoles can be prepared by reaction of a protected 2-trimethylsilylimidazole with a suitable electrophile by the method described by F.H. Pinkerton and S.F. Thames, *J. Het. Chem.*, **9**, 67 (1972), which can be further elaborated as desired. Alternatively,  $R^6$  may also be introduced by nickel catalyzed cross-coupling of Grignard reagents with 2-(methylthio)imidazoles (Scheme 31) as described by E. Wenkert and T.W. Ferreira, *J. Chem. Soc., Chem. Commun.*, 840, (1982); E. Wenkert et al., *J. Chem. Soc., Chem. Commun.*, 637, (1979); and H. Sugimura and H. Takei, *Bull. Chem. Soc. Japan*, **58**, 664 (1985). The 2-(methylthio)imidazoles can be produced by the procedure described in German Patent No. 2,618,370 and the references cited therein.

## Scheme 31




As shown in Schemes 32-35, elaboration of  $R^8$  can be accomplished by procedures described in Schemes 3, 28 and 30b and by chain extension reactions familiar to those skilled in the art in which  $R^8$  bears a reactive terminal functional group, e.g. -OH, halogen, -CHO, -CO<sub>2</sub>R, -CO<sub>2</sub>H, -CH=CH<sub>2</sub>, -NH<sub>2</sub>, -NO<sub>2</sub>, -CN,



etc., or by degradation reactions such as conversion of an ester to an acid or an alkene to an aldehyde.

Specifically, the hydroxymethyl group can be activated for the displacement reaction by reacting with thionyl chloride, POCl<sub>3</sub> or with carbon tetrachloride/triphenylphosphine to form a corresponding chloro derivative. By a similar reaction bromo and iodo derivatives can be obtained. The hydroxymethyl group can also be activated by forming the corresponding p-toluenesulfonate, methanesulfonate and trifluoromethane sulfonate derivatives. The hydroxyl group can be converted to its corresponding fluoro compound by various fluorinating agents such as DAST as shown in Scheme 32.

Scheme 32






Also as shown in Scheme 32, the hydroxyl group can be converted to thiolacetic acid derivative (215), J. Y. Gauthier, *Tet. Lett.*, 15 (1986), and to thiol derivative (216) by subsequent hydrolysis.

The hydroxymethyl group on compound (17) can be readily oxidized to an aldehyde group by means of manganese dioxide or ceric ammonium nitrate. The aldehyde group will undergo chain extension reactions such as the Wittig and Wittig-Horner reactions and enter into typical carbon-carbon bond forming reactions with Grignard and lithium reagents as well as with compounds bearing activated methylene groups. Alternatively, the hydroxymethyl group can be oxidized directly to an acid functionality which can in turn be converted to ester and amide derivatives. The esters and amides can be prepared directly from the aldehydes by manganese dioxide oxidation in the presence of sodium cyanide and an alcohol or amine, *J. Am. Chem. Soc.*, 90, 5616 (1968) and *J. Chem. Soc. (C)*, 2355 (1971).

As shown in Scheme 33, the chlorine on compound (25) can be displaced by the anion of dialkyl malonate to give the corresponding malonate derivative (217). The saponification of (217) with NaOH (or KOH) gives the corresponding diacid which can be decarboxylated to give the corresponding propionic acid derivative (218) by heating to 120°C. Alternatively, (218) can be directly obtained by refluxing (217) with a mineral acid such as HCl or sulfuric acid. The free acid (218) can be esterified by heating in a medium of the various alcohols and a catalytic amount of mineral acids such as HCl or sulfuric acid to give the corresponding esters (219). Alternatively the esters can be obtained by reacting the free acid (218) and the corresponding alcohols in the presence of coupling reagents such as DDQ or EEDQ. A similar reaction with various mono-substituted and disubstituted amines produces the corresponding amides (220). A similar reaction with various mercaptans produces the corresponding thioesters.

Scheme 33



As shown in Scheme 34, the chloro group on (25) can be displaced by the sodium salt or potassium salt of the alkyl, aryl or arylalkyl mercaptans to give the corresponding sulfide derivatives (221). The amine derivative (222) can be obtained by treating (25) with ammonia or with the corresponding mono-substituted amines. Alternatively, the chloro group may be displaced by sodium azide to give an azide intermediate  
5 which upon reduction with H<sub>2</sub> over a noble metal catalyst or with a reducing agent such as chromous chloride (W. K. Warburton, J. Chem. Soc., 2651 (1961)) yields (222) where R<sup>10</sup> and R<sup>11</sup> are hydrogen. This amine can be subsequently alkylated with alkyl halides, or reductively alkylated with aldehydes and ketones to give alkyl derivatives of (222). The amines (222) are converted to the corresponding carbamates (224), sulfonamides (225), amides (226) or ureas (227) by standard procedures illustrated in Scheme 34 and  
10 familiar to one skilled in the art. The nitro compound (223) can be obtained by the treatment of (25) with sodium nitrite or potassium nitrite. The nitrate (228) may be synthesized by treatment of (25) with AgNO<sub>3</sub>, A. F. Ferris, et al., J. Am. Chem. Soc., 75, 4078 (1953).

15

20

25

30

35


40

45


50

55

Scheme 34




## Scheme 34 (Cont'd)



20 The reaction between the thiopyridyl ester (229) and a suitable Grignard reagent produces the ketones (230).

## Scheme 35



45 The compounds of this invention and their preparation can be understood further by the following examples, which do not constitute a limitation of the invention. In these examples, unless otherwise indicated, all temperatures are in degrees centigrade and parts and percentages are by weight.

## Example 1


## 50 PART A: Preparation of 2-Butyl-4-chloro-1-(4-cyanobenzyl)-5-hydroxymethylimidazole

To a solution of 2-butyl-4-chloro-5-hydroxymethylimidazole (prepared as described in U.S. 4,355,040; 3.56 g, 40 mmol, 1 eq) in 300 mL methanol was added dropwise a freshly prepared sodium methoxide solution (0.92 g Na, 40 mmol, 1 eq, in 30 mL MeOH). After stirring for 0.5 hours, the methanol was removed in vacuo and the resultant glass was dissolved in 100 mL DMF. To this mixture was added a solution of  $\alpha$ -bromo-p-tolunitrile (8.60 g, 44 mmol, 1.1 eq) in DMF and the entire contents stirred overnight under N<sub>2</sub> at room temperature. The solvent was then removed in vacuo and the residue dissolved in 300 mL ethyl acetate and 300 mL H<sub>2</sub>O. The layers were separated and the aqueous layer was extracted twice with 300

mL portions of ethyl acetate. The organic layers were dried and evaporated and the crude product flash chromatographed over silica gel in 1:1 hexane/ethyl acetate to give 6.83 g of one regioisomer as a white solid; m.p. 92.5-98.0°. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.65 (d, 2H, J = 8Hz); 7.13 (d, 2H, J = 8Hz); 5.30 (s, 2H); 4.46 (s, 2H); 2.49 (t, 2H, J = 7Hz); 1.59 (m, 2H); 1.28 (m, 2H); 0.84 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>16</sub>H<sub>18</sub>N<sub>3</sub>OCl: 303.1138. Found: 303.1124.

Continued elution gave 3.56 g of the second regioisomer as a white solid, listed below as the first entry in Table 1.

The intermediates shown below were prepared or could be prepared in accordance with the procedure described in Example 1, Part A using the appropriately substituted imidazole and benzyl halide as starting material.




R <sup>1</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
4-CN	n-butyl	CH <sub>2</sub> OH	Cl	98.0-100.0
4-NO <sub>2</sub>	n-butyl	Cl	CH <sub>2</sub> OH	56.8- 59.5
4-NO <sub>2</sub>	n-butyl	CH <sub>2</sub> OH	Cl	114.5-116.5
2-CN	n-butyl	Cl	CH <sub>2</sub> OH	93.0- 95.5

#### PART B: Preparation of 2-Butyl-4-chloro-1-(4-cyanobenzyl)-5-cyanomethylimidazole

Thionyl chloride (3.60 mL, 49 mmol, 5 eq) was slowly dripped into a solution of 2-butyl-4-chloro-1-(4-cyanobenzyl)-5-hydroxymethylimidazole (3.0 g, 9.9 mmol, 1 eq) in a minimum of CHCl<sub>3</sub>. The mixture was stirred for 2 hours at room temperature after which the solvent was removed *in vacuo* and the residue suspended in toluene (200 mL). The toluene was removed on the rotary evaporator and this procedure was repeated again to remove all traces of thionyl chloride. The chloride was then dissolved in DMSO (minimum to dissolve) and added to a solution of sodium cyanide (2.90 g, 59 mmol, 6 eq) in DMSO (200 mL). The solution was stirred overnight under N<sub>2</sub> at room temperature after which 500 mL H<sub>2</sub>O was added and the aqueous layer was extracted three times with 300 mL of ethyl acetate. The organic layers were dried and concentrated and the residue flash chromatographed in 4:1 hexane/ethyl acetate over silica gel to give 1.62 g of a light yellow solid; m.p. 109.5-113.0°. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 2H, J = 10Hz); 7.12 (d, 2H, J = 10Hz); 3.51 (s, 2H); 2.60 (t, 2H, J = 7Hz); 1.70 (m, 2H); 1.40 (m, 2H); 0.90 (t, 3H, J = 7Hz). Mass spectrum shows M<sup>+</sup> = 312/314. Mass Calcd. for C<sub>17</sub>H<sub>17</sub>ClN<sub>4</sub>: 312.1139, Found 312.1126.

The intermediates shown below were prepared, or could be prepared, in accordance with the procedure described in Example 1, Part B using the appropriately substituted imidazole and benzyl halide as starting material.



R <sup>1</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
4-CN	n-butyl	CH <sub>2</sub> CN	Cl	(oil) <sup>a</sup>
4-NO <sub>2</sub>	n-butyl	Cl	CH <sub>2</sub> CN	117.0-119
4-NO <sub>2</sub>	n-butyl	CH <sub>2</sub> CN	Cl	(oil) <sup>b</sup>
2-CN	n-butyl	Cl	CH <sub>2</sub> CN	(oil) <sup>c</sup>
3-CN	n-butyl	Cl	CH <sub>2</sub> CN	(oil) <sup>d</sup>

a NMR (200 MHz, CDCl<sub>3</sub>) δ 7.66 (d, 2H, J = 7Hz); 7.12 (d, 2H, J = 7Hz); 5.15 (s, 2H); 3.69 (s, 2H); 2.56 (t, 2H, J = 7Hz); 1.62 (t of t, 2H, J = 7,7Hz); 1.33 (t of q, 2H, J = 7,7Hz); 0.87 (t, 3H, J = 7Hz).

b NMR (200 MHz, CDCl<sub>3</sub>) δ 8.24 (d, 2H, J = 10Hz); 7.18 (d, 2H, J = 10Hz); 5.20 (s, 2H); 3.67 (s, 2H); 2.55 (t, 2H, J = 7Hz); 1.64 (m, 2H); 1.34 (m, 2H); 0.85 (t, 3H, J = 7Hz).

c NMR (200 MHz, CDCl<sub>3</sub>) δ 7.80 (d, 1H, J = 10Hz); 7.64 (d of d, 1H, J = 10,10Hz); 7.53 (d of d, 1H, J = 10,10Hz); 6.74 (d, 1H, J = 10Hz); 5.37 (s, 2H); 3.64 (s, 2H); 2.55 (t, 2H, J = 7Hz); 1.67 (m, 2H); 1.34 (m, 2H); 0.85 (t, 3H, J = 7Hz).

d NMR (200 MHz, CDCl<sub>3</sub>) δ 7.66 (d, 1H, J = 7Hz); 7.54 (d of d, 1H, J = 7,7Hz); 7.33 (s, 1H); 7.25 (d, 1H, J = 7Hz); 5.25 (s, 2H); 3.56 (s, 2H); 2.61 (t, 2H, J = 7Hz); 1.69 (m, 2H); 1.35 (m, 2H); 0.91 (t, 3H, J = 7Hz).

#### PART C: Preparation of 2-Butyl-1-(4-carboxybenzyl)-4-chloroimidazole-5-acetic acid

2-Butyl-4-chloro-1-(4-cyanobenzyl)-5-(cyanomethyl)imidazole (0.5 g) and a solution of 1:1 12 N HCl/glacial acetic acid (10 mL) were mixed and refluxed for 6 hours. The solvents were removed by rotary evaporation and the resultant solids were washed with isopropanol, and filtered. The mother liquor was flash chromatographed on silica gel in 1:1 hexane/ethyl acetate to give 60 mg of product. Further flushing of the column with isopropanol followed by preparatory TLC of the evaporated residue gave an additional 100 mg of product. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.90 (d, 2H, J = 8Hz); 7.12 (d, 2H, J = 8Hz); 5.30 (s, 2H); 3.08 (s, 2H); 2.50 (t, 2H, J = 7Hz); 1.49 (m, 2H); 1.24 (m, 2H); 0.79 (t, 3H, J = 7Hz). Mass. Calcd. for C<sub>13</sub>H<sub>19</sub>ClN<sub>2</sub>O<sub>4</sub>: 350.1033. Found 350.1066.

#### Example 2

##### PART A: Preparation of 2-Butyl-4-chloro-1-(4-nitrobenzyl)imidazole-5-acetic acid

2-Butyl-4-chloro-5-(cyanomethyl)-1-(4-nitrobenzyl)imidazole (7.08 g) and a 1:1 mixture of 12 N HCl and glacial acetic acid (175 mL) were mixed and refluxed for 6 hours. The solvents were removed by rotary evaporation and water (300 mL) was then added to the residue. After a few minutes, the product precipitated and was collected and dried to give 7.35 g of a solid; m.p. 207.0-210.0°. NMR (200 MHz, DMSO-d<sub>6</sub>/CDCl<sub>3</sub>) δ 8.20 (d, 2H, J = 10Hz); 7.22 (d, 2H, J = 10Hz); 5.28 (s, 2H); 3.42 (s, 2H); 2.52 (t, 2H, J = 7Hz); 1.64 (m, 2H); 1.34 (m, 2H); 0.86 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>16</sub>H<sub>18</sub>ClN<sub>3</sub>O<sub>4</sub>; C, 54.63; H, 5.16; N, 11.94. Found: C, 54.52; H, 5.05; N, 12.21.

**PART B: Preparation of Methyl 2-butyl-4-chloro-1-(4-nitrobenzyl)imidazole-5-acetate**


2-Butyl-4-chloro-1-(4-nitrobenzyl)imidazole-5-acetic acid (7.35 g, 20.9 mmol, 1eq); 3.1M HCl in dioxane (34.0 mL, 105.4 mmol, 5 eq) and 100 mL methanol were mixed and refluxed for 7.5 hours. The solvents were removed by rotary evaporation and the residue taken up in methylene chloride and 1 N NaOH (300 mL each). The layers were separated and the organic layer washed two more times with 1N NaOH (300 mL each), dried and concentrated to give 5.43 g of a light pink solid; m.p. 97.5-100.0°. NMR (200 MHz, DMSO- $d_6$ )  $\delta$  8.23 (d, 2H, J = 9Hz); 7.33 (d, 2H, J = 9Hz); 5.50 (s, 2H); 3.73 (s, 2H); 3.40 (s, 3H); 2.66 (t, 2H, J = 7Hz); 1.53 (m, 2H); 1.22 (m, 2H); 0.76 (t, 3H, J = 7Hz). Mass Calcd. for  $C_{17}H_{20}N_3O_4Cl$ : 365.1140. Found: 365.1158.

Methyl 2-butyl-5-chloro-1-(4-nitrobenzyl)imidazole-5-acetate was also prepared by the procedure described in Example 2 Part B from 2-butyl-5-chloro-1-(4-nitrobenzyl)imidazole-5-acetic acid. NMR (200 MHz,  $CDCl_3$ )  $\delta$  8.23 (d, 2H, J = 10Hz); 7.20 (d, 2H, J = 10Hz); 5.21 (s, 2H); 3.75 (s, 3H); 3.67 (s, 2H); 2.58 (t of t, 2H, J = 7Hz); 1.32 (q of t, 2H, J = 7Hz); 0.86 (t, 3H, J = 7Hz). Mass Calcd. for  $C_{17}H_{20}ClN_3O_4$ ; 365.1142. Found 365.1132.

**PART C: Methyl 2-butyl-4-chloro-1-(4-aminobenzyl)imidazole-5-acetate**

A mixture of methyl 2-butyl-4-chloro-1-(4-nitrobenzyl)imidazole-5-acetate (5.00 g, 13.7 mmol, 1 eq), iron (2.67 g, 47.8 mmol, 3.5 eq), glacial acetic acid (5.47 mL, 95.3 mmol, 7 eq), and methanol (250 mL) was refluxed for 5.5 hours. The solvent was removed by rotary evaporation. The residue was diluted with water (300 mL) and extracted five times with 300 mL portions of ethyl acetate. The organic layers were dried and concentrated. The residue was flash chromatographed in 75:25 hexane/ethyl acetate over silica gel to give 4.53 g of a golden yellow oil which crystallized after standing for several days. NMR (200 MHz,  $CDCl_3$ )  $\delta$  6.72 (d, 2H, J = 7Hz); 6.60 (d, 2H, J = 7Hz); 4.99 (s, 2H); 3.61 (s, 3H); 3.47 (s, 2H); 2.60 (t, 2H, J = 7Hz); 1.68 (m, 2H); 1.35 (m, 2H); 0.86 (t, 3H, J = 7Hz). Mass spectrum shows  $M^+ = 335/337$ . Mass Calcd. for  $C_{17}H_{22}N_3O_2Cl$ : 335.1400. Found: 335.1407.

The following intermediates were prepared by the procedure described in Example 2, Part C from the corresponding nitro intermediates:



R <sup>1</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
4-NH <sub>2</sub>	n-butyl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	Cl	(oil) <sup>a</sup>
4-NH <sub>2</sub>	n-butyl	Cl	OCOCH <sub>3</sub>	(oil) <sup>b</sup>
4-NH <sub>2</sub>	n-butyl	Cl	CH <sub>2</sub> OH	(oil) <sup>c</sup>

a NMR (200 MHz,  $CDCl_3$ )  $\delta$  6.85 (d, 2H, J = 7Hz); 6.63 (d, 2H, J = 7Hz); 4.95 (s, 2H); 3.69 (s, 3H); 2.57 (t, 2H, J = 7Hz); 1.59 (t of t, 2H, J = 7,7Hz); 1.30 (t of q, 2H, J = 7,7Hz); 0.86 (t, 3H, J = 7Hz).

b NMR (200 MHz,  $CDCl_3$ )  $\delta$  6.74 (d, 2H, J = 10Hz); 6.60 (d, 2H, J = 10Hz); 4.97 (s, 2H); 4.95 (s, 2H); 3.56 (t, 2H, J = 7Hz); 1.86 (s, 3H); 1.64 (t of t, 2H, J = 7,7Hz); 1.33 (t of q, 2H, J = 7,7Hz); 0.85 (t, 3H, J = 7Hz).

c NMR (200 MHz,  $CDCl_3$ )  $\delta$  6.80 (d, 2H, J = 10Hz); 6.69 (d, 2H, J = 10Hz); 5.05 (s, 2H); 4.43 (s, 2H); 2.56 (t, 2H, J = 7Hz); 1.56 (t of t, 2H, J = 7,7Hz); 1.26 (t of q, 2H, J = 7,7Hz); 0.83 (t, 3H, J = 7Hz).



PART D: Preparation of Methyl 2-butyl-1-[4-(2-carboxybenzamido)benzyl]-4-chloroimidazole-5-acetate

A chloroform solution (10 mL) of methyl 2-butyl-4-chloro-1-(4-aminobenzyl)imidazole-5-acetate (500 mg, 1.5 mmol, 1 eq) was mixed with a chloroform solution (10 mL) of phthalic anhydride (221 mg, 1.5 mmol, 1 eq). After five minutes of stirring at room temperature, product began to precipitate. After 24 hours, the product was filtered, washed with a minimum amount of  $\text{CHCl}_3$  and dried to give 400 mg of a white solid. After some evaporation, the mother liquor yielded an additional 220 mg of product, both of which had identical melting points; m.p. 109.5 - 112.5°. NMR (200 MHz,  $\text{DMSO-d}_6$ )  $\delta$  10.37 (s, 1H); 7.85 (d, 2H,  $J = 8\text{Hz}$ ); 7.71-7.50 (m, 5H); 6.96 (d, 2H,  $J = 10\text{Hz}$ ); 5.12 (s, 2H); 3.60 (s, 2H); 3.49 (s, 3H); 2.55 (t, 2,  $J = 7\text{Hz}$ ); 1.52 (m, 2H); 1.27 (m, 2H); 0.83 (t, 3H,  $J = 7\text{Hz}$ ). The carboxylic acid could be titrated with 1.000 N NaOH to form the sodium salt. High resolution mass spectrum shows M-18 (loss of  $\text{H}_2\text{O}$ ). Calcd. Mass for  $\text{C}_{25}\text{H}_{26}\text{ClN}_3\text{O}_5$ : 465.1455. Found: 465.1440.

Example 3

15

PART A: Preparation of 2-Butyl-5-chloro-1-(4-nitrobenzyl)imidazole-4-acetic acid

2-Butyl-5-chloro-4-cyanomethyl-1-(4-nitrobenzyl)imidazole (4.48 g) was converted to the corresponding carboxylic acid by the procedure described in Example 2, Part A. No product precipitated upon the addition of water (300 mL) until the pH was raised to about 3 with conc. ammonium hydroxide to liberate the imidazole from its HCl salt. The precipitated solids were amorphous and ethyl acetate (5 x 300 mL) was used to extract the product. The organic layers were dried and concentrated to give 3.93 g of a yellow solid. Recrystallization from hexane/ethyl acetate gave 3.06 g of a white solid; m.p. = 138.0-139.5°. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.25 (d, 2H,  $J = 10\text{Hz}$ ); 7.21 (d, 2H,  $J = 10\text{Hz}$ ); 5.23 (s, 2H); 3.30 (s, 2H); 2.63 (t, 2H,  $J = 7\text{Hz}$ ); 1.63 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.32 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.87 (t, 3H,  $J = 7\text{Hz}$ ). Anal. Calcd. for  $\text{C}_{16}\text{H}_{18}\text{ClN}_3\text{O}_4$ ; C, 54.63; H, 5.16; N, 11.94. Found: C, 54.75; H, 5.29; N, 12.14.

PART B: Preparation of Methyl 2-butyl-1-[4-(2-carboxybenzamido)benzyl]-5-chloroimidazole-4-acetate

2-Butyl-5-chloro-1-(4-nitrobenzyl)imidazole-4-acetic acid (Part A) was carried on to methyl 2-butyl-1-[4-(2-carboxybenzamido)benzyl]-5-chloroimidazole-4-acetate; m.p. 150.5-152.5° by the procedure described in Example 2. NMR (200 MHz,  $\text{DMSO-d}_6$ )  $\delta$  13.00 (bs, 1H); 10.40 (s, 1H); 7.87 (d, 1H,  $J = 8\text{Hz}$ ); 7.67 (d, 2H,  $J = 8\text{Hz}$ ); 7.71-7.52 (m, 3H); 7.02 (d, 2H,  $J = 8\text{Hz}$ ); 5.13 (s, 2H); 3.61 (s, 3H); 3.52 (s, 2H); 2.59 (t, 2H,  $J = 7\text{Hz}$ ); 2.53 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.28 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.82 (t, 3H,  $J = 7\text{Hz}$ ). Mass Calcd. for  $\text{C}_{25}\text{H}_{26}\text{ClN}_3\text{O}_5 \cdot \text{H}_2\text{O}$ : 465.1455. Found, 465.1460.

Example 4PART A: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-(4-nitrobenzyl)imidazole

40

2-n-butyl-4-chloro-5-hydroxymethyl-1-(4-nitrobenzyl)imidazole (10.5 g, 32.4 mmol, 1 eq), conc. sulfuric acid (26 mL) and methanol (300 mL) were mixed and refluxed overnight. The solvent was removed in vacuo and the residue taken up in water (about 300 mL). The pH was adjusted to 5 with 1N NaOH and then this aqueous portion extracted with ethyl acetate (3 x 250 mL). The organic layers were collected, dried ( $\text{MgSO}_4$ ) and the solvent removed in vacuo to yield 11.57 g of an amber oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.22 (d, 2H,  $J = 8\text{Hz}$ ); 7.15 (d, 2H,  $J = 8\text{Hz}$ ); 5.26 (s, 2H); 4.25 (s, 2H); 3.23 (s, 3H); 2.52 (t, 2H,  $J = 7\text{Hz}$ ); 1.64 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.28 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.81 (t, 3H,  $J = 7\text{Hz}$ ). Anal. Calcd. for  $\text{C}_{16}\text{H}_{20}\text{ClN}_3\text{O}_3 \cdot (\text{H}_2\text{O})_{0.5}$ ; C, 55.41; H, 6.10; Cl, 10.22. Found: C, 55.21; H, 6.22; Cl, 9.92.

PART B: Preparation of 1-(4-Aminobenzyl)-2-n-butyl-4-chloro-5-(methoxymethyl)imidazole

To a solution of 2-n-butyl-4-chloro-5-methoxymethyl-1-(4-nitrobenzyl)imidazole (11.22 g) in methanol (100 mL) under  $\text{N}_2$  was carefully added 1.0 g of 10% palladium on charcoal. Hydrogen gas was then bubbled through the solution for 4 hours. The solution was filtered through Celite® and the solvent removed in vacuo to yield 9.23 g of an amber oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.99 (s, 1H); 6.78 (d of d, 4H,  $J = 5,5\text{Hz}$ ); 5.05 (s, 2H); 4.24 (s, 2H); 3.27 (s, 3H); 2.59 (t, 2H,  $J = 7\text{Hz}$ ); 1.62 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.32 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.84 (t, 3H,  $J = 7\text{Hz}$ ). Mass Calcd. for  $\text{C}_{16}\text{H}_{23}\text{ClN}_3\text{O}$ ; 307.1451. Found: 307.1460.



PART C: Preparation of 2-Butyl-1-[4-(2-carboxybenzamido)benzyl]-4-chloro-5-(methoxymethyl)imidazole

The above compound was prepared from 1-(4-aminobenzyl)-2-n-butyl-4-chloro-5-(methoxymethyl) im-  
idazole (3.00 g, 9.7 mmol, 1 eq) and phthalic anhydride (1.44 g, 9.7 mmol, 1 eq) using the procedure of  
5 Example 2, Part D. Work-up yielded 1.71 g of an off-white powder, which was washed with acetonitrile. The  
insoluble material was filtered and dried to yield 1.17 g of a white powder; m.p. 165.5-166.5 °C. NMR (200  
MHz, DMSO-d<sub>6</sub>) δ 13.01 (m, 1H); 10.39 (s, 1H); 7.87 (d, 1H, J = 7Hz); 7.75-7.46 (m, 5H); 7.03 (d, 2H, J =  
8Hz); 5.16 (s, 2H); 4.30 (s, 2H); 3.20 (s, 3H); 2.54 (t, 2H, J = 7Hz); 1.54 (t of t, 2H, J = 7,7Hz); 1.30 (t of q,  
10 2H, J = 7,7Hz); 0.83 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>24</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>4</sub>:C, 63.22; H, 5.75; Cl, 7.78. Found: C,  
63.54; H, 5.76; Cl, 7.58.

Examples 5-18 shown in Table 1 were prepared or could be prepared by the procedures described in  
Examples 2-4 from the appropriately substituted aniline derivative and a suitable anhydride or acid chloride.  
Other solvents, such as benzene or ethyl acetate may be substituted for chloroform.

15

20

25

30

35

40

45

50

55

Table 1






Ex. No.	R	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>MP(°C)</u>
5		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	(oil) <sup>a</sup>
6		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	138.0-141.0
7		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	184.0-186.0
8		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	169.0-170.5
9		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	172.0-173.5

Table 1 (cont'd.)











Ex. No.	R	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
10		n-butyl	Cl	$\text{CH}_2\overset{\text{O}}{\overset{\parallel}{\text{C}}}\text{CH}_3$	140.0-144.5
11		n-butyl	Cl	$\text{CH}_2\text{CO}_2\text{CH}_3$	129-131
12		n-butyl	Cl	$\text{CH}_2\text{CO}_2\text{CH}_3$	119-121
13		n-butyl	Cl	$\text{CH}_2\text{CO}_2\text{CH}_3$	148-151
14		n-butyl	Cl	$\text{CH}_2\text{CO}_2\text{CH}_3$	159-160
15		n-butyl	Cl	$\text{CH}_2\text{CO}_2\text{CH}_3$	175-176
					

Table 1 (cont'd.)

Ex. No.	R	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP (°C)
16		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	199.0-200.0
17		n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>	173.5-177.0
18		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	151-153
<p>a NMR (200 MHz, CDCl<sub>3</sub>) δ 9.48 (bs, 1H); 7.87-7.61 (m, 2H); 7.5-7.04 (m, 8H); 6.69 (d, 2H, J = 9Hz); 4.98 (s, 2H); 3.45 (s, 3H); 3.40 (s, 2H); 2.56 (m, 2H); 1.48 (m, 2H); 1.26 (m, 2H); 0.72 (t, 3H, J = 7Hz).</p>					

## Example 19

## Preparation of 2-Butyl-4-chloro-5-hydroxymethyl-1-(4-carboxybenzyl)imidazole

The title compound was prepared from 2-butyl-4-chloro-5-hydroxymethyl-1-(4-cyanobenzyl)imidazole by the method described in Example 2, Part A. NMR (200 MHz, CDCl<sub>3</sub> + DMSO-d<sub>6</sub>) δ 7.96 (d, 2H, J = 8Hz); 7.13 (d, 2H, J = 8Hz); 5.33 (s, 2H); 4.40 (s, 2H); 2.50 (t, 2H, J = 7Hz); 1.57 (t of t, 2H, J = 7,7Hz); 1.27 (t of q, 2H, J = 7,7Hz); 0.85 (t, 3H, J = 7Hz).

## Example 20

## Preparation of 5-Acetoxymethyl-2-butyl-1-(4-carboxybenzyl)-4-chloroimidazole

2-Butyl-1-(4-carboxybenzyl)-4-chloro-5-(hydroxymethyl)imidazole (2.00 g, 6.2 mmol, 1 eq), acetic anhydride (1.46 mL, 15.5 mmol, 2.5 eq), triethylamine (2.59 mL, 18.6 mmol, 3 eq) and THF (50 mL) were mixed and stirred for 3 days. Water (200 mL) was added to the solution and the mixture was stirred for 0.5 hours. The pH was lowered to 5 with conc. HCl and the mixture extracted with ethyl acetate (3 x 100 mL). The organic layers were dried (MgSO<sub>4</sub>) and concentrated to give 2.47 g of a brown oil. This product (2.16 g) was dissolved in a minimum of ethyl acetate and dicyclohexylamine (DCHA) (1.18 mL, 1 eq) was added and mixed. The solution was allowed to slowly evaporate overnight. The DCHA salt so obtained (1.43 g) was subsequently taken up in ethyl acetate (100 mL), and washed with 1 N HCl (3 x 100 mL), followed by brine.

The organic layer was dried (MgSO<sub>4</sub>) and concentrated to give a yellow oil (670 mg). NMR (200 MHz, CDCl<sub>3</sub>) δ 8.09 (d, 2H, J = 10Hz); 7.05 (d, 2H, J = 10Hz); 5.20 (s, 2H); 4.98 (s, 2H); 2.58 (t, 2H, J = 7Hz); 1.82 (t of t, 2H, J = 7,7Hz); 1.33 (t of q, 2H, J = 7,7Hz); 0.86 (t, 3, J = 7Hz). Anal. Calcd. for C<sub>18</sub>H<sub>21</sub>ClN<sub>2</sub>O<sub>4</sub>; C, 59.26; H, 5.80, N, 7.68. Found: C, 58.89; H, 6.17; N, 7.39. Mass Calcd. for C<sub>18</sub>H<sub>21</sub>ClN<sub>2</sub>O<sub>4</sub>: 364.1200. Found: 364.1167.


#### Example 21

##### Preparation of Methyl 2-butyl-4-chloro-1-[4-(trifluoromethylsulfonamido)benzyl]imidazole-5-acetate

A solution of triflic anhydride (0.88 mL, 5.2 mmol, 1 eq) in methylene chloride (5 mL) was dripped into a solution of methyl 2-butyl-1-(4-aminobenzyl)-4-chloroimidazole-5-acetate (1.74 g, 5.2 mmol, 1 eq) and triethylamine (1.44 mL, 10.4 mmol, 2 eq) in 20 mL of methylene chloride at -78 °C. The solution was kept at -78 °C for 1 hour after which it was allowed to warm to room temperature. After 24 hours, the reaction was quenched with water (100 mL) and the pH adjusted to 5 with conc. HCl and the aqueous extracted with methylene chloride (5 x 100 mL). The organic layers were dried (MgSO<sub>4</sub>), concentrated, and the residue flash chromatographed in 1:1 hexane/ethyl acetate on silica gel. The crystalline product which formed in the 1:1 hexane/ethyl acetate solution while the crude product was being applied to the column was isolated (1.03 g). Chromatography of the mother liquor yielded an additional 1.03 g of the title compound as a white solid; m.p. 154.0-157.0 °. The product could be titrated with 1 equivalent of 1.000 N NaOH. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.32 (d, 2H, J = 10Hz); 6.91 (d, 2H, J = 10Hz); 5.15 (s, 2H); 3.62 (s, 3H); 3.46 (s, 2H); 2.55 (t, 2H, J = 7Hz); 1.56 (m, 2H); 1.26 (m, 2H); 0.72 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>18</sub>H<sub>21</sub>N<sub>3</sub>O<sub>4</sub>SF<sub>3</sub>Cl: 467.0890. Found: 467.0872.

Examples 22-25 in Table 2 were prepared or could be prepared by the procedure described in the above example employing the appropriately substituted 1-(aminobenzyl)-imidazole, which in some instances is followed by ester hydrolysis familiar to one skilled in the art.

Table 2



Ex. No.	R <sup>1</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP (°C)
22	NHSO <sub>2</sub> CF <sub>3</sub>	n-butyl	Cl	CH <sub>2</sub> OH	
23	NHSO <sub>2</sub> CF <sub>3</sub>	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>	
24	NHSO <sub>2</sub> CF <sub>3</sub>	n-butyl	Cl	CH <sub>2</sub> OCH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	
25	NHSO <sub>2</sub> CF <sub>3</sub>	n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> H	(oil) <sup>a</sup>

<sup>a</sup> NMR (200 MHz, CDCl<sub>3</sub>) δ 7.29 (d, 2H, J = 10Hz); 6.64 (d, 2H, J = 10Hz); 5.11 (s, 2H); 3.45 (s, 2H); 2.56 (t, 2H, J = 7Hz); 1.60 (m, 2H); 1.30 (m, 2H); 0.85 (t, 3H, J = 7Hz)

#### Example 26

##### Preparation of 2-Butyl-4-chloro-5-[(1H-tetrazol-5-yl)methyl]-1-[3-(1H-tetrazol-5-yl)benzyl]imidazole

2-Butyl-4-chloro-1-(3-cyanobenzyl)-5-(cyanomethyl)imidazole (2.00 g, 6.4 mmol, 1 eq); ammonium chloride (0.91 g, 17 mmol, 2.7 eq); sodium azide (1.11 g, 17 mmol, 2.7 eq) and DMF (25 mL) were mixed and stirred at 80°C for 24 hours. The mixture was filtered and the solvent removed by rotary evaporation. The residue was dissolved in water (100 mL) and methylene chloride (100 mL). The layers were separated and the aqueous layer extracted again with methylene chloride (2 x 100 mL). The aqueous was then acidified with conc. HCl to pH of 3. The solid which precipitated was collected and dried to give 560 mg of the title compound as a tan solid; m.p. 254° (darken), 258° (dec.). The product when titrated with 1.000 N NaOH showed the presence of exactly two acidic functionalities. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.79 (d, 1H, J = 7Hz); 7.69 (s, 1H); 7.53 (t, 1H, J = 7Hz); 7.10 (d, 1H, J = 7Hz); 5.37 (s, 2H); 4.23 (s, 2H); 2.57 (t, 2H, J = 7Hz); 1.53 (t of t, 2H, J = 7Hz); 1.27 (t of q, 2H, J = 7 Hz); 0.80 (t, 3H, J = 7Hz); Anal. Calcd. for C<sub>17</sub>H<sub>19</sub>ClN<sub>10</sub>: C, 51.19; H, 4.80. Found: C, 51.04; H, 4.69.

#### Example 27

##### Preparation of 2-Butyl-4-chloro-5-[(1H-tetrazol-5-yl)methyl]-1-[4-(1H-tetrazol-5-yl)benzyl]imidazole

The title compound was prepared from 2-butyl-4-chloro-1-(4-cyanobenzyl)-5-(cyanomethyl)imidazole by the procedure described in Example 26; m.p. 228 (dark), 229.0-230° (dec). Titration with 1.000 N NaOH

showed the presence of exactly two acid functionalities. NMR (200 MHz, DMSO- $d_6$ )  $\delta$  7.95 (d, 2,  $J$  = 7Hz); 7.13 (d, 2,  $J$  = 7Hz); 5.34 (s, 2); 4.23 (s, 2); 2.53 (t, 2,  $J$  = 7Hz); 1.50 (t of t, 2,  $J$  = 7,7Hz); 1.26 (t of q, 2,  $J$  = 7Hz); 0.79 (t, 3,  $J$  = 7Hz); 1R 3420 br, 1930 br, 740  $\text{cm}^{-1}$ . Mass Calcd. for  $\text{C}_{13}\text{H}_{19}\text{ClN}_3\text{O}$ : 398.1482. Found: 398.1509.

5

#### Example 28

##### Preparation of 2-Butyl-4-chloro-5-hydroxymethyl-1-(4-N-phthalimidobenzyl)imidazole

10 1-(4-Aminobenzyl)-2-butyl-4-chloro-5-(hydroxymethyl)imidazole (1.00 g, 3.4 mmol, 1 eq) in 20 mL of methylene chloride was dripped into a stirred solution of phthaloyl chloride (0.49 mL, 3.4 mmol, 1 eq), triethylamine (0.95 mL, 6.82 mmol, 2 eq) and methylene chloride (500 mL). After 11 days, the solvent was removed by rotary evaporation and the residue flash chromatographed in 1:1 hexane/ethyl acetate over silica gel to give 240 mg of the title compound as a light yellow glassy solid; m.p. 65.0-73.5°, NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  (key peaks only) 7.97 (m, 2H); 7.79 (m, 2H); 7.43 (d, 2,  $J$  = 10Hz); 7.11 (d, 2H,  $J$  = 10Hz); 15 4.50 (s, 2H); 2.57 (t, 2H,  $J$  = 7Hz); 1.67 (m, 2H); 1.34 (m, 2H); 0.87 (t, 3H,  $J$  = 7Hz). Mass Calcd. for  $\text{C}_{23}\text{H}_{22}\text{ClN}_3\text{O}_3$ : 423.1349. Found: 423.1324.

#### Example 29

20

##### Preparation of Methyl 2-butyl-4-chloro-1-(4-N-phthalimidobenzyl)imidazole-5-acetate

Methyl 2-butyl-1-[4-(2-carboxybenzamido)benzyl]-4-chloroimidazole-5-acetate (1.00 g), methanol (50 mL) and 3.6 mL of 3.1 N HCl in dioxane were refluxed for 6 days. The solvent was removed in vacuo and the residue taken up in ethyl acetate (100 mL). The organic phase was washed with 1 N NaOH (2 x 100 mL) and brine (1 x 100 mL), dried ( $\text{MgSO}_4$ ) and concentrated. The residue was flash chromatographed over silica gel in 75:25 hexane/ethyl acetate to give 400 mg of an oil which eventually crystallized; m.p. 141.5 - 143.0°. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.92 (m, 2H); 7.80 (m, 2H); 7.43 (d, 2H,  $J$  = 10Hz); 7.08 (d, 2H,  $J$  = 10Hz); 5.17 (s, 2H); 3.62 (s, 3H); 3.50 (s, 2H); 2.62 (t, 2H,  $J$  = 7Hz); 1.71 (t of t, 2H,  $J$  = 7,7Hz); 1.36 (t of q, 20 2H,  $J$  = 7,7Hz); 0.89 (t, 3H,  $J$  = 7Hz). Mass Calcd. for  $\text{C}_{25}\text{H}_{24}\text{ClN}_3\text{O}_4$ : 465.1455. Found: 465.1440.

30

#### Example 30

##### Preparation of Methyl 2-butyl-4-chloro-1-[4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl]imidazole-5-acetate

35

Methyl-1-(4-aminobenzyl)-2-butyl-4-chloro-5-imidazoleacetate (1.00 g, 2.98 mmol, 1 eq), N-(trifluoromethanesulfonyl)anthranoyl chloride which is described in EP 003836, (0.86 g, 2.99 mmol, 1 eq), and sodium bicarbonate (1.25 g, 14.9 mmol, 5 eq) were mixed and stirred in 50 mL methylene chloride (acid chloride was added last). The reaction was worked up after 2.5 hours by filtering, removing the solvent from the filtrate in vacuo and recrystallizing the residue from ethyl acetate/hexane to give 1.07 g of light yellow crystals; m.p. 151.0 - 152.0°. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  9.32 (s, 1H); 8.02 (d, 1H,  $J$  = 10Hz); 7.79 (d, 1H,  $J$  = 10Hz); 7.56 (d of d, 2H,  $J$  = 10, 10Hz); 7.50 (d, 2H,  $J$  = 10Hz); 7.78 (d of d, 1H,  $J$  = 10, 10Hz); 6.86 (d, 2H,  $J$  = 10Hz); 5.10 (s, 2H); 3.58 (s, 3H); 3.45 (s, 2H); 2.45 (t, 2H,  $J$  = 7Hz); 1.52 (t of t, 2H,  $J$  = 7,7Hz); 1.22 (t of q, 2H,  $J$  = 7,7Hz); 0.75 (t, 3H,  $J$  = 7Hz). Titration of the product with 1.000 N NaOH shows the presence of exactly one acidic functionality. Anal. Calcd. for  $\text{C}_{25}\text{H}_{26}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : C, 51.15; H, 4.46; N, 9.54. Found: C, 50.95; H, 4.26; N, 9.67. Mass Calcd. for  $\text{C}_{25}\text{H}_{26}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : 586.1264. Found: 586.1222.

45

#### Example 31

50

##### Preparation of 2-Butyl-4-chloro-1-[4-((N-trifluoro methanesulfonyl)anthranilamido)benzyl]imidazole-5-acetic acid

Methyl 2-butyl-4-chloro-1-[4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl]imidazole-5-acetate (400 mg, 0.66 mmol, 1 eq) was stirred in 1.0 N NaOH (0.66 mL, 0.66 mmol, 1 eq) for 3 hours under  $\text{N}_2$ . The pH was adjusted to 5 with 1.0 N HCl and the product precipitate was collected and dried affording 120 mg of the title compound as a white solid. The NMR spectrum shows the methyl ester to be missing. Mass spectrum shows M- $\text{CO}_2$  peak. Mass Calcd. for  $\text{C}_{23}\text{H}_{24}\text{ClF}_3\text{N}_4\text{O}_3\text{S}$ : 528.1209. Found: 528.1236.

55

Example 32

Preparation of 2-Butyl-1-[4-(2-carboxybenzamido)benzyl]-4-chloroimidazole-5-acetic acid

5     The title compound was prepared from methyl 2-butyl-1-[4-(2-carboxybenzamido)benzyl]-4-chloroimidazole-5-acetate by the procedure described in Example 31; m.p. 170.5 - 175.0°.

     Examples 33-53 in Table 3 were prepared or could be prepared by the procedures described in Examples 30 and 31 using the appropriate aniline and acid chloride starting materials.

10

15

20

25

30

35

40


45

50

55



Table 3



Ex. No.	R	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
33		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	(oil) <sup>a</sup>
34		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	
35		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	226-228
36		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	153-156 (dec.)
37		n-propyl	Cl	CH <sub>2</sub> OH	
38		n-hexyl	H	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	

Table 3 (cont'd.)


















Ex. No.	R	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
39		n-propyl	Cl	CH <sub>2</sub> OH	
40		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	
41		n-propyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	
42		n-butyl	Cl	CH <sub>2</sub> OH	
43		CH <sub>3</sub> CH <sub>2</sub> CH=CH-	Cl	CH <sub>2</sub> OH	
44		n-butyl	Cl	CH <sub>2</sub> OCOCH <sub>3</sub>	
45		n-butyl	Cl	CH <sub>2</sub> OCOCH <sub>3</sub>	
46		n-butyl	CH <sub>2</sub> CO <sub>2</sub> H	Cl	

Table 3 (cont'd.)

Ex. No.	R	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	MP(°C)
47		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> H	
48		n-butyl	Cl	n-butyl	
49		n-butyl	CH <sub>2</sub> CO <sub>2</sub> H	Cl	
50		n-hexyl	Cl	CH <sub>2</sub> CO <sub>2</sub> H	
51		n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	74.0-79.5
52		n-butyl	Cl		200.5-205.0
53		n-propyl	Cl		

a NMR (200 MHz, CDCl<sub>3</sub>) δ 8.69 (s, 1H); 7.82 (s, 1H); 7.75 (d, 1H, J= 7Hz); 7.59 (d, 2H, J= 10Hz); 7.55 (d, 1H, J= 7Hz); 7.45 (t, 1H, J= 7Hz); 6.87 (d, 2H, J= 10Hz); 5.06 (s, 2H); 3.60 (s, 3H); 3.46 (s, 2H); 2.54 (t, 2H, J= 7Hz); 1.55 (t of t, 2H, J= 7,7Hz); 1.24 (t of q, 2H, J= 7,7Hz); 0.78 (t, 3H, J= 7Hz).

Example 54PART A: Preparation of Ethyl n-heptylimidate hydrochloride

5 To a solution of caprylonitrile (30 g, 0.24 mol) in 25 mL of absolute ethanol cooled to 0 ° was bubbled HCl gas (9.6 g, 0.26 mol). After 7 days at 0 ° the viscous solution was diluted with 250 mL of anhydrous ether and the precipitated product was filtered with suction onto a coarse frit and washed liberally with ether before placing under a vacuum to remove residual solvent. The product was stored under nitrogen at 0 ° to yield 22 g (44%) of a white solid. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 4.40 (q, 2H, J = 7Hz); 3.30 (m, 4H); 2.45 (m, 10 4H); 1.40-0.75 (m, 12H). Mass Spec. 172 (M-Cl).

PART B: Preparation of 2-Heptyl-5-(hydroxymethyl)imidazole

In a high-pressure (bomb) reactor was placed ethyl n-heptylimidate hydrochloride (22 g, 0.11 mol), 1,3-  
15 dihydroxyacetone dimer (9.5 g, 0.053 mol) and liquid ammonia (60 g, 3.5 mol). The reactor was sealed and heated to 70 ° for 12 hours. The crude product (24.7 g) was purified by flash chromatography (silica gel, 300 g; 10:1 EtOAc/EtOH) to give 12.7 g (61%) of a light yellow solid; m.p. 82-84 °. NMR (200 MHz, CDCl<sub>3</sub>/Acetone-d<sub>6</sub>) δ 6.75 (s, 1H); 4.50 (s, 2H); 4.50-4.25 (br s, 2H); 2.60 (t, 2H, 8Hz); 1.75-1.60 (m, 2H); 1.40-1.15 (m, 8H); 0.95-0.75 (m, 3H). Mass Spec. 196, 167 (M-Et), 149 (M-Et-H<sub>2</sub>O).

PART C: Preparation of 4-Chloro-2-heptyl-5-hydroxymethylimidazole

To a solution of 2-heptyl-5-(hydroxymethyl)imidazole (10.0 g, 51 mmol) in EtOH/1,4-dioxane (1:1; 600 mL) was added N-chlorosuccinimide (7.9 g, 59 mmol). After being stirred for 1 hour at room temperature  
25 the solvents were removed on a rotary evaporator and the solid residue was partitioned between ethyl acetate and water (300 mL each). The organic phase was washed with water (150 mL), dried (MgSO<sub>4</sub>), filtered and concentrated to afford 12.4 g crude product. Recrystallization (1:1 EtOAc/hexane, 60 mL) gave 5.7 g (45%) of white crystals; m.p. 134-140 °. NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD) δ 4.50 (s, 2H); 4.00-3.80 (br s, 2H); 2.65 (t, 2H, 5Hz); 1.80-1.60 (m, 2H); 1.40-1.20 (m, 8H); 0.90-0.80 (m, 3H). Mass Spec. 230.

PART D: Preparation of 4-Chloro-2-heptyl-5-(hydroxymethyl)-1-(4-nitrobenzyl)imidazole

To a solution of 4-chloro-2-heptyl-5-(hydroxymethyl)imidazole (5.2 g, 20.7 mmol) in dry DMF (100 mL) was added anhydrous K<sub>2</sub>CO<sub>3</sub> (4.3 g, 31.1 mmol) followed by 4-nitrobenzylbromide (5.4 g, 24.9 mmol). The  
35 solution was stirred 3-5 hours at 65-70 °. The reaction mixture was poured into a separatory funnel containing EtOAc and H<sub>2</sub>O (300 mL each). The aqueous phase was extracted with EtOAc (150 mL) and the combined organic phases were washed three times with H<sub>2</sub>O (150 mL) before being dried (MgSO<sub>4</sub>), filtered and concentrated to give 9.0 g brown crude oil. Chromatography (silica gel, 450 g; 1:1 EtOAc/hexanes) gave 1.3 g (17% overall, 35% of theoretical); m.p. 110-115 °. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.20 (d, 2H, 5Hz);  
40 7.20 (d, 2H, 5Hz); 5.35 (s, 2H); 4.45 (s, 2H); 3.10-3.00 (m, 1H); 2.50 (t, 2H, 5Hz); 1.75-1.50 (m, 2H); 1.40-1.10 (m, 8H); 0.90-0.75 (m, 3H). Mass Spec. 365.

PART E: Preparation of 1-(4-Aminobenzyl)-4-chloro-2-heptyl-5-hydroxymethylimidazole

To a solution of 4-chloro-2-heptyl-5-hydroxymethyl-1-(4-nitrobenzyl)imidazole (1.00 g, 2.7 mmol) in EtOH (30 mL) and glacial acetic acid (5 mL) was added iron powder (2.5 g, 44.8 mmol). The mixture was stirred while being refluxed for 20 minutes. The solution was cooled, the iron was removed by filtration, and the solution was partitioned between EtOAc and 20% aq. K<sub>2</sub>CO<sub>3</sub> (150 mL each). The organic phase was washed with saturated aqueous NaCl, dried (MgSO<sub>4</sub>), filtered and concentrated to afford 0.8 g yellow-  
50 orange oil. Flash chromatography (silica gel, 25 g; EtOAc/hexanes, 1:1) gave 0.74 g (80%) of yellow-orange oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 6.80-6.60 (ABq, 4H, 7Hz, 32Hz); 5.10 (s, 2H); 4.45 (s, 2H); 3.75-3.60 (m, 2H); 2.55 (t, 2H, 5Hz); 1.75-1.65 (m, 2H); 1.30-1.15 (m, 8H); 0.90-0.80 (m, 3H). Mass Spec. 335.

PART F: Preparation of 4-Chloro-2-heptyl-5-hydroxymethyl-1-[4-((N-trifluoromethanesulfonyl)anthranilamido)-benzyl]imidazole

To a solution of 1-(4-aminobenzyl)-4-chloro-2-heptyl-5-(hydroxymethyl)imidazole (211 mg, 0.63 mmol) in dry methylene chloride (10 mL) was added anhydrous sodium bicarbonate (263 mg, 3.1 mmol) followed

by N-(trifluoromethanesulfonyl)anthranoyl chloride (180 mg, 0.63 mmol). After 2 hours the mixture was filtered, the filtrate was concentrated and the residue was purified by flash chromatography (silica gel, 10 g; EtOAc) to provide 298 mg (81%) of pale yellow solid; m.p. 90-95 ° (dec.). NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD) δ 7.75-6.80 (m, 8H); 5.10 (s, 2H); 4.40 (s, 2H); 2.50 (t, 2H, 7Hz); 1.75-1.50 (m, 2H); 1.35-1.15 (m, 8H); 0.95-0.80 (m, 3H). Mass Spec - no mass ion observed due to apparent decomposition; 424 (M-NHSO<sub>2</sub>CF<sub>3</sub>-CH<sub>3</sub>).

#### Example 55

##### PART A: Preparation of Ethyl 3-methoxypropylimidate hydrochloride

This compound was prepared according to the procedure described in Example 54, Part A. From 3-methoxypropionitrile (30 g, 0.35 mol) and hydrogen chloride (14.1 g, 0.39 mol) in ethanol (25 mL) there was obtained 37.7 g (64%) white solid. Mass Spec. 132 (M-Cl).

##### PART B: Preparation of 5-Hydroxymethyl-2-(2-methoxyethyl)imidazole

This compound was prepared according to the procedure described in Example 54, Part B. From ethyl 3-methoxypropylimidate (36.7 g, 0.22 mol), 1,3-dihydroxyacetone dimer (19.7 g, 0.11 mol) and liquid ammonia (90 g, 5.3 mol) there was obtained 14.0 g (41%) of an off-white solid following chromatography, m.p. 100-107 °. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 6.70 (s, 1H); 4.30 (s, 2H); 3.6 (t, 2H, 5Hz); 3.20 (s, 3H); 2.80 (t, 2H, 5Hz). Mass Spec. 156.

##### PART C: Preparation of 4-Chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole

This compound was prepared according to the procedure described in Example 54, Part C. From 4-hydroxymethyl-2-(2-methoxyethyl)imidazole (13.5 g, 81.7 mmol) and N-chlorosuccinimide (13.8 g, 103 mmol) was obtained 4.8 g (29%) of light yellow solid following chromatography (silica gel, 500 g; EtOAc); m.p. 102-108 °. NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD) δ 4.50 (s, 2H); 3.65 (m, 4H); 3.40 (s, 3H); 2.90 (t, 2H, 5Hz). Mass Spec. 190.

##### PART D: Preparation of 4-Chloro-5-hydroxymethyl-2-(2-methoxyethyl)-1-(4-nitrobenzyl)imidazole

This compound was prepared according to the procedure described in Example 54, Part D. From 4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole (4.3 g, 22.6 g) was obtained 2.2 g (30% overall, 60% of theoretical) of light yellow solid; m.p. 91-95 °. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.15 (d, 2H, 8Hz); 7.20 (d, 2H, 8Hz); 5.45 (s, 2H); 4.45 (s, 2H); 3.60 (t, 2H, 5Hz); 3.20 (s, 3H); 3.15 (s, 1H); 2.80 (t, 2H, 5Hz). Mass Spec. 325.

##### PART E: Preparation of 1-(4-Aminobenzyl)-4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole

This compound was prepared according to the procedure described in Example 54, Part E. From 4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)-1-(4-nitrobenzyl)imidazole (2.2 g, 6.75 mmol) and iron powder (6.7 g, 120 mmol) there was obtained 1.6 g (80%) of light yellow solid; m.p. 164-167 °. NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD) δ 6.80 (d, 2H, 7Hz); 6.65 (d, 2H, 7Hz); 5.15 (s, 2H); 4.45 (s, 2H); 4.30 (s, 3H); 3.60 (t, 2H, 5Hz); 3.25 (s, 3H); 2.8 (t, 2H, 5Hz). Mass Spec. 295.

##### PART F: Preparation of 1-[4-(2-Carboxybenzamido)benzyl]-4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole

To an acetonitrile solution (12 mL) of 1-(4-aminobenzyl)-4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole (150 mg, 0.51 mmol) was added an acetonitrile solution (2 mL) of phthalic anhydride (75 mg, 0.51 mmol). After stirring overnight at room temperature a light yellow precipitate was produced. The mixture was cooled to 0 °, filtered with suction onto a fine fritted funnel and the solid was washed with cold acetonitrile, chloroform and finally ether (2 mL each) to afford 180 mg (80%) of light tan solid, m.p. 185-186 ° (dec.). NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD) δ 8.05-6.95 (m, 8H); 5.30 (s, 2H); 4.50 (s, 2H); 3.60 (t, 2H, 5Hz); 3.25 (s, 3H); 2.8 (t, 2H, 5Hz). Mass Spec. Calcd. for C<sub>22</sub>H<sub>18</sub>ClN<sub>3</sub>O<sub>3</sub> (M-2H<sub>2</sub>O): 407.1037. Found: 407.1031.

Example 56Preparation of 4-Chloro-5-hydroxymethyl-2-(2-methoxyethyl)-1-[4-((N-trifluoromethanesulfonyl)-anthranilamido)benzyl]imidazole

5

This compound was prepared according to the procedure described in Example 54, Part F. From 1-(4-aminobenzyl)-4-chloro-5-hydroxymethyl-2-(2-methoxyethyl)imidazole (200 mg, 0.68 mmol), N-(trifluoromethanesulfonyl)anthranoyl chloride (190 mg, 0.68 mmol) and sodium bicarbonate (280 mg, 3.3 mmol) in acetonitrile (5 mL) was obtained 300 mg (81%) of tan solid after chromatography (silica gel, 20 g; EtOAc/EtOH, 20:1); m.p. 75-95° (slow dec.); one spot by TLC. NMR (200 MHz, CDCl<sub>3</sub>/CD<sub>3</sub>OD)  $\delta$  8.00-6.80 (m, 8H); 5.15 (s, 2H); 4.45 (s, 2H); 3.60 (t, 2H, 5Hz); 3.15 (s, 3H); 2.75 (t, 2H, 5Hz).

10

The following compounds listed in Table 4 were prepared by the procedures described in Examples 54, Parts D, E and 54, Part F or 55, Part F.

15

20

25

30


35

40

45














50

55

Table 4

<u>Ex. No.</u>	<u>R</u>	<u>R<sup>6</sup></u>	<u>MP(°C)</u>
57		ethyl	(amorphous solid) <sup>a</sup>
58		i-propyl	(amorphous solid) <sup>b</sup>
59		n-butyl	(amorphous solid) <sup>c</sup>
60		n-pentyl	(amorphous solid) <sup>d</sup>
61			(amorphous solid) <sup>e</sup>
62		ethyl	188-189.5 (free acid)

Table 4 (continued)

Ex. No.	R	R <sup>6</sup>	MP (°C)
5 63		n-propyl	181.5-183 (free acid)
10 64		n-butyl	188.5-189.5 (Na+ salt)
15 65		n-pentyl	170.5-171.5
20 66		n-hexyl	171-171.5
25 67		n-heptyl	181-182
30 68			
35 69			
40 70			150-152
45 71			175-177



a NMR  $\delta$  8.05 (d, 1H); 7.62 (d, 2H); 7.52 (d, 1H); 7.30 (t, 1H); 7.17 (m, 3H); 6.93 (m, 2H); 5.13 (s, 2H); 2.61 (quart., 2H); 1.15 (t, 3H).

b NMR  $\delta$  8.04 (d, 1H); 7.63 (d, 2H); 7.51 (d, 1H); 7.28 (t, 1H); 7.13 (m, 3H); 6.89 (m, 2H); 5.14 (s, 2H); 3.11 (sept., 1H); 1.11 (d, 6H).

c NMR  $\delta$  8.05 (d, 1H); 7.64 (d, 2H); 7.52 (d, 1H); 7.30 (t, 1H); 7.17 (m, 3H); 6.92 (m, 2H); 5.15 (s, 2H); 2.66 (t, 2H); 1.53 (quint., 2H); 1.28 (sext., 2H); 0.83 (t, 3H).

d NMR  $\delta$  8.07 (d, 1H); 7.68 (d, 2H); 7.52 (m, 2H); 7.30 (m, 4H); 6.93 (t, 1H); 5.29 (s, 2H); 2.83 (t, 2H); 1.56 (m, 2H); 1.24 (m, 4H); 0.82 (t, 3H).

e NMR  $\delta$  8.03 (d, 1H); 7.61 (d, 2H); 7.51 (d, 1H); 7.28 (t, 1H); 7.10 (m, 3H); 6.91 (t, 1H); 6.78 (s, 1H); 5.09 (s, 2H); 2.46 (d, 2H); 1.62 (m, 6H); 0.99 (m, 5H).

#### Example 72

##### PART A: Preparation of 5-Hydroxymethyl-2-mercapto-1-(4-nitrobenzyl)imidazole

A mixture of 4-nitrobenzylamine hydrochloride (75 g, 0.40 mol), 1,3-dihydroxyacetone dimer (32.1, 0.17 mol) and potassium thiocyanate (51.9 g, 0.53 mol) in n-butanol (250 mL) and glacial acetic acid (40 mL) was stirred vigorously at room temperature for 48 hours. The mixture was suction filtered and the solid was washed thrice with water (300 mL) and thrice with ether (300 mL) before being dried overnight under vacuum to give 70.9 g (75%) of a yellow tan powder; m.p. 214-215° (dec.). NMR (200 MHz, DMSO- $d_6$ )  $\delta$  12.25 (s, 1H; absent in D<sub>2</sub>O shake); 8.20 (d, 2H, 8Hz); 7.40 (d, 2H, 8Hz); 6.90 (s, 1H); 5.40 (s, 2H); 5.25 (t, 1H, 5Hz; absent in D<sub>2</sub>O shake); 4.15 (d, 2H, 5Hz; s in D<sub>2</sub>O shake). Mass Spec. 265.

##### PART B: Preparation of 5-Hydroxymethyl-2-methylthio-1-(4-nitrobenzyl)imidazole

An ethanolic solution of sodium ethoxide was prepared by the gradual addition of sodium hydride (0.70 g of 60% NaH in mineral oil, 17.6 mmol) to absolute ethanol (150 mL). To this 5-hydroxymethyl-2-mercapto-1-(4-nitrobenzyl)imidazole (3.9 g, 14.7 mmol) was added and after being stirred 5-10 minutes, iodomethane (2.5 g, 1.1 mL, 17.6 mmol) was added. After being stirred 3 hours at room temperature, the mixture was concentrated on a rotary evaporator and the residue was partitioned between ethyl acetate (500 mL) and water (250 mL). The aqueous phase was further extracted with ethyl acetate (250 mL) and the combined organic phases were washed with water (150 mL), saturated aqueous sodium chloride (150 mL), dried (MgSO<sub>4</sub>), filtered and concentrated to leave 4.1 g of yellow-brown solid. Recrystallization from ethyl acetate

gave 2.6 g (64%) of light yellow-brown powder; m.p. 160-162°. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.20 (d, 2H, 7Hz); 7.30 (d, 2H, 7Hz); 6.95 (s, 1H); 5.40 (s, 2H); 5.20 (t, 1H, 5Hz; absent in D<sub>2</sub>O shake); 4.40 (d, 3H, 5Hz; s in D<sub>2</sub>O shake); 3.40 (s, 2H; monohydrate; δ 3.5 in D<sub>2</sub>O); 2.45 (s, 3H). Mass Spec. 279.

5 PART C: Preparation of 1-(4-Aminobenzyl)-5-hydroxymethyl-2-(methylthio)imidazole

This compound was prepared according to the procedure described in Example 54, Part E, from 5-hydroxymethyl-2-methylthio-1-(4-nitrobenzyl)imidazole (21 g, 75.2 mmol) and iron powder (75 g, 1.3 mol) there was obtained 13.5 g (72%) of a yellow hygroscopic solid. NMR (200 MHz, CDCl<sub>3</sub>) δ 6.90 (s, 1H); 6.85-6.45 (q, 4H, 5Hz, 51Hz); 5.10 (s, 2H); 4.40 (s, 2H); 2.40 (s, 3H). Mass Spec. 249.

PART D: Preparation of 1-[4-(2-Carboxybenzamido)benzyl]-5-hydroxymethyl-2-(methylthio)imidazole

This compound was prepared according to the procedure described in Example 55, Part F, though in this case the reaction was run in chloroform and the filtered product was washed with chloroform and ether. From 1-(4-aminobenzyl)-5-hydroxymethyl-2-(methylthio)imidazole (323 mg, 1.3 mmol) and phthalic anhydride (192 mg, 1.3 mmol) there was obtained 488 mg (95%) of the title compound as a yellow powder; m.p. 115-118° (dec.). NMR (200 MHz, CDCl<sub>3</sub>/DMSO-d<sub>6</sub>) δ 9.80 (s, 1H); 8.00-6.85 (m, 9H); 5.20 (s, 2H); 4.40 (s, 2H); 2.50 (s, 3H). Mass Spec. 379 (M-H<sub>2</sub>O).

20 Example 73

Preparation of 1-[4-(2-Carboxybenzamido)benzyl]-5-hydroxymethyl-2-methoxyimidazole

25 By repeating Example 72, Parts C and D, but substituting 5-hydroxymethyl-2-methoxy-1-(4-nitrobenzyl)imidazole as starting material in Part C, the compound 1-[4-(2-carboxybenzamido)benzyl]-5-hydroxymethyl-2-methoxyimidazole can be prepared.

30 Example 74

**PART A: Preparation of *trans*-2-(Trifluoromethanesulfonamido)cyclohexanecarboxylic acid**

35 Ethyl *trans*-2-(trifluoromethanesulfonamido) cyclohexanecarboxylate was synthesized from ethyl *trans*-2-aminocyclohexanecarboxylate [E. J. Moriconi and P. H. Mazzocchi, *J. Org. Chem.*, **31**, 1372 (1966)] by the procedure described in Example 21. The crude product (2.59 g, 8.55 mmol, 1 eq) was then hydrolyzed by refluxing in 1.00N NaOH (26.5 mL, 26.5 mmol, 3.1 eq) overnight under N<sub>2</sub>. Water (100 mL) was then added and the pH adjusted to 3 using 1N HCl. The aqueous was extracted with ethyl acetate (3 x 100 mL), the organic layers dried (MgSO<sub>4</sub>) and concentrated to yield a crystalline white solid which was recrystallized from *n*-butyl chloride. Obtained 1.71 g of product; m.p. 114.5-118.5°. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 12.47 (bs, 1H); 9.52 (bs, 1H); 2.35 (d of d of d, 1H, J = 10, 10, 4Hz); 2.10-1.13 (m, 9H). Anal. Calcd. for C<sub>8</sub>H<sub>12</sub>F<sub>3</sub>NO<sub>4</sub>S: C, 34.91; H, 4.39; N, 5.09. Found, C, 34.73; H, 4.22; N, 5.04.

45 **PART B: Preparation of Methyl 2-butyl-4-chloro-1-[4-(*trans*-2-(trifluoromethanesulfonamido)cyclohexanecarboxamido)benzyl]imidazole-5-acetate and methyl 2-butyl-4-chloro-1-[4-(*cis*-2-(trifluoromethanesulfonamido)cyclohexanecarboxamido)benzyl]imidazole-5-acetate**

55 *trans*-2-(Trifluoromethanesulfonamido)cyclohexanecarboxylic acid (500 mg, 1.82 mmol, 1 eq) and thionyl chloride (2.30 mL, 31.5 mmol, 17.3 eq) were mixed and refluxed for 2 hours. The excess thionyl chloride was removed in vacuo and the residue suspended in toluene. The toluene was removed by rotary

evaporation and the procedure repeated to remove traces of thionyl chloride. Final rotary evaporation yielded 460 mg of white crystalline acid chloride product which was used without further purification (IR 1789  $\text{cm}^{-1}$ ).

Methyl 2-butyl-4-chloro-1-(4-aminobenzyl)imidazole-5-acetate (530 mg, 1.57 mmol, 1 eq), trans-2-(trifluoromethanesulfonamido)cyclohexanoyl chloride (460 mg, 1.57 mmol, 1 eq) and sodium bicarbonate (400 mg, 4.70 mmol, 3 eq) were mixed and stirred in chloroform (20 mL) overnight. Water (100 mL) was then added, and the pH adjusted to 4 with 1N HCl. The aqueous was extracted with methylene chloride (3 x 100 mL) and the organic layers dried and concentrated. Gradient flash chromatography of the residue in 60:40 hexane/ethyl acetate to 100% ethyl acetate over silica gel yielded two isomers; both of which were isolated as glasses. The faster eluting product being the minor cis isomer (170 mg) while the slower being the major trans isomer (520 mg).

trans-Isomer; NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.18 (s, 1H); 7.42 (d, 2H,  $J = 10\text{Hz}$ ); 6.84 (d, 2H,  $J = 10\text{Hz}$ ); 6.47 (bd, 1H,  $J = 8\text{Hz}$ ); 5.07 (s, 2H); 3.72 (m, 1H); 3.57 (s, 3H); 3.47 (s, 2H); 2.53 (t, 2H, 7Hz); 2.24-1.12 (m, 13Hz); 0.82 (t, 3H,  $J = 7\text{Hz}$ ). Anal. Calcd. for  $\text{C}_{25}\text{H}_{32}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : C, 50.63; H, 5.44; N, 9.45. Found: C, 50.64; H, 5.44; N, 9.16. Mass Calcd. for  $\text{C}_{25}\text{H}_{32}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : 592.1734. Found: 592.1731.

cis-Isomer; NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.94 (s, 1H); 7.42 (d, 2H,  $J = 10\text{Hz}$ ); 6.88 (d, 2H,  $J = 10\text{Hz}$ ); 6.52 (bd, 2H,  $J = 8\text{Hz}$ ); 5.11 (s, 2H); 3.75 (m, 1H); 3.63 (s, 3H); 3.48 (s, 2H); 2.56 (t, 2H, 7Hz); 2.29-1.25 (m, 13H); 0.86 (t, 3H,  $J = 7\text{Hz}$ ). Anal. Calcd. for  $\text{C}_{25}\text{H}_{32}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : C, 50.63; H, 5.44. Found: C, 49.87; H, 5.65. Mass Calcd. for  $\text{C}_{25}\text{H}_{32}\text{ClF}_3\text{N}_4\text{O}_5\text{S}$ : 592.1734. Found: 592.1689.

## Example 75

### PART A: Preparation of 2-Butyl-4,5-dicyanoimidazole

Ethyl pentanimidate hydrochloride (42.66 g, 257.8 mmol, 1 eq), diaminomaleonitrile (27.90 g, 258.1 mmol, 1 eq) and pyridine (400 mL) were mixed and refluxed for 48 hours under  $\text{N}_2$ . The solvent was removed by rotary evaporation.

The residue was taken up in ethyl acetate and filtered through a pad (3" x 4") of florisil. The solvent was removed in vacuo and the residue flash chromatographed in 60:40 hexane/ethyl acetate over silica gel to give 16.59 g of a yellow solid which was used in the following step without further purification. An analytical sample was prepared by recrystallizing the crude product (3.03 g) from ether/hexane to give 1.55 g of yellow crystals; m.p. 108.0-109.0°. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  2.86 (t, 2H,  $J = 7\text{Hz}$ ); 1.77 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.41 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.98 (t, 3H,  $J = 7\text{Hz}$ ). Anal. Calcd. for  $\text{C}_9\text{H}_{10}\text{N}_4$ : C, 62.05; H, 5.79; N, 32.16. Found: c, 62.28; H, 5.81; N, 32.22. Mass spectrum shows M-H peak. Mass Calcd. for  $\text{C}_9\text{H}_{10}\text{N}_4\text{-H}$ : 173.0827. Found: 173.0785.

### PART B: Preparation of 2-Butyl-4,5-dicyano-1-(4-nitrobenzyl)imidazole

2-n-Butyl-4,5-dicyano-1-(4-nitrobenzyl)imidazole was prepared from 2-n-butyl-4,5-dicyanoimidazole by the procedure in Example 1, Part A using 4-nitrobenzyl bromide as the alkylating agent. The product was obtained as an oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.29 (d, 2H,  $J = 10\text{Hz}$ ); 7.29 (d, 2H,  $J = 10\text{Hz}$ ); 5.36 (s, 2H); 2.67 (t, 2H,  $J = 7\text{Hz}$ ); 1.70 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.36 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.86 (t, 3H,  $J = 7\text{Hz}$ ). Mass Calcd. for  $\text{C}_{16}\text{H}_{15}\text{N}_5\text{O}_2$ : 309.1225. Found: 309.1211.

### PART C: Preparation of 1-(4-Aminobenzyl)-2-butyl-4,5-dicyanoimidazole

A mixture of 2-butyl-4,5-dicyano-1-(4-nitrobenzyl)imidazole (2.00 g, 6.5 mmol, 1 eq), tin dichloride dihydrate (7.30 g, 32.3 mmol, 5 eq) and ethanol (13 mL) was stirred and heated at 70° for 50 minutes. The reaction was terminated by pouring the mixture onto ice and adjusting the pH to 8 with saturated aqueous  $\text{NaHCO}_3$ . The aqueous mixture was extracted with ethyl acetate (3 x 100 mL) and the organic layers were dried ( $\text{MgSO}_4$ ) and concentrated to give a thick amber oil. This oil was flash chromatographed over silica gel in 75:25 to 70:30 hexane/ethyl acetate yielding 330 mg of yellow crystals; m.p. 99.0-103.5°. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  6.97 (d, 2H,  $J = 10\text{Hz}$ ); 6.68 (d, 2H,  $J = 10\text{Hz}$ ); 5.10 (s, 2H); 2.69 (t, 2H,  $J = 7\text{Hz}$ ); 1.72 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.38 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.91 (t, 3H,  $J = 7\text{Hz}$ ). Mass Calcd. for  $\text{C}_{16}\text{H}_{17}\text{N}_5$ : 279.1483. Found: 279.1489.

PART D: Preparation of 2-Butyl-4,5-dicyano-1-[4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl]-imidazole

The title compound was prepared by the procedure described in Example 30 starting with 1-(4-aminobenzyl)-2-butyl-4,5-dicyanoimidazole and N-(trifluoromethanesulfonyl)anthranilic acid chloride. NMR (200 MHz, CDCl<sub>3</sub> + DMSO-d<sub>6</sub>) δ 7.98 (d, 1H, J = 7Hz); 7.32 (d, 2H, J = 7Hz); 7.62 (d, 1H, J = 7Hz); 7.47 (d of d, 1H, J = 7,7Hz); 7.24 (d of d, 1H, J = 7,7Hz); 7.15 (d, 2, J = 7,7Hz); 5.32 (s, 2H); 2.75 (t, 2H, J = 7Hz); 1.70 (t of t, 2H, J = 7,7Hz); 1.37 (t of q, 2H, J = 7,7Hz); 0.92 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>24</sub>H<sub>21</sub>F<sub>3</sub>N<sub>5</sub>O<sub>3</sub>S: 503.1348. Found: 530.1343.

Example 76PART A: Preparation of Methyl 1-[4-(N-benzylamino)benzyl]-2-butyl-4-chloroimidazole-5-acetate

A mixture of methyl 1-(4-aminobenzyl)-2-butyl-4-chloroimidazole-5-acetate (1.00 g, 3.0 mmol, 1 eq), benzaldehyde (0.30 mL, 3.0 mmol, 1 eq), 4A\* powdered molecular sieves (enough to make a slurry) and 40 mL THF was stirred overnight. The next day, more benzaldehyde (0.2 mL) and acidic Al<sub>2</sub>O<sub>3</sub> (activity 1, 1g) were added and the slurry stirred another 24 hours. The solids were filtered and the solvent from the filtrate removed in vacuo. The residue was dissolved in methanol (10 mL) and sodium cyanoborohydride was added (0.19 g, 3.0 mmol, 1 eq). The mixture was stirred for 24 hours, after which the solvent was removed in vacuo to yield a green oil which was flash chromatographed over silica gel in 70:30 hexane/ethyl acetate to give 740 mg of product as an oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.42 - 7.24 (m, 5H); 6.74 (d, 2H, J = 7Hz); 6.56 (d, 2H, J = 7Hz); 4.98 (s, 2H); 4.31 (s, 2H); 3.61 (s, 3H); 3.48 (s, 2H); 2.60 (t, 2H, J = 7Hz); 1.67 (t of t, 2H, J = 7,7Hz); 1.35 (t of q, 2H, J = 7,7Hz); 0.89 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>24</sub>H<sub>28</sub>ClN<sub>3</sub>O<sub>2</sub>: 425.1868. Found: 425.1853.

PART B: Preparation of Methyl 2-butyl-1-[4-(N-benzyl-N-(2-(trifluoromethanesulfonamido)benzoyl)amino)-benzyl]-4-chloroimidazole-5-acetate

The title compound was prepared from the compound of Part A by the procedure described in Example 30. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.59 (d, 1H, J = 10Hz); 7.33-7.16 (m, 6H); 6.89 (d, 2H, J = 10Hz); 6.76 (d, 2H, J = 10Hz); 6.93-6.70 (m, 2H); 5.12 (s, 2H); 5.02 (s, 2H); 3.55 (s, 3H); 3.39 (s, 2H); 2.47 (t, 2H, J = 7Hz); 1.64 (t of t, 2H, J = 7,7Hz); 1.30 (t of q, 2H, J = 7,7Hz); 0.88 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>32</sub>H<sub>32</sub>ClF<sub>3</sub>N<sub>4</sub>O<sub>5</sub>S: C, 56.76; H, 4.76; N, 8.27. Found: C, 56.64; H, 4.90; N, 7.98.

Example 77PART A: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[N-methyl-4-aminobenzyl]imidazole

1-(4-Aminobenzyl)-2-n-butyl-4-chloro-5-(methoxymethyl)imidazole (10.94 g) and ethyl formate (150 mL) were mixed and refluxed overnight. The excess ethyl formate was removed in vacuo and another 150 mL added and the mixture was refluxed overnight again. The excess ethyl formate was removed in vacuo and the residue flash chromatographed over silica gel in 1:1 hexane/ethyl acetate to yield 9.52 g of a golden oil which slowly crystallized after several days. This oil (9.40 g, 28 mmol, 1 eq) was dissolved in THF and to it LAH (1M in THF, 84.0 mL, 84 mmol, 3 eq) was slowly added via syringe under N<sub>2</sub>. After stirring for 1 h, the mixture was worked up as described in Fieser and Fieser, V.1 pg. 584 (Steinhardt procedure) to yield 8.47 g of an orange oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 6.84 (d, 2H, J = 10Hz); 6.55 (d, 2H, J = 10Hz); 5.02 (s, 2H); 4.26 (s, 2H); 3.27 (s, 3H); 2.81 (s, 3H); 2.58 (t, 2H, J = 7Hz); 1.67 (t of t, 2H, J = 7,7Hz); 1.35 (t of q, 2H, J = 7,7Hz); 0.87 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>17</sub>H<sub>24</sub>ClN<sub>3</sub>O: C, 63.44; H, 7.52; N, 13.06. Found: C, 63.60; H, 7.61; N, 12.86.

PART B: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[4-(N-methyl-2-carboxy-3,6-dichlorobenzamido)benzyl]imidazole

2-n-Butyl-4-chloro-5-methoxymethyl-1-[N-methyl-4-aminobenzyl]imidazole (2.00 g, 6.2 mmol, 1 eq) and 3,6-dichlorophthalic anhydride (1.35 g, 6.2 mmol, 1 eq) were reacted by the procedure described in Example 2, Part D to give 2.37 g of a white powder; m.p. 120.0-123.5°. The NMR shows a 7:2 mixture of conformers in DMSO-d<sub>6</sub>. NMR (200 MHz, DMSO-d<sub>6</sub>) δ (major conformer only) 14.25 (m, 1H); 7.76-6.85 (m,



6H); 5.09 (s, 2H); 4.18 (s, 2H); 3.06 (s, 3H); 2.37 (t, 2H, J = 7Hz); 1.38 (t of t, 2H, J = 7,7Hz); 1.21 (t of q, 2H, J = 7,7Hz); 0.77 (t, 3H, J = 7Hz). Anal. Calcd. for  $C_{25}H_{26}Cl_3N_3O_4$ : C, 55.72; H, 4.86; Cl, 19.74. Found: C, 55.48; H, 4.88; Cl, 19.77.

## 5 Example 78

### PART A: Preparation of 2-n-Butyl-1-(4-carbomethoxybenzyl)-4-chloro-5-(methoxymethyl)imidazole

2-Butyl-4-chloro-5-hydroxymethyl-1-(4-carboxybenzyl)imidazole (17.6 g), methanol (500 mL) and conc. sulfuric acid (50 mL) were mixed and refluxed overnight. Potassium carbonate (100 g) was then carefully added to the solution which was cooled over ice. The reaction mixture was then stirred for 2.5 hours. The solvent was removed in vacuo and the residue dissolved in water (1 L). This aqueous mixture was extracted with ethyl acetate (3 x 400 mL). The organic layers were combined, dried ( $MgSO_4$ ) and the solvent removed in vacuo to yield 15.2 g of an oil. NMR (200 MHz,  $DMSO-d_6$ )  $\delta$  8.46 (d, 2H, J = 9Hz); 7.68 (d, 2H, J = 9Hz); 5.82 (s, 2H); 4.80 (s, 2H); 4.37 (s, 3H); 3.66 (s, 3H); 3.02 (t, 2H, J = 7Hz); 2.01 (t of t, 2H, J = 7,7Hz); 1.77 (t of q, 2H, J = 7,7Hz); 1.33 (t, 3H, J = 7Hz). Anal. Calcd. for  $C_{18}H_{23}ClN_2O_3$ : C, 61.62; H, 6.61; N, 7.99. Found: C, 61.79; H, 6.78; N, 7.82.

### PART B: Preparation of 2-n-Butyl-1-(4-carboxybenzyl)-4-chloro-5-(methoxymethyl)imidazole

2-n-Butyl-1-(4-carbomethoxybenzyl)-4-chloro-5-(methoxymethyl)imidazole (15.2 g, 43.3 mmol, 1 eq), 0.5 N KOH in methanol (130 mL, 65.0 mmol, 1.5 eq), water (10 mL) and methanol (50 mL) were mixed and refluxed for 4 hours. The solvent was removed in vacuo and the residue dissolved in water (300 mL). The pH was adjusted to 4 with conc. HCl and this aqueous mixture extracted with ethyl acetate (3 x 300 mL). The organic layers were combined, dried ( $MgSO_4$ ), the solvent removed in vacuo and the crude residue recrystallized from hexane/butyl chloride to yield 9.6 g of white solid; m.p. 126.5-127.5°. NMR (200 MHz,  $DMSO-d_6$ )  $\delta$  12.95 (bs, 1H); 7.93 (d, 2H, J = 9Hz); 7.16 (d, 2H, J = 9Hz); 5.30 (s, 2H); 4.31 (s, 2H); 3.19 (s, 3H); 2.50 (t, 2H, J = 7Hz); 1.49 (t of t, 2H, J = 7,7Hz); 1.24 (t of q, 2H, J = 7,7Hz); 0.80 (t, 3H, J = 7Hz). Anal. Calcd. for  $C_{17}H_{21}ClN_2O_3$ : C, 60.62; H, 6.29; N, 8.32. Found: C, 60.89; H, 6.10; N, 8.03.

### PART C: Preparation of 2-n-Butyl-1-[4-(N-(2-carboxyphenyl)carboxamido)benzyl]-4-chloro-5-methoxymethyl)imidazole


2-n-Butyl-1-(4-carboxybenzyl)-4-chloro-5-(methoxymethyl)imidazole (6.00 g, 17.8 mmol, 1 eq), thionyl chloride (13.0 mL, 178 mmol, 10 eq) and chloroform (100 mL) were mixed and refluxed for 6 h. The solvent was removed in vacuo, and the residue dissolved in toluene. The solvent was removed on the rotary evaporator and the evaporation from toluene repeated to remove all of the thionyl chloride. This yielded 6.0 g of acid chloride as an amber gum. IR 1776, 1745  $cm^{-1}$ . Anthranilic acid (0.737 g, 5.36 mmol, 1 eq) was dissolved in 1.000 N NaOH (10.75 mL, 10.7 mmol, 2 eq) and water (100 mL) and cooled over ice. The aforementioned acid chloride (1.91 g, 5.36 mmol, 1 eq) dissolved in THF (50 mL) was slowly added via a dropping funnel to the stirred and cooled anthranilic acid solution. The following day, more anthranilic acid (74 mg, 0.536 mmol, 0.1 eq) was added to bring the reaction to completion. After 1.5 h, the solution was acidified to pH = 5 with 1N HCl and extracted with ethyl acetate (1 x 100mL). The ethyl acetate layer was then washed with water (3 x 50 mL), and brine (1 x 50 mL), dried ( $MgSO_4$ ) and the solvent removed in vacuo to yield 2.28 g of a brown glass. This glass was dissolved in a minimum amount of ethyl acetate and dicyclohexylamine ("DCHA", 1 eq) was added thereto. The salt did not crystallize and therefore was flash chromatographed over silica gel starting in 100% ethyl acetate and finishing in 1:1 ethyl acetate/isopropanol to yield 1.44 g of an oil. This oil was dissolved in ethyl acetate (100 mL) and a minimum of methanol, and washed with 1N HCl (2x50mL). The ethyl acetate layer was dried ( $MgSO_4$ ) and the solvent removed in vacuo to yield 0.52 g of an amber oil. NMR (200 MHz,  $CDCl_3$ )  $\delta$  12.53 (s, 1H); 8.91 (d, 1H, J = 8Hz); 8.23 (d, 1H, J = 7Hz); 8.08 (d, 3H, J = 7Hz); 7.62 (t, 1H, J = 6Hz); 7.11 (t, 2H, J = 7Hz); 5.30 (s, 2H); 4.30 (s, 2H); 3.30 (s, 3H); 2.72 (t, 2H, J = 7Hz); 1.72 (t of t, 2H, J = 7,7Hz); 1.31 (t of q, 2H, J = 7,7Hz); 0.83 (t, 3H, J = 7Hz). Anal. Calcd. for  $C_{25}H_{25}ClN_3O_4 \cdot (H_2O)_{1.5}$ : C, 59.81; H, 5.85; Cl, 7.36. Found: C, 59.78; H, 6.38; Cl, 7.51.

Examples 79-84 in Table 5 were made or could be made by procedures described in Example 78 and by methods familiar to one skilled in the art.

Table 5

5


10



15

<u>Ex. No.</u>	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>R</u>	<u>MP (°C)</u>
79	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		( > 300 ) <sup>a</sup>

20



25

30

35






40

45

50

55

Table 5 (continued)

Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R	MP (°C)
80	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		(glass) <sup>b</sup>
81	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		(white solid) <sup>c</sup>
82	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>	 L-isomer	149-152
83	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>	 L-isomer	134.5-136.0
84	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		

a NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.01 (d, 2H, J= 7Hz); 7.17 (d, 2H, J= 7Hz); 5.31 (s, 2H); 4.27 (s, 2H); 3.18 (s, 3H); 2.50 (t, 2H, J= 7Hz); 1.50 (t of t, 2H, J= 7,7Hz); 1.21 (t of q, 2H, J= 7,7Hz); 0.80 (t, 3H, J= 7Hz).



b NMR (200 MHz, CDCl<sub>3</sub>) δ 11.52 (s, 1H) 8.55 (d, 1H, J= 7 Hz); 8.0 (d, 2H, J= 7Hz) 7.41 (t, 1H, J= 7Hz); 7.14 (d, 2H, J= 7Hz); 7.04 (d, 1H, J= 7Hz); 5.30 (s, 2H); 4.25 (s, 2H); 3.30 (s, 3H); 2.73 (t, 2H, J= 7Hz); 2.60 (s, 3H); 1.68 (t of t (br), 2H); 1.29 (t of q, 2H, J= 7,7Hz); 0.81 (t, 3H, J= 7Hz).

c NMR (200 MHz, CDCl<sub>3</sub>) δ 12.05 (s, 1H); 8.88 (d, 1H, J= 7Hz); 8.23 (d, 2H, J= 8Hz); 8.11 (d, 1H, J= 7Hz); 7.51 (t, 1H, J= 7Hz); 7.25-7.11 (m, 3H); 5.29 (s, 2H); 4.31 (s, 2H); 3.29 (s, 3H); 2.62 (t, 2H, J= 7Hz); 1.63 (t of t, 2H, J= 7,7Hz); 1.26 (t of q, 2H, J= 7,7Hz); 0.75 (t, 3H, J= 7Hz) IR: 1621, 753 cm<sup>-1</sup>.

## Example 85

## PART A: Preparation of Methyl 4'-methylbiphenyl-3-carboxylate

To a stirred solution of 25.2 g of methyl 3-iodobenzoate and 21.0 g of 4-iodotoluene at 180-190° under nitrogen was added 30.3 g of copper powder portionwise over 1 hour. When approximately one-third of the copper had been added, the reaction initiated and the temperature increased spontaneously to 240°. The mixture was allowed to cool to 210°, then was held at 210° during the addition of the remaining copper and for an additional hour. The mixture was allowed to cool to room temperature and was filtered employing benzene as solvent; the resulting filtrate was concentrated in vacuum to provide the crude product. Column chromatography on silica gel (elution = 50-100% benzene/hexane) followed by distillation furnished 7.60 g of methyl 4'-methylbiphenyl-3-carboxylate [bp: 114-115°C (0.025 torr)] as a colorless oil; NMR (200 MHz, CDCl<sub>3</sub>): δ 8.27 (br s, 1H); 7.99 (d, 1H); 7.77 (d, 1H); 7.50 (t, 1H); 7.39 (A<sub>2</sub>B<sub>2</sub>, 4H); 3.94 (s, 3H); 2.41 (s, 3H). The following methylbiphenyl starting materials were prepared employing the above procedure.

		NMR (200 MHz, CDCl <sub>3</sub> )
20	a) 	δ 7.78 (d, 1H); 7.46 (d, 1H); 7.35 (t, 2H); 7.19 (s, 4H); 3.64 (s, 3H); 2.37 (s, 3H)
25	b) 	δ 7.80 (d of d, 1H); 7.57 (t of d, 1H); 7.41 (m, 2H); 7.19 (s, 4H); 2.37 (s, 3H)
30		

Alternatively methyl 4'-methylbiphenyl-2-carboxylate (compound a) and tert-butyl 4'-methylbiphenyl-2-carboxylate can be prepared by chemistry described by A. Meyers via the following five-step procedure.

## Step 1: Preparation of 2-Methoxybenzoyl chloride

To 30 g of 2-anisic acid in 500 mL of round-bottom flask was added dropwise 50 mL of thionyl chloride. After all of the thionyl chloride was added the reaction mixture was stirred at room temperature for 18 hours. Excess thionyl chloride was then distilled off by water aspirator and the remaining liquid was vacuum distilled (82°/0.7 mm Hg). Desired 2-methoxybenzoyl chloride was obtained as a colorless liquid, 32 g.

## Step 2: Preparation of 4,4-Dimethyl-2-(2-methoxyphenyl)oxazoline

20 g of 2-Amino-2-methyl-1-propanol was dissolved in 100 mL of methylene chloride and the mixture was cooled with ice. Meanwhile, 17 g of 2-methoxybenzoyl chloride prepared from Step 1 was placed in a dropping funnel, diluted with 50 mL of methylene chloride and added dropwise. After the addition of the acid chloride, the cooling ice bath was removed and the reaction mixture was stirred at room temperature for 2 hours.

The reaction mixture was concentrated to remove the solvent and the solids obtained were triturated with water, collected by filtration and washed with water. Thus obtained solids were dried in vacuo to give a colorless light solid, 20.5 g.

The solid was placed in 200 mL of round-bottom flask and 22 mL of thionyl chloride was added slowly to the solid without any solvent. At the beginning of the addition the reaction was vigorous but was controllable. After the addition of thionyl chloride was complete, the yellow reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 200 mL of ether and the resulting solids were collected and washed with ether. The solids were dissolved in 100 mL of water and the pH of the



solution was adjusted to 10 by adding 1N NaOH. The aqueous solution was extracted into ether 3 times. The combined ether extracts were dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated to give the desired product as a white solid, 18 g, m.p. 70-72°.

5 Step 3: Preparation of 2-(4'-Methylbiphenyl-2-yl)-4,4-dimethyloxazoline

4-Methylphenyl Grignard reagent was prepared from 2.5 g of magnesium and 13 mL of 4-bromotoluene in 200 mL of anhydrous THF. The Grignard reagent was added to 10 g of the product from Step 2 in 100 mL of anhydrous THF and the reaction mixture was stirred at room temperature for 2 hours. The reaction  
10 mixture was concentrated and the residue was treated with 200 mL of saturated  $\text{NH}_4\text{Cl}$  solution and the mixture was stirred at room temperature for 30 minutes. The aqueous solution was then extracted with ethyl acetate. The crude product obtained upon concentration of the ethyl acetate extracts were purified by flash column chromatography (silica gel, hexane:ethyl acetate = 2:1) to give the desired compound as a colorless liquid, 11.8 g.

15 Step 4: Preparation of 4'-Methylbiphenyl-2-carboxylic acid

A mixture of 10 g of the product from Step 3 and 200 mL of 4.5 N HCl was refluxed for 12 hours. During this period of time the desired compound was isolated as a brownish oil floating on the surface of  
20 the reaction medium. The reaction mixture was cooled to room temperature. The product which was oily initially began to solidify upon cooling. The product was extracted with ethyl ether. Upon concentration of the ether extract the desired product was obtained as a colorless solid, 7 g, m.p. 140-142°.

25 Step 5: Esterification of 4'-methylbiphenyl-2-carboxylic acid

Preparation of methyl 4'-methylbiphenyl-2-carboxylate

To 100 mL of methanol was added dropwise 5 mL of acetyl chloride with ice cooling. After stirring the mixture for 15 minutes, 5 g of the acid from Step 4 was added at once and the mixture was refluxed for 4  
30 hours. The reaction mixture was concentrated to remove the solvent and the desired methyl ester was obtained as a thick liquid, 5 g.

Preparation of tert-butyl 4'-methylbiphenyl-2-carboxylate

35 To a solution of 42.4 g of 4'-methylbiphenyl-2-carboxylic acid in 200 mL of methylene chloride at 0° was added dropwise 20 mL of oxalyl chloride. The reaction was allowed to warm to 25° and then was stirred at 25° for 3 hours. The solvent was removed in vacuo. The residue was dissolved in benzene, and the benzene then removed in vacuo to provide 46.1 g of crude acid chloride.




The acid chloride prepared above was dissolved in 600 mL of tetrahydrofuran. To this solution at 0°  
40 was added 26.0 g of potassium t-butoxide portionwise such that the reaction temperature did not exceed 15-20°C. The resulting mixture was then allowed to stir at 25°C for 1 hour. The reaction mixture was poured into water, and the resulting emulsion was extracted with diethyl ether. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Distillation provided 49.5 g of tert-butyl 4'-methylbiphenyl-2-carboxylate (bp 115-120°/0.05 torr). NMR (200 MHz,  
45  $\text{CDCl}_3$ ):  $\delta$  7.73 (d of d, 1H), 7.46-7.27 (m, 3H); 7.18 (s, 4H); 2.40 (s, 3H); 1.30 (s, 9H).

PART B: Preparation of Methyl 4'-bromomethylbiphenyl-3-carboxylate

50 A solution of 7.31 g of Methyl 4'-methylbiphenyl-3-carboxylate, 5.75 g of N-bromosuccinimide, 0.125 g of azo(bisisobutyronitrile), and 500 mL of carbon tetrachloride was refluxed for 3 hours. After cooling to room temperature the resulting suspension was filtered and then concentrated in vacuo to provide 9.90 g of crude methyl 4'-bromomethylbiphenyl-3-carboxylate which was used in a subsequent reaction without further purification; NMR (200 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.28 (s, 1H); 8.05 (d, 1H); 7.79 (d, 1H); 7.67-7.48 (m, 5H); 4.55 (s, 2H); 3.98 (s, 3H).

55

The following bromomethylbiphenyl intermediates were prepared employing the above procedure.

5	a) 	NMR (200 MHz, CDCl <sub>3</sub> )  δ 7.82 (d, 1H); 7.59-7.23 (m, 7H); 4.52 (s, 2H); 3.62 (s, 3H)
10		
15	b) 	δ 7.86 (d of d, 1H); 7.62 (t of d, 1H); 7.53-7.21 (m, 6H); 4.52 (s, 2H)
20		
25	c) 	δ 7.79 (d, 1H); 7.56-7.24 (m, 7H); 4.51 (s, 2H); 1.25 (s, 9H).
30		









PART C: Preparation of 1-[(3'-Carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole

To a suspension of 1.43 g of sodium methoxide in 20 mL of dimethylformamide at 25° was added a solution of 5.00 g of 2-butyl-4(5)-chloro-5(4)-hydroxymethyl imidazole in 15 mL of DMF. The resulting mixture was stirred at 25° for 0.25 hours, and then to this mixture was added dropwise a solution of 9.90 g of methyl 4'-bromomethylbiphenyl-3-carboxylate in 15 mL of DMF. Finally, the reaction mixture was stirred at 40° for 4 hours. After cooling to 25°, the solvent was removed in vacuo. The residue was dissolved in ethyl acetate, and this solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product contains two regioisomers, the faster moving one by TLC being the more potent isomer. Column chromatography on silica gel (elution: 10-25% ethyl acetate/benzene) afforded 3.85 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole (m.p. 162-163°), the regioisomer of higher R<sub>f</sub>; NMR (200 MHz, CDCl<sub>3</sub>) 8.24 (s, 1H); 8.03 (d, 1H); 7.76 (d, 1H); 7.52 (t, 1H); 7.33 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.27 (s, 2H); 4.52 (d, 2H); 3.93 (s, 3H); 2.60 (t, 2H); 1.89 (t, 1H); 1.67 (quint., 2H); 1.35 (sext., 2H); 0.88 (t, 3H).

PART D: Preparation of 1-[(3'-Carbomethoxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethylimidazole

A mixture of 1.00 g of 10% palladium/carbon and 1.00 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethyl imidazole in 20 mL of methanol was stirred at 25° for five minutes. Hydrogen gas was bubbled into the solution, and the mixture was stirred under H<sub>2</sub>(g) (1 atm.) at 25° for 3.5 hours. The mixture was filtered, and the resulting solution concentrated in vacuo. Column chromatography (elution: 0-5% methanol/chloroform) furnished 0.33 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl imidazole. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.20 (s, 1H); 7.98 (d, 2H); 7.65 (t, 1H); 7.41 (A<sub>2</sub>M<sub>2</sub>, 4H); 6.80 (s, 1H); 5.30 (s, 2H); 5.12 (t, 1H); 4.37 (d, 2H); 3.90 (s, 3H); 2.52 (t, 2H); 1.51 (quint., 2H); 1.27 (sext., 2H); 0.80 (t, 3H).

The following intermediates shown below were also prepared by the procedures described in Part C or Parts C and D above.

5						
10						
15						
20	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>		<u>MP (°C)</u>	
25	n-butyl	Cl	CH <sub>2</sub> OH		162-163	
30	n-butyl	Cl	CH <sub>2</sub> OH		(oil) <sup>a</sup>	
35	n-butyl	H	CH <sub>2</sub> OH		139-141	
40	n-butyl	I	CH <sub>2</sub> OH		125-126	
45	n-butyl	CH <sub>2</sub> OH Cl			116-118	
50	n-butyl	CH <sub>2</sub> OH Cl			122-124	
55						

a NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.82 (d of d, 1H);  
 7.58 (t of d, 1H); 7.44 (t of d, 1H); 7.35 (d  
 of d, 1H); 7.11 ( $\text{A}_2\text{B}_2$ , 4H); 5.21 (s, 2H);  
 4.46 (s, 2H); 2.59 (t, 2H); 1.60 (quint, 2H);  
 1.29 (sext., 2H); 0.82 (t, 3H).

PART E: Preparation of 1-[(3'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole


A solution of 0.30 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole in 16 mL of ethanol and 8 mL of 10% aqueous sodium hydroxide was refluxed for 5 hours. After cooling, the reaction mixture was filtered, and the solvent was removed in vacuo. The residue was dissolved in water, and the solution was acidified to pH 3.5 using hydrochloric acid. The precipitated solid was recovered by filtration and recrystallized from aqueous ethanol to furnish 0.24 g of 1-[(3'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole (m.p. 180-181°); NMR (200 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.26 (s, 1H); 8.04 (d, 1H); 7.77 (d, 1H); 7.52 (t, 1H); 7.36 ( $\text{A}_2\text{M}_2$ , 4H); 5.30 (s, 2H); 4.48 (s, 2H); 2.57 (t, 2H); 1.64 (quint., 2H); 1.34 (sext., 2H); 0.87 (t, 3H).






Example 86

PART A: Preparation of 1-[(3'-Carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole

A solution of 5.00 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole and 1.0 mL of conc. sulfuric acid in 200 mL of methanol was refluxed for 20 hours. After cooling, the solvent was removed in vacuo, and the residue was poured into saturated sodium bicarbonate solution. The resulting mixture was extracted with methylene chloride, and the combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography on silica gel (elution: 0-20% ethyl acetate/benzene) furnished 5.35 g of 1-[(3'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole; NMR (200 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.26 (t, 1H); 8.03 (d of t, 1H); 7.76 (d of t, 1H); 7.51 (t, 1H); 7.33 ( $\text{A}_2\text{M}_2$ , 4H); 5.20 (s, 2H); 4.31 (s, 2H); 3.94 (s, 3H); 3.27 (s, 3H); 2.59 (t, 2H); 1.68 (quint., 2H); 1.34 (sext., 2H); 0.87 (t, 3H).

The following intermediates were prepared or could be prepared using the above described procedure.



5	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>		<u>MP (°C)</u>
10	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		oil <sup>a</sup>
15	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		oil <sup>b</sup>
20	n-butyl	Cl			oil <sup>c</sup>

25 a -NMR (200 MHz, CDCl<sub>3</sub>) δ 7.82 (d, 1H, J = 7Hz);  
 7.50 (t, 1H, J = 7Hz); 7.38 (t, 1H, J = 7Hz);  
 7.30 (d, 1H, J = 7Hz); 7.26 (d, 2H, J = 10Hz);  
 7.00 (d, 2H, J = 10Hz); 5.14 (s, 2H); 4.32 (s,  
 30 2H); 3.63 (s, 3H); 3.28 (s, 3H); 2.60 (t, 2H,  
 J = 7Hz); 1.70 (t of t, 2H, J = 7.7Hz); 1.36 (t  
 of q, 2H, J = 7.7Hz); 0.89 (t, 3H, J = 7Hz).

35 b -NMR (200 MHz, CDCl<sub>3</sub>) δ 7.88 (d of d, 1H);  
 7.63 (t of d, 1H); 7.51 (t of d, 1H); 7.41 (d  
 of d, 1H); 7.17 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.20 (s, 2H);  
 4.30 (s, 2H); 3.27 (s, 3H); 2.59 (t, 2H);  
 40 1.67 (quint., 2H); 1.35 (sext., 2H); 0.87  
 (t, 3H).

45 c -NMR (200 MHz, CDCl<sub>3</sub>) δ 7.84 (d, 1H); 7.53  
 (t, 1H); 7.40 (t, 1H); 7.29 (m, 3H); 7.04 (d,  
 2H); 5.22 (s, 2H); 4.36 (s, 2H); 3.65 (s,  
 50 3H); 3.61 (sept., 1H); 2.59 (t, 2H); 1.68  
 (quint., 2H); 1.33 (sext., 2H); 1.14 (d, 6H);  
 0.88 (t, 3H).

55

PART B: Preparation of 1-[(3'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole

By the procedure described in Example 85, Part E, 3.35 g of the title compound was prepared from 5.35 g of 1-[(3'-carbomethoxy)biphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  8.33 (s, 1H); 8.11 (d, 1H); 7.80 (d, 1H); 7.55 (t, 1H); 7.34 (A<sub>2</sub>M<sub>2</sub>, 4H); 5.21 (s, 2H); 4.32 (s, 2H); 3.27 (s, 3H); 2.63 (t, 2H); 1.68 (quint., 2H); 1.34 (sext., 2H); 0.86 (t, 3H).

Example 8710 Preparation of 1-[(3'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-acetoxymethylimidazole

A solution of 0.10 g of 1-[(3'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole, 5 mg of N,N-dimethylaminopyridine, 0.10 mL of acetic anhydride, and 0.14 mL of triethylamine in 8 mL of tetrahydrofuran was stirred for 4.5 hours at 25°. The reaction mixture was poured into water, and dilute aqueous sodium hydroxide was added until the pH of the solution remained in the range of pH 8-9. The solution was then acidified to pH 3.5 using 10% aqueous hydrochloric acid and extracted with ethyl acetate. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate and concentrated. Column chromatography on silica gel (elution: 0.5% i-propanol/chloroform) furnished 0.065 g of 1-[(3'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-acetoxymethylimidazole, m.p. 172-173°; NMR (200 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.17 (s, 1H); 7.93 (t, 2H); 7.61 (t, 1H); 7.43 (A<sub>2</sub>M<sub>2</sub>, 4H); 5.32 (s, 2H); 4.99 (s, 2H); 2.60 (t, 2H); 1.76 (s, 3H); 1.53 (quint., 2H); 1.28 (sext., 2H); 0.82 (t, 3H).

Example 8825 Preparation of 1-[(3'-Trimethylacetoxymethoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole

To a solution of 1.25 g of 1-[(3'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole in 10 mL of dimethylformamide at 25° was added 0.17 g of sodium methoxide followed after 5 minutes by 0.45 g of chloromethyl trimethylacetate. The mixture was stirred at 25° for 4 days. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate. This solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography on silica gel afforded 1.38 g of the product as a glassy solid. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.87 (d, 1H); 7.54 (t, 1H); 7.43 (t, 1H); 7.29 (d, 1H); 7.11 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.72 (s, 2H); 5.24 (s, 2H); 4.51 (s, 2H); 2.61 (t, 2H); 2.06 (br s, 1H); 1.68 (quint., 2H); 1.36 (sext., 2H); 1.17 (s, 9H); 0.88 (t, 3H).

Example 89PART A: Preparation of 4'-methylbiphenyl-2-carboxylic acid

Methyl 4'-methylbiphenyl-2-carboxylate (10.0 g, 44.2 mmol, 1 eq), 0.5 N KOH in methanol (265.5 mL, 133 mmol, 3 eq), and water (50 mL) were mixed and refluxed under N<sub>2</sub>. After 5 hours, the solvent was removed *in vacuo* and water (200 mL) and ethyl acetate (200 mL) added. The aqueous layer was acidified with concentrated hydrochloric acid to a pH of 3 and the layers were separated. The aqueous phase was extracted with ethyl acetate (2 x 200 mL), the organic layers collected, dried (MgSO<sub>4</sub>) and the solvent removed *in vacuo* to yield 8.71 g of a white solid; m.p. 140.0-145.0. NMR (200 MHz, DMSO-d<sub>6</sub>)  $\delta$  7.72 (d, 1H, J = 7Hz); 7.56 (t, 1H, J = 7Hz); 7.45 (d, 1H, J = 7Hz); 7.40 (t, 1H, J = 7Hz); 7.25 (s, 4H); 2.36 (s, 3H). Anal. Calcd. for C<sub>14</sub>H<sub>12</sub>O<sub>2</sub>; C, 79.23; H, 5.70. Found: C, 79.22; H, 5.47.

50 PART B: Preparation of 4'-Methyl-2-cyanobiphenyl

4'-Methylbiphenyl-2-carboxylic acid (8.71 g, 41 mmol, 1 eq) and thionyl chloride (30.0 mL, 411 mmol, 10 eq) were mixed and refluxed for 2 hours. The excess thionyl chloride was removed *in vacuo* and the residue was taken up in toluene. The toluene was removed by rotary evaporation and this toluene evaporation procedure was repeated to ensure that all of the thionyl chloride was removed. The crude acid chloride was then added slowly to cold (0° C) concentrated NH<sub>4</sub>OH (50 mL) so that the temperature was kept below 15°. After 15 minutes of stirring, water (100 mL) was added and solids precipitated. These were collected, washed well with water and dried under high vacuum over P<sub>2</sub>O<sub>5</sub> in a dessicator overnight to yield



7.45 g of a white solid; m.p. 126.0-128.5°. NMR (200 MHz, DMSO- $d_6$ )  $\delta$  7.65-7.14 (m, 10H); 2.32 (s, 3H). Anal. Calcd. for  $C_{14}H_{13}NO$ : C, 79.59; H, 6.20; N, 6.63. Found C, 79.29; H, 6.09; N, 6.52.

The above product amide (7.45 g, 35 mmol, 1 eq) and thionyl chloride (25.7 mL, 353 mmol, 10 eq) were mixed and refluxed for 3 hours. The thionyl chloride was removed using the same procedure as described above. The residue was washed with a little hexane which partly solubilized the product, but removed the impurity as well to yield 6.64 g of white solid; m.p. 44.0-47.0°. NMR (200 MHz, DMSO- $d_6$ )  $\delta$  7.95 (d, 1H,  $J$  = 8Hz); 7.78 (t, 1H,  $J$  = 7Hz); 7.69-7.32 (m, 6H); 2.39 (s, 3H). Anal. Calcd. for  $C_{14}H_{11}N$ : C, 87.01; H, 5.74. Found: C, 86.44; H, 5.88.

#### 10 PART C: Preparation of 4'-bromomethyl-2-cyanobiphenyl

4'-methyl-2-cyanobiphenyl (5.59 g) was brominated in the benzylic position by the procedure in Example 85, Part B using benzoyl peroxide as an initiator. The product was recrystallized from ether to yield 4.7 g of product; m.p. 114.5-120.0°. NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.82-7.37 (m, 8H); 4.50 (s, 2H). Anal. Calcd. for  $C_{14}H_{10}BrN$ : C, 61.79; H, 3.70; N, 5.15. Found: C, 62.15; H, 3.45; N, 4.98.

#### PART D: Preparation of 2-n-butyl-4-chloro-1-[(2'-cyanobiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole

4'-Bromomethyl-2-cyanobiphenyl (4.6 g) was alkylated onto 2-n-butyl-4-chloro-5-(hydroxymethyl)-imidazole by the procedure described in Example 1, Part A. Work-up and flash chromatography in 1:1 hexane/ethyl acetate over silica gel to separate the regioisomeric products yielded 2.53 g of the faster eluting isomer. Recrystallization from acetonitrile yielded 1.57 g of analytically pure product; m.p. 153.5-155.5°. NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.82-7.43 (m, 6); 7.12 (d, 2,  $J$  = 8Hz); 5.32 (s, 2); 4.52 (s, 2); 2.62 (t, 2,  $J$  = 7Hz); 1.70 (t of t, 2,  $J$  = 7,7Hz); 1.39 (t of q, 2,  $J$  = 7,7Hz); 0.90 (t, 3,  $J$  = 7Hz). Anal. Calcd. for  $C_{22}H_{22}ClN_3O$ : C, 69.56; H, 5.84; N, 11.06. Found: C, 69.45; H, 5.89; N, 10.79.

#### PART E: Preparation of 2-n-butyl-4-chloro-5-hydroxymethyl-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-imidazole

2-n-Butyl-4-chloro-1-[(2'-cyanobiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole (11.93 g) was converted to the above product by the procedure described in Example 90, Part C. The product was purified by flash chromatography in 100% ethyl acetate to 100% ethanol over silica gel to yield 5.60 g of a light yellow solid. Recrystallization from acetonitrile yielded 4.36 g of light yellow crystals which still melted broadly. The crystals were taken up in 100 mL of hot acetonitrile. The solid that did not dissolve was filtered off to yield 1.04 g of product as a light yellow solid; m.p. 183.5-184.5°. Upon cooling, the mother liquor yielded an additional 1.03 g of product as a light yellow solid; m.p. 179.0-180.0°. NMR (200 MHz, DMSO- $d_6$ )  $\delta$  7.75-7.48 (m, 4H); 7.07 (d, 2H,  $J$  = 9Hz); 7.04 (d, 2H,  $J$  = 9Hz); 5.24 (s, 2H); 5.24 (bs, 1H); 4.34 (s, 2H); 2.48 (t, 2H,  $J$  = 7Hz); 1.48 (t of t, 2H,  $J$  = 7,7Hz); 1.27 (t of q, 2H,  $J$  = 7,7Hz); 0.81 (t, 3H,  $J$  = 7Hz). Anal. Calcd. for  $C_{22}H_{23}ClN_6O$ : C, 62.48; H, 5.48; Cl, 8.38. Found for the solids which did not dissolve in 100 mL of acetonitrile: C, 62.73; H, 5.50; Cl, 8.26. Found for the solids obtained from the mother liquor: C, 62.40; H, 5.23; Cl, 8.35.

#### Example 90

#### 45 PART A: Preparation of 2-n-Butyl-4-chloro-5-chloromethyl-1-[(2'-cyanobiphenyl-4-yl)methyl]imidazole•HCl salt

2-n-Butyl-4-chloro-5-hydroxymethyl-1-[(2'-cyanobiphenyl-4-yl)methyl]imidazole (15.00 g, 39.3 mmol, 1 eq) was converted to the chloride by the procedure in Example 1, Part B. The reaction time was 5 hours. The crude solid product was washed with ether to remove the yellow color. The solid white powdery product was then dried under high vacuum, yield 10.02 g; m.p. 152.0-154.0°. NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.85-7.46 (m, 6H); 7.20 (d, 2H,  $J$  = 10Hz); 5.47 (s, 2H); 4.50 (s, 2H); 3.06 (t, 2H,  $J$  = 7Hz); 1.82 (t of t, 2H,  $J$  = 7,7Hz); 1.45 (t of q, 2H,  $J$  = 7,7Hz); 0.94 (t, 3H,  $J$  = 7Hz). Mass Calcd. for  $C_{22}H_{21}Cl_2N_3$ : 397.1113. Found: 397.1105.

55

PART B: Preparation of 2-n-Butyl-4-chloro-1-[(2'-cyanobiphenyl-4-yl)methyl]-5-(methoxymethyl)imidazole

2-n-Butyl-4-chloro-5-chloromethyl-1-[(2'-cyanobiphenyl-4-yl)methyl]imidazole•HCl salt (5.00 g, 11.5 mmol, 1 eq), sodium methoxide (1.37 g, 25.3 mmol, 2.2 eq) and methanol (100 mL) were mixed and stirred  
 5 for 3 days. The solvent was removed in vacuo and ethyl acetate (200 mL) and water (200 mL) added. The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 200 mL). The organic layers were dried (MgSO<sub>4</sub>), the solvent removed in vacuo and the residue flash chromatographed over silica gel in 1:1 hexane/ethyl acetate to yield 4.06 g of a clear light yellow oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.82-7.43 (m, 6); 7.10 (d, 2H, J = 7Hz); 5.23 (s, 2H); 4.32 (s, 2H); 3.30 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.70 (t of t, 2H,  
 10 J = 7,7Hz); 1.38 (t of q, 2H, J = 7,7Hz); 0.89 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>23</sub>H<sub>24</sub>ClN<sub>3</sub>O: C, 68.11; H, 6.54; Cl, 9.58. Found: C, 68.70; H, 6.11; Cl, 9.51. Mass Calcd. for C<sub>23</sub>H<sub>24</sub>ClN<sub>3</sub>O: 393.1607. Found: 393.1616.

15 PART C: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-methyl)-imidazole

2-n-Butyl-4-chloro-1-[2'-cyanobiphenyl-4-yl)methyl]-5-methoxymethyl)imidazole (3.94 g, 10 mmol, 1 eq), sodium azide (1.95 g, 30 mmol, 3 eq), and ammonium chloride (1.60 g, 30 mmol, 3 eq) were mixed and stirred in DMF (150 mL) in a round bottom flask connected to a reflux condenser under N<sub>2</sub>. An oil bath with  
 20 a temperature controller was then used to heat the reaction at 100 °C for 2 days, after which the temperature was raised to 120 °C for 6 days. The reaction was cooled and 3 more equivalents each of ammonium chloride and sodium azide were added. The reaction was again heated for 5 more days at 120 °C. The reaction was cooled, the inorganic salts filtered, and the filtrate solvent removed in vacuo. Water (200 mL) and ethyl acetate (200 mL) were added to the residue and the layers were separated. The  
 25 aqueous layer was extracted with ethyl acetate (2 x 200 mL), the organic layers were collected, dried (MgSO<sub>4</sub>) and the solvent removed in vacuo, to yield a dark yellow oil. Flash chromatography in 100% ethyl acetate yielded 3.54 g of a white glass. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.83 (d, 1H, J = 7Hz); 7.59 (t, 1H, J = 7Hz); 7.50 (t, 1H, J = 7Hz); 7.39 (d, 1H, J = 7Hz); 7.03 (d, 2H, J = 8Hz); 6.73 (d, 2H, J = 8Hz); 5.08 (s, 2H); 4.12 (s, 2H); 3.18 (s, 3H); 2.32 (t, 2H, J = 7Hz); 1.52 (t of t, 2H, J = 7,7Hz); 1.28 (t of q, 2H, J = 7,7Hz); 0.83  
 30 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>23</sub>H<sub>25</sub>ClN<sub>6</sub>O: 436.1178. Found: 436.1750.

CAUTION! The above reaction although uneventful in our hands can be potentially explosive! Crystals that sublimed and collected in the reflux condenser during the reaction were not analyzed, but potentially could be ammonium azide. Hydrazoic acid, which is shock sensitive, could also be potentially produced during the reaction and work-up. Extreme care should be taken!

35 Example 91

PART A: Preparation of 2-butyl-4(5)-hydroxymethyl-5(4)-nitroimidazole

40 To a solution of 5.75 g of 2-butyl-4(5)-hydroxymethylimidazole (prepared as described in U.S. 4,355,040) in 200 mL of aqueous methanol at 25 °C was added concentrated hydrochloric acid until the pH of the solution reached pH 3. The solvent was then removed in vacuo, and the residue was dissolved in 100 mL of chloroform. To this solution at 25 °C was added dropwise 15.0 mL of thionyl chloride, and the mixture was refluxed for 1 hour. After cooling, the solvent and excess thionyl chloride were removed in vacuo to  
 45 provide a viscous yellow oil.

To a solution of 20 mL of concentrated sulfuric acid and 10 mL of concentrated nitric acid at -10 °C was added a solution of the yellow oil, prepared above, in 10 mL of concentrated sulfuric acid. The resulting mixture was heated on a steam bath for 2 hours. After cooling, the reaction mixture was poured onto water-ice, and the resulting emulsion was extracted with chloroform. The combined organic phases were washed  
 50 with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. The residue was then dissolved in 100 mL of 1:12-propanol/water. The solution was then refluxed for 16 hours. Finally, after cooling, the solution was concentrated in vacuo. Column chromatography (elution: methanol/chloroform) afforded 2.64 g of 2-butyl-4(5)-hydroxymethyl-5(4)-nitroimidazole. NMR (200 MHz, DMSO-d<sub>6</sub>): δ 12.92 (br s, 1H); 5.80 (br t, 1H); 4.82 (d, 2H); 2.60 (t, 2H); 1.61 (quint., 2H); 1.25 (sext., 2H); 0.84 (t,  
 55 3H).



PART B: Preparation of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-nitroimidazole

This compound was prepared according to the procedure described in Example 85, Part C. From 2.64 g of 2-butyl-4(5)-hydroxymethyl-5(4)-nitroimidazole and 5.55 g of tert-butyl 4'-bromomethylbiphenyl-2-carboxylate there was obtained 2.05 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-nitroimidazole. NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.79 (d, 1H); 7.45 (m, 2H); 7.33 (d, 1H); 7.28 (d, 1H); 7.03 (d, 2H); 5.34 (s, 2H); 4.87 (s, 2H); 2.81 (br s, 1H); 2.67 (t, 2H); 1.73 (quint., 2H); 1.37 (sext. 2H); 1.27 (s, 9H); 0.90 (t, 3H).

PART C: Preparation of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-nitroimidazole

A solution of 1.98 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-nitroimidazole, 20 mL of trifluoroacetic acid, and 20 mL of methylene chloride was stirred at 25° for 1 hour. At this point, the solution was poured into water. The resulting mixture was adjusted to pH 3 using 10% sodium hydroxide solution and then extracted with chloroform. The combined organic phases were washed with brine, dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo*. Column chromatography (elution: methanol/chloroform) provided 1.49 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-nitroimidazole; m.p. 204-205.5°. NMR (200 MHz, DMSO-d<sub>6</sub>):  $\delta$  7.71 (d, 1H); 7.56 (t, 1H); 7.43 (t, 1H); 7.32 (m, 3H); 7.15 (d, 2H); 5.63 (br s, 1H); 5.42 (s, 2H); 4.83 (s, 2H); 2.54 (t, 2H); 1.50 (quint., 2H); 1.24 (sext., 2H); 0.76 (t, 3H).

Example 92

PART A: Preparation of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-4-iodo-5-(2-methoxyethoxymethoxymethyl)imidazole

To a solution of 5.56 mL of 1.6 M n-butyllithium/hexane in 80 mL of tetrahydrofuran at 0° was added dropwise 1.15 mL of t-butanol. To the solution was added 3.28 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-iodoimidazole followed by 1.15 mL of 2-methoxyethoxymethyl chloride. The resulting solution was stirred at 25° for 16 hours. The mixture was diluted with diethyl ether, washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography afforded 2.61 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-4-iodo-5-(2-methoxyethoxymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.78 (d, 1H); 7.43 (m, 2H); 7.28 (m, 3H); 6.98 (d, 2H); 5.26 (s, 2H); 4.69 (s, 2H); 4.45 (s, 2H); 3.68 (m, 2H); 3.57 (m, 2H); 3.37 (s, 3H); 2.58 (t, 2H); 1.67 (quint., 2H); 1.34 (sext., 2H); 1.26 (s, 9H); 0.87 (t, 3H).

PART B: Preparation of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-(2-methoxyethoxymethoxymethyl)-4-trifluoromethylimidazole

To a suspension of 22.4 g of cadmium powder powder in 50 mL of dimethylformamide at 25° was added dropwise 8.60 mL of bromochlorodifluoromethane. The resulting mixture was stirred at 25° for 2 hours and then was filtered through a medium-fritted Schlenk funnel under nitrogen pressure to provide a dark brown solution of the trifluoromethyl cadmium reagent.

To a mixture of 15 mL of the above solution and 20 mL of hexamethylphosphoric triamide at 0° was added 2.10 g of copper(I)bromide followed by 2.61 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-4-iodo-5-(2-methoxyethoxymethoxymethyl)imidazole in 5 mL of dimethylformamide. The reaction mixture was stirred at 70-75° for 6 hours. After cooling, the mixture was diluted with water and then extracted with methylene chloride. The combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography (elution: ethyl acetate/hexane) afforded 2.30 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-(2-methoxyethoxymethoxymethyl)-4-trifluoromethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.79 (d, 1H); 7.46 (m, 2H); 7.28 (m, 3H); 7.00 (d, 2H); 5.28 (s, 2H); 4.71 (s, 2H); 4.58 (s, 2H); 3.66 (m, 2H); 3.54 (m, 2H); 3.38 (s, 3H); 2.62 (t, 2H); 1.70 (quint., 2H); 1.36 (sext., 2H); 1.27 (s, 9H); 0.88 (t, 3H).

PART C: Preparation of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-trifluoromethylimidazole

A solution of 2.30 g of 1-[(2'-tert-butoxycarbonylbiphenyl-4-yl)methyl]-2-butyl-5-(2-methoxy ethox-  
 ymethoxymethyl)-5-trifluoromethylimidazole in 200 mL of 1.5 M aqueous tetrafluoroboric acid/acetonitrile  
 was stirred at 25 ° for 18 hours, and then the mixture was poured into water. The resulting aqueous solution  
 was adjusted to pH 3 employing saturated sodium bicarbonate solution and then was extracted with  
 chloroform. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate,  
 filtered, and concentrated. Column chromatography (elution: methanol/chloroform) provided 1.38 g of 1-[(2'-  
 carboxybiphenyl-4-yl)methyl]-2-butyl-5-hydroxymethyl-4-trifluoromethylimidazole (m.p. 198-199.5 °). NMR  
 (200 MHz, DMSO-d<sub>6</sub>): δ 7.75 (d, 1H); 7.54 (t, 1H); 7.43 (t, 1H); 7.32 (m, 3H); 7.10 (d, 2H); 5.36 (s, 2H); 4.51  
 (s, 2H); 2.56 (t, 2H); 1.56 (quint., 2H); 1.30 (sext., 2H); 0.83 (t, 3H).

Example 93

15

PART A: Preparation of 4-azidomethyl-2'-methoxycarbonylbiphenyl

To a stirred solution of 4-bromomethyl-2'-methoxycarbonylbiphenyl (150 g, 0.49 mol) in dry DMF (500  
 ml) was added NaN<sub>3</sub> (80 g, 1.23 mol, 2.5 eq). The mixture was stirred at room temperature overnight (ca.  
 18 hours), filtered, and the filtrate was partitioned between ethyl acetate and H<sub>2</sub>O (500 ml each). The  
 organic phase was washed twice more with H<sub>2</sub>O, once with saturated aqueous NaCl solution and dried over  
 anhydrous magnesium sulfate before being filtered and concentrated to leave 111.3 g (85%) of a yellow oil,  
 used in the following step without further purification. NMR (CDCl<sub>3</sub>, TMS, δ) 7.9-7.1 (m, 8H); 4.35 (s, 2H);  
 3.55 (s, 3H) IR V<sub>max</sub> 2487 cm<sup>-1</sup>.

25

PART B: Preparation of 4-aminomethyl-2'-methoxycarbonylbiphenyl hydrochloride

The azido compound prepared above was dissolved in liter of methanol. The solution was divided into  
 three equal volumes and placed in 500 ml Parr bottles. To each flask was added 6.7 g of 5% Pd on carbon  
 (Caution: Pyrophoric! add under a N<sub>2</sub> atmosphere). The flasks were shaken on a Parr hydrogenator under  
 40-50 psi H<sub>2</sub> for 4-5 hours (overnight is also acceptable). The mixture was suction filtered through a bed of  
 Celite® and the filtrate was concentrated to leave a viscous yellow residue (88 g). This was dissolved in  
 EtOAc (500 ml) to which was added with stirring a solution of EtOAc saturated with anhydrous HCl (100-150  
 ml) until precipitation was complete. The amine hydrochloride as produced was suction filtered, washed  
 with EtOAc and hexanes and dried under vacuum to afford 48.5 g (40% overall from the bromide) white  
 solid; m.p. 204-208 °. NMR (CDCl<sub>3</sub>, CD<sub>3</sub>OD; TMS) δ 7.9-7.25 (m, 8H); 4.2 (s, 2H); 4.1-3.8 (br, 3H; shifts in  
 D<sub>2</sub>O); 3.6 (s, 3H). HRMS calcd. for C<sub>15</sub>H<sub>15</sub>NO<sub>2</sub> (free base); M/Z 241.1103; Found: M/Z: 241.1045.

40

PART C: Preparation of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-propylthio-5-hydroxymethylimidazole

The title compound was prepared from methyl 4'-aminomethylbiphenyl-2-carboxylate by the procedures  
 described in Examples 72, A and B, and 85E; m.p. 194-195 °.


The 4-biphenylmethyl compounds in Table 6 were prepared or could be prepared by the procedures  
 illustrated in Examples 85-92 or by procedures previously described.

45

50

55

Table 6










Ex. No.	$\underline{R^6}$	$\underline{R^7}$	$\underline{R^8}$		MP(°C)
94	n-butyl Cl		CH <sub>2</sub> OH		168-169.5
95	n-butyl CH <sub>2</sub> OH		Cl		197-198
96	n-butyl H		CH <sub>2</sub> OH		154-155
97	n-butyl H		CH <sub>2</sub> OH		(amorphous solid) <sup>a</sup>
98	n-butyl Cl		CH <sub>2</sub> OCH <sub>3</sub>		166.5-169.0
99	n-butyl Cl		CH <sub>2</sub> OCH(CH <sub>3</sub> ) <sub>2</sub>		156-158

Table 6 (continued)












Ex. No.	$R^6$	$R^7$	$R^8$		MP(°C)
100	n-butyl	Br	CH <sub>2</sub> OH		175-178
101	n-butyl	F	CH <sub>2</sub> OH		
102	n-butyl	I	CH <sub>2</sub> OH		165 (dec)
103		Cl	CH <sub>2</sub> OH		
104		Cl	CH <sub>2</sub> OH		
105	n-butyl	CH <sub>2</sub> OH	I		205 (dec)
106	n-butyl	Cl	CH <sub>2</sub> OH		185-186
107	ethyl	Cl	CH <sub>2</sub> OH		153-156

Table 6 (continued)








5						
	<u>Ex. No.</u>	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>		<u>MP (°C)</u>
10	108	n-propyl	Cl	CH <sub>2</sub> OH		198-200
15	109	n-pentyl	Cl	CH <sub>2</sub> OH		(amorphous <sup>b</sup> solid)
20	110	n-hexyl	Cl	CH <sub>2</sub> OH		84-88
25	111	n-butyl	Cl	CH <sub>2</sub> SH		
30	112	n-butyl	Cl	CH <sub>2</sub> O- 		
35						
40						
45						
50						
55						

Table 6 (continued)


















Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>		MP(°C)
113	n-propyl	Cl	CH <sub>2</sub> OH		(amorphous <sup>c</sup> solid)
114	n-propyl	Cl	CHO		(amorphous <sup>d</sup> solid)
115	n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> H		221-222
116	n-butyl	Cl	CH(CH <sub>3</sub> )CO <sub>2</sub> H		118-120
117	n-butyl	CH <sub>2</sub> OH	NO <sub>2</sub>		154-157
118	n-butyl	CH <sub>2</sub> OH	Cl		(white <sup>e</sup> powder)
119	n-butyl	NO <sub>2</sub>	CH <sub>2</sub> OH		

Table 6 (continued)

Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>		MP (°C)
120	n-butyl	Cl			
121	n-butyl	Cl	CH <sub>2</sub> OCOCH <sub>3</sub>		157-159
122	n-butyl	Cl	CH <sub>2</sub> OCOCH <sub>2</sub> CH <sub>2</sub> - 		
123	n-C <sub>4</sub> H <sub>9</sub> S	H	CH <sub>2</sub> OH		190-191
124	 CH <sub>2</sub> S	H	CH <sub>2</sub> OH		194.5-195.5

a -NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.69 (dd, 1H);  
 7.54 (d of t, 1H); 7.43 (d of t, 1H); 7.33  
 (d, 1H); 7.16 (A<sub>2</sub>B<sub>2</sub>, 4H); 6.76 (s, 1H);  
 5.24 (s, 2H); 4.34 (s, 2H); 2.50 (t, 2H);  
 1.49 (quint, 2H); 1.25 (sext, 2H); 0.80 (t,  
 3H).

b -NMR (200 MHz, DMSO- $d_6$ )  $\delta$  7.70 (d, 1H).

7.55 (t, 1H), 7.42 (t, 1H), 7.28 (m, 3H),

7.10 (d, 2H), 5.28 (s, 2H), 4.34 (s, 2H),

2.49 (t, 2H), 1.49 (m, 2H), 1.18 (m, 4H),

0.79 (t, 3H).

c -NMR (200 MHz,  $CDCl_3/CD_3OD$ ):  $\delta$

7.82-6.93 (m, 8H); 5.21 (s, 2H); 4.47 (s,

2H); 2.55 (t,  $J = 7.5$  Hz, 2H); 1.70-1.59 (m,

2H); 0.92 (t,  $J = 7.5$  Hz, 3H).

d -NMR (200 MHz,  $CDCl_3$ ) 9.65 (s, 1H);

7.95-6.96 (m, 8H); 5.51 (s, 2H); 2.59 (t,  $J =$

7.5 Hz, 2H); 1.70-1.63 (m, 2H); 0.92 (t,  $J =$

7.5 Hz, 3H).

e -NMR (200 MHz,  $CDCl_3$ )  $\delta$  7.76 (d, 1H,  $J =$

7 Hz); 7.57 (t, 1H,  $J = 7$  Hz); 7.49 (t, 1H,  $J =$

7 Hz); 7.40 (d, 1H,  $J = 7$  Hz); 7.02 (d, 2H,  $J =$

8 Hz); 6.81 (d, 2H,  $J = 8$  Hz); 5.03 (s, 2H);

4.28 (s, 2H); 2.46 (t, 2H,  $J = 7$  Hz); 1.47 (t

of t, 2H,  $J = 7.7$  Hz); 1.17 (t of q, 2H,  $J =$

7.7 Hz); 0.73 (t, 3H,  $J = 7$  Hz).

#### Example 125

##### Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde

A mixture of 1.46 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole and 7.30 g of activated manganese dioxide in 40 ml of tetrahydrofuran was stirred at 25°C for 5 days. The mixture was filtered through Celite®, and the filtrate was concentrated in vacuo. Column chromatography on silica gel (elution: 2-10% methanol/chloroform) followed by recrystallization from ethyl acetate afforded 0.71 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde (m.p. 154-158°C (dec.)). NMR (200 MHz, DMSO- $d_6$ )  $\delta$  12.85 (br s, 1H), 9.77 (s, 1H), 7.77 (d, 1H), 7.62 (t, 1H), 7.50 (t, 1H), 7.40 (d, 1H), 7.26 ( $A_2B_2$ , 4H), 5.67 (s, 2H), 2.70 (t, 2H), 1.56 (quint., 2H), 1.28 (sext., 2H), 0.83 (t, 3H).

#### Example 126

##### Preparation of Methyl 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxylate

To a mixture of 1.45 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde and 0.91 g of sodium cyanide in 20 mL of methanol at 25°C was added 0.32 mL of acetic acid followed by 7.25 g of manganese dioxide. The resulting mixture was stirred at 25°C for 40 hours. The reaction mixture was filtered through Celite®, and the filtrate diluted with water. The aqueous solution was adjusted to pH 3 using hydrochloric acid and extracted with methylene chloride. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product was recrystallized from diethyl ether to afford 0.90 g of methyl 1-[(2'-carboxybiphenyl-4-yl)methyl]-



2-butyl-4-chloroimidazole-5-carboxylate (m.p. 154-155 °C). NMR (200 MHz, DMSO- $d_6$ ):  $\delta$  12.75 (br s, 1H), 7.73 (d, 1H), 7.58 (t, 1H), 7.46 (t, 1H), 7.34 (m, 3H), 7.07 (d, 2H), 5.63 (s, 2H), 3.78 (s, 3H), 2.67 (t, 2H), 1.56 (quint., 2H), 1.29 (sext., 2H), 0.83 (t, 3H).

#### 5 Example 127

##### Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxamide

Anhydrous ammonia was bubbled into 40 mL of i-propanol until the solvent was saturated. To this  
 10 solution at 25 °C was added 0.49 g of powdered sodium cyanide, then 0.80 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde, and finally 3.48 g of manganese dioxide. This mixture was stirred at 25 °C for 65 hours. The reaction mixture was filtered through Celite®, and the filtrate concentrated in vacuo. The residue was dissolved in water, and the aqueous solution was adjusted to pH 3 using hydrochloric acid and then extracted with methylene chloride. The combined organic phases were  
 15 washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 0-10% i-propanol (chloroform) provided 0.22 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxamide as a white solid (m.p. 200-202 °C). NMR (200 MHz, DMSO- $d_6$ ):  $\delta$  12.74 (br s, 1H); 7.71 (d, 2H); 7.56 (t, 1H), 7.48-7.30 (m, 6H); 7.09 (s, 2H); 5.57 (s, 2H); 2.59 (t, 2H); 1.51 (quint., 2H); 1.26 (sext., 2H); 0.80 (s, 3H).

20

#### Example 128

##### PART A: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde

25

A mixture of 2.06 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole and 3.08 g of activated manganese dioxide in 20 mL of methylene chloride at 25 °C was stirred for 40 hours. The reaction mixture was filtered through Celite®, and the filtrate concentrated in vacuo. Column chromatography (elution: ethyl acetate/benzene) provided 1.15 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde. NMR (200 MHz,  $CDCl_3$ )  $\delta$  9.76 (s, 1H);  
 30 7.83 (d of d, 1H); 7.52 (t of d, 1H); 7.40 (t of d, 1H); 7.31 (d of d, 1H); 7.17 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.58 (s, 2H); 3.63 (s, 3H); 2.67 (t, 2H); 1.70 (quint., 2H); 1.38 (sext., 2H); 0.90 (t, 3H).

##### PART B: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-2-(1-bromobutyl)-4-chloroimidazole-5-carboxaldehyde

35

A mixture of 1.12 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxaldehyde and 0.49 g of N-bromosuccinimide in 40 mL of  $CCl_4$  was irradiated (UV-lamp, pyrex filter) for 0.5 hours. The reaction mixture was filtered, and the filtrate was concentrated in vacuo. Column chromatography (elution: ethyl acetate/benzene) afforded 0.54 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-bromobutyl)-4-chloroimidazole-5-carboxaldehyde. NMR (200 MHz,  $CDCl_3$ )  $\delta$  9.87 (s, 1H); 7.86 (d, 1H); 7.54 (t, 1H); 7.46 (t, 1H); 7.30 (m, 3H); 7.11 (d, 2H); 6.16 (d, 1H); 5.32 (d, 1H); 4.79 (t, 1H); 3.65 (s, 3H); 2.32 (m, 2H); 1.34 (sext., 2H); 0.83 (t, 3H).

##### PART C: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloroimidazole-5-carboxaldehyde

45

A solution of 0.54 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-bromobutyl)-4-chloroimidazole-5-carboxaldehyde and 0.33 mL of 1,8-diazabicyclo[4.5.0]undec-7-ene in 10 mL of tetrahydrofuran was stirred  
 50 at 25 °C for 18 hours. the reaction mixture was diluted with diethyl ether, washed with dilute hydrochloric acid, water, and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography (elution: ethyl acetate/benzene) furnished 0.26 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloroimidazole-5-carboxaldehyde. NMR (200 MHz,  $CDCl_3$ )  $\delta$  9.75 (s, 1H); 7.82 (d, 1H); 7.51 (t, 1H); 7.40 (t, 1H); 7.33-7.07 (m, 6H); 6.27 (d, 1H); 5.62 (s, 2H); 3.62 (s, 3H); 2.30 (quint., 2H);  
 55 1.09 (t, 3H).

PART D: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole

To a solution of 0.26 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloroimidazole-5-carboxaldehyde in 10 mL of methanol at 0 °C was added 0.24 g of sodium borohydride portionwise over 0.5 hours. The mixture was stirred for an additional 0.5 hours at 0 °C and then poured into a solution at 10% sodium hydroxide in water. The resulting mixture was extracted with ethyl acetate, and the combined organic phases were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography (elution: ethyl acetate/benzene) provided 0.23 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.84 (d, 1H); 7.53 (t, 1H); 7.40 (t, 1H); 7.29 (m, 3H); 7.08 (d, 2H); 6.86 (d of t, 1H); 6.17 (d, 1H); 5.30 (s, 2H); 4.54 (br s, 2H); 3.63 (s, 3H); 2.23 (quint., 2H); 1.04 (t, 3H).

PART E: Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole

This compound was prepared according to the procedure described in Example 85, Part E. From 0.23 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole there was obtained 0.16 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole (m.p. 198.5-199.5 °C). NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.71 (d, 1H); 7.56 (t, 1H); 7.44 (t, 1H); 7.32 (m, 3H); 7.11 (d, 2H); 6.62 (d of t, 1H); 6.39 (d, 1H); 5.38 (s, 2H); 5.33 (br s, 1H); 4.35 (br s, 2H); 2.18 (quint., 2H); 0.99 (t, 3H).

Example 129

Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloroimidazole-5-carboxaldehyde

This compound was prepared according to the procedure of Example 125. From 0.50 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloro-5-hydroxymethylimidazole and 2.50 g of manganese dioxide was obtained 0.24 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-(1-trans-butenyl)-4-chloroimidazole-5-carboxaldehyde (m.p. 164-166 °C). NMR (200 MHz, DMSO-d<sub>6</sub>) δ 12.79 (br s, 1H); 9.70 (s, 1H); 7.72 (d, 1H); 7.57 (t, 1H); 7.46 (t, 1H); 7.33 (m, 3H); 7.15 (d, 2H); 7.01 (d of t, 1H); 6.65 (d, 1H); 5.71 (s, 2H); 2.28 (quint., 2H); 1.04 (t, 3H).

The compounds in Table 7 were prepared or could be prepared employing the procedures described in Examples 125-129 or by procedures described previously.

Table 7















5					
10					
15	Ex. No.	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	 <u>MP (°C)</u>
20	130	n-butyl	H	CHO	 (amorphous <sup>a</sup> solid)
25	131	n-butyl	CF <sub>3</sub>	CHO	 132-134
30	132	n-butyl	Cl	CHO	 82.7
40	133	n-butyl	CF <sub>3</sub>	CHO	
45					
50					
55					

Table 7 (continued)

5	Ex. No. $R^6$	$R^7$	$R^8$		MP (°C)
10	134 n-butyl	Cl	CONHCH <sub>3</sub>		(solid) <sup>b</sup>
15	135 n-butyl	Cl	CON(CH <sub>3</sub> ) <sub>2</sub>		(solid) <sup>c</sup>
20	136 CH <sub>3</sub> CH=CH-	Cl	CH <sub>2</sub> OH		
25	137 CH <sub>3</sub> CH <sub>2</sub> CH=CH-	CF <sub>3</sub>	CH <sub>2</sub> OH		
35	138 CH <sub>3</sub> CH <sub>2</sub> CH=CH-	Cl	CHO		
40	139 CH <sub>3</sub> CH <sub>2</sub> CH=CH-	Cl	CH <sub>2</sub> OH		
45	140 CH <sub>3</sub> CH <sub>2</sub> CH=CH-	Cl	CHO		

55

- a -NMR (200 MHz, DMSO- $d_6$ )  $\delta$  12.76 (br s, 1H); 9.67 (s, 1H); 7.93 (s, 1H); 7.71 (d, 1H); 7.55 (t, 1H); 7.43 (t, 1H); 7.30 (m, 3H); 7.06 (d, 2H); 5.63 (s, 2H); 2.67 (t, 2H); 2.57 (quint., 2H); 2.27 (sext., 2H); 0.81 (t, 3H).
- b -NMR (200 MHz, DMSO- $d_6$ )  $\delta$  12.75 (br s, 1H); 8.10 (br quart., 1H); 7.72 (d, 1H); 7.57 (t, 1H); 7.45 (t, 1H); 7.32 (m, 3H); 7.10 (d, 2H); 5.51 (s, 2H); 2.75 (d, 3H); 2.58 (t, 2H); 1.52 (quint., 2H); 1.27 (sext., 2H); 0.81 (t, 3H).
- c -NMR (200 MHz, DMSO- $d_6$ )  $\delta$  12.77 (br s, 1H); 7.73 (d, 1H); 7.57 (t, 1H); 7.45 (t, 1H); 7.33 (m, 3H); 7.09 (d, 2H); 5.20 (br s, 2H); 2.83 (s, 3H); 2.73 (t, 2H); 2.66 (s, 3H); 1.63 (quint., 2H); 1.36 (sext., 2H); 0.89 (t, 3H).

#### Example 141

##### PART A: Preparation of 1-[2'-Aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole


A solution of 4.40 g of 1-[(2'-nitrobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole, 2.10 g of iron powder, 4.25 mL of glacial acetic acid, and 200 mL of methanol was refluxed for 5 hours. After cooling, the solvent was removed *in vacuo*, and the residue was dissolved in ethyl acetate. The precipitated iron salts were removed by filtration through Celite®, and the resulting solution was washed with water and brine, dried over anhydrous sodium sulfate and concentrated. Column chromatography on silica gel (elution: 10-30% ethyl acetate/benzene) furnished 2.95 g of 1-[2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole; NMR (200 MHz,  $CDCl_3$ ):  $\delta$  7.43 (d, 2H); 7.19-7.04 (m, 4H); 6.80 (m, 2H); 5.19 (s, 2H); 4.33 (s, 2H); 3.70 (br s, 1H); 3.28 (s, 3H); 2.59 (t, 2H); 1.67 (quint., 2H); 1.34 (sext., 2H); 0.87 (t, 3H).

##### PART B: Preparation of 1-[2'-Trifluoromethanesulfonamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole

To a solution of 2.95 g of 1-[(2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole and 1.07 mL of triethylamine in 30 mL of methylene chloride at -78° was added 2.59 mL of trifluoromethanesulfonic anhydride dropwise at such rate that the reaction temperature remains below -50°. Following the addition, the reaction mixture was allowed to warm slowly to 25°. At the point the mixture was poured into dilute aqueous acetic acid. The resulting suspension was stirred vigorously for several minutes and then extracted with methylene chloride. The combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography on silica gel (elution: 20-50% ethyl acetate/benzene) afforded 0.80 g of 1-[(2'-trifluoromethanesulfonamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-methoxymethylimidazole, m.p. 148-150°; NMR (200 MHz,  $CDCl_3$ ):  $\delta$  7.60 (d, 1H); 7.44-7.27 (m, 5H); 7.07 (d, 2H); 5.20 (s, 2H); 4.29 (s, 2H); 3.27 (s, 3H); 2.57 (t, 2H); 1.65 (quint., 2H); 1.35 (sext., 2H); 0.88 (t, 3H).

Examples 142 to 147 can or could be prepared by the procedures described in Example 141 using the appropriate starting material.

Table 8



Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
142	n-butyl	H	CH <sub>2</sub> OCH <sub>3</sub>		
143	n-hexyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		
144	n-butyl	Cl	CH <sub>2</sub> OH		171-172
145	FCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -	Cl	CH <sub>2</sub> OH		
146	HO <sub>2</sub> CCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -	Cl	CH <sub>2</sub> OH		
147	CH <sub>3</sub> O <sub>2</sub> CCH <sub>2</sub> CH <sub>2</sub> -	Cl	CH <sub>2</sub> OH		

## Example 148

PART A: Preparation of 2-Butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(chloromethyl)-imidazole•HCl salt

5 2-Butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(chloromethyl)imidazole•HCl salt was prepared from 2-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(hydroxymethyl)imidazole using the procedure of Example 1, Part B; m.p. 156.0-161.0°. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.90 (d, 1H, 7Hz); 7.56 (t, 1H, J = 7Hz); 7.45 (t, 1H, J = 7Hz); 7.43-7.26 (m, 3H); 7.12 (d, 2H, J = 8Hz); 5.47 (s, 2H); 4.48 (s, 2H); 3.70 (s, 10 3H); 3.14 (t, 2H, J = 7Hz); 1.80 (t of t, 2H, J = 7.7Hz); 1.44 (t of q, 2H, J = 7.7Hz); 0.92 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>23</sub>H<sub>24</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>2</sub>•HCl: C, 59.05; H, 5.39; N, 5.99. Found: C, 58.80; H, 5.48; N, 5.69. Mass Calcd. for C<sub>23</sub>H<sub>24</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>2</sub>: 430.1215. Found 430.1215.

PART B: Preparation of 5-Azidomethyl-2-n-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole

15 2-Butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(chloromethyl)imidazole•HCl salt (3.31 g, 7.67 mmol, 1 eq), sodium azide (1.50 g, 23.0 mmol, 3 eq) and DMSO (100 mL) were mixed and stirred overnight. Water was then added (500 mL) and the aqueous extracted with ethyl acetate (3 x 300 mL). The organic layers were dried (MgSO<sub>4</sub>) and concentrated to yield 3.48 g of product as an oil. NMR (200 MHz, 20 CDCl<sub>3</sub>) δ 7.85 (d, 1H, J = 7Hz); 7.54 (t, 1H, J = 7Hz); 7.40 (t, 1H, J = 7Hz); 7.28 (d, 2H, J = 8Hz); 7.00 (d, 2H, J = 8Hz); 5.20 (s, 2H); 4.23 (s, 2H); 3.67 (s, 3H); 2.63 (t, 2H, J = 7Hz); 1.73 (t of t, 2H, J = 7.7Hz); 1.39 (t of q, 2H, J = 7.7Hz); 0.91 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>23</sub>H<sub>24</sub>ClN<sub>5</sub>O<sub>2</sub>: 438.1697. Found: 438.1669.

PART C: Preparation of 5-Aminomethyl-2-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole

25 5-Azidomethyl-2-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole (3.48 g) was hydrogenated at 1 atm in methanol (100 mL) over 10% palladium/carbon (0.5 g). After 1 hour, the mixture was filtered through Celite® and the solvent removed *in vacuo* to give product (2.80 g) as an oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.84 (d, 1H, J = 7Hz); 7.52 (t, 1H, J = 7Hz); 7.40 (t, 1H, J = 7Hz); 7.30 (d, 1H, J = 7Hz); 7.26 30 (d, 2H, J = 8Hz); 7.02 (d, 2H, J = 8Hz); 5.27 (s, 2H); 3.74 (s, 2H); 3.65 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.67 (t of t, 2H, J = 7.7Hz); 1.36 (t of q, 2H, J = 7.7Hz); 0.86 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>23</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>2</sub>••(DMSO)<sub>0.5</sub>: C, 63.91; H, 6.48; N, 9.32. Found: C, 63.78; H, 6.30; N, 9.14

PART D: Preparation of 5-Aminomethyl-2-butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloroimidazole


35 5-Aminomethyl-2-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole (1.64 g, 3.98 mmol, 1 eq), 0.5N KOH in methanol (11.96 mL, 5.98 mmol, 1.5 eq), water (1.0 mL) and methanol (20 mL) were mixed and refluxed under N<sub>2</sub> overnight. The solution was then brought to neutrality with 1N HCl and the solvents removed *in vacuo*. The residue was taken up in DMF and the salts filtered off. The DMF was then 40 removed *in vacuo* to yield 1.76 g of a glass. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.50 (d, 1H, J = 7Hz); 7.40-7.18 (m, 5H); 6.92 (d, 2H, J = 8Hz); 6.50 (bm, 3H); 5.26 (s, 2H); 3.60 (s, 2H); 2.55 (t, 2H, J = 7Hz); 1.51 (t of t, 2H, J = 7.7Hz); 1.27 (t of q, 2H, J = 7.7Hz); 0.81 (t, 3H, J = 7Hz).

PART E: Preparation of 2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloro-5-(ethoxycarbonylaminomethyl)imidazole

45 2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloro-5-(ethoxycarbonylaminomethyl)imidazole was prepared from 5-aminomethyl-2-n-butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloroimidazole using ethyl chloroformate and the Schotten-Baumann procedure described in Example 209, Part B; m.p. 144.0-147.0°. 50 NMR (200 MHz, DMSO-d<sub>6</sub>) δ 12.74 (s, 1H); 7.73 (d, 1H, J = 7Hz); 7.63-7.27 (m, 5H); 7.03 (d, 2H, J = 10Hz); 5.27 (s, 2H); 4.60 (bd, 2H, J = 7Hz); 3.90 (q, 2H, J = 7Hz); 3.34 (s, 2H); 2.47 (t, 2H, J = 7Hz); 1.48 (t of t, 2H, J = 7.7Hz); 1.24 (t of q, 2H, J = 7.7Hz); 1.06 (t, 3H, J = 7Hz); 0.78 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>25</sub>H<sub>28</sub>ClN<sub>3</sub>O<sub>4</sub>•(H<sub>2</sub>O)<sub>0.33</sub>: C, 63.17; H, 6.06; N, 8.83. Found: C, 63.30; H, 6.35; N, 8.44.

55 Examples 149-159 in Table 9 were prepared or could be prepared using the appropriate chloroformate by the procedure described in Example 148, Parts D and E (the order of which may be interchanged by one skilled in the art) i.e., starting with the amino ester from Part C, reacting it with a chloroformate under Schotten-Baumann type conditions followed by hydrolyzing the ester if necessary.

Table 9




Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R	R <sup>13</sup>	MP (°C)
149	n-butyl	Cl	C <sub>6</sub> H <sub>5</sub>	CO <sub>2</sub> H	198.0-200.0
150	n-butyl	Cl	CH <sub>3</sub>	CO <sub>2</sub> H	151.0-155.0
151	n-butyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CO <sub>2</sub> H	115.5-117.0
152	n-butyl	Cl	CH <sub>2</sub> (CH <sub>3</sub> ) <sub>2</sub>	CO <sub>2</sub> H	135.5-138.0
153	n-butyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CO <sub>2</sub> H	123.0-125.0
154	n-butyl	Cl	1-adamantyl	CO <sub>2</sub> H	170.0-172.0
155	n-propyl	Cl	CH <sub>3</sub>	CO <sub>2</sub> H	
156	n-butyl	Cl	CH <sub>3</sub>		202.0-204.5
157	n-butyl	Cl	(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>		
158	n-propyl	Cl	CH <sub>3</sub>		
159	n-propyl	H	CH <sub>2</sub> CH <sub>3</sub>		

Examples 160-164 in Table 10 were prepared or could be prepared from 2-n-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-5-chloro-4-(hydroxymethyl)imidazole using the procedures in Example 148.



Table 10



Ex. No.	R <sup>6</sup>	R <sup>8</sup>	R	R <sup>13</sup>	MP (°C)
160	n-butyl	Cl	CH <sub>3</sub>	COOH	200-205
161	n-butyl	Cl	CH <sub>2</sub> CH <sub>3</sub>	COOH	
162	n-butyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	COOH	166.5-169.5
163	n-butyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	COOH	
164	n-butyl	Cl	CH(CH <sub>3</sub> ) <sub>2</sub>	COOH	

## EXAMPLE 165

PART A: Preparation of 2-n-Butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(1-naphthylaminocarbonylaminomethyl)imidazole


5-Aminomethyl-2-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole (1.00 g, 2.4 mmol, 1 eq) and 1-naphthyl isocyanate (0.35 mL, 2.4 mmol, 1 eq), were mixed and stirred in chloroform at room temperature for 3 days. The solvent was removed in vacuo and the residue was purified by flash chromatography over silica gel in 1:1 hexane/ethyl acetate to yield 770 mg of a white glass. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.83 (d, 3H, J = 6Hz); 7.67 (d, 1H, J = 6Hz); 7.56-7.18 (m, 9H); 6.97 (d, 2H, J = 7Hz); 6.74 (s, 1H); 5.27 (s, 2H); 4.74 (s, 1H); 4.39 (d, 2H, J = 7Hz); 3.58 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.43-1.21 (m, 4H); 0.85 (t, 3H, J = 7Hz).

PART B: Preparation of 2-n-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloro-5-(1-naphthylaminocarbonylaminomethyl)imidazole

The title compound was prepared from 2-n-butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(1-naphthylaminocarbonylaminomethyl)imidazole by the hydrolysis procedure described in Example 148, Part D. Work-up yielded 380 mg of white crystalline solid; m.p. 169-175°. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.45 (s, 1H); 8.05-7.03 (m, 15H); 6.97 (s, 1H); 5.34 (s, 2H); 4.30 (d, 2H, J = 5Hz); 2.52 (t, 2H, J = 7Hz); 1.48 (t of t, 2H, J = 7,7Hz); 1.21 (t of q, 2H, J = 7,7Hz); 0.85 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>33</sub>H<sub>31</sub>ClN<sub>4</sub>O<sub>3</sub> · (H<sub>2</sub>O)<sub>0.5</sub>: C, 68.77; H, 5.60; N, 9.70. Found: C, 68.88; H, 5.67; N, 9.70.

Examples 166-172 in Table 11 were prepared or could be prepared using the appropriate isocyanate by the procedure described in Example 165.

Table 11



Ex. No.	R <sup>6</sup>	R <sup>8</sup>	R	R <sup>13</sup>	MP (°C)
166	n-Bu	Cl	CH <sub>3</sub>	CO <sub>2</sub> H	187-193
167	n-Bu	Cl	CH <sub>2</sub> CH <sub>3</sub>	CO <sub>2</sub> H	
168	n-Bu	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CO H	
169	n-Bu	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CO <sub>2</sub> H	
170	n-Bu	Cl	CH(CH <sub>3</sub> ) <sub>2</sub>	CO H	
171	n-Bu	Cl		CO <sub>2</sub> H	163-166
172	n-Bu	Cl	1-adamantyl		

## Example 173

## Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[(2'-((tetrazol-5-yl)aminocarbonyl)biphenyl-4-yl)methyl]-imidazole

2-n-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]-4-chloro-5-(methoxymethyl)imidazole (1.0 g) was first converted to the corresponding acid chloride and then coupled to 5-aminotetrazole by the procedure in Example 78, Part C to yield 0.87 g of a yellow glass. Flash chromatography in 100% ethyl acetate over silica gel yielded 77.1 mg of a white solid; m.p. 169-173°. NMR (200 MHz, CDCl<sub>3</sub>, DMSO-d<sub>6</sub>) δ 12.0 (br s, 1H); 7.73-7.30 (m, 6H); 7.00 (d, 2H, J = 7Hz); 5.18 (s, 2H); 4.23 (s, 2H); 2.55 (t, 2H, J = 7Hz) 1.63 (t of t, 2H, J = 7,7Hz); 1.31 (t of q, 2H, J = 7,7Hz); 0.84 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>24</sub>H<sub>26</sub>ClN<sub>7</sub>O<sub>2</sub>•(H<sub>2</sub>O)<sub>2</sub>: C, 55.87; H, 5.86. Found: C, 56.01; H, 6.01.

## Example 174

## PART A: Preparation of 2-n-Butyl-4-chloro-1-[(2'-(hydroxymethyl)biphenyl-4-yl)methyl]-5-(methoxymethyl)-imidazole

2-n-Butyl-1-[2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(methoxymethyl)imidazole (5.62 g, 13 mmol, 1 eq) was dissolved in THF (50 mL) and to it was slowly added a 1M lithium aluminum hydride

solution in THF (39.5 mL, 39 mmol, 3 eq). The resultant mixture was refluxed under N<sub>2</sub> for 2 hours and worked up according to Fieser and Fieser, V.1, p. 584 (Steinhardt procedure) to yield 4.68 g of a light yellow oil which slowly crystallized. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.57 (bd, 1H, J = 7Hz); 7.47-7.20 (m, 5H); 7.03 (d, 2H, J = 9Hz); 5.18 (s, 2H); 4.58 (s, 2H); 4.32 (s, 2H); 3.28 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.67 (t of t, 2H, J = 7, 7Hz); 1.35 (t of q, 2H, J = 7,7Hz); 0.86 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>23</sub>H<sub>27</sub>ClN<sub>2</sub>O<sub>2</sub>: C, 69.25; H, 6.82; Cl, 8.89. Found: C, 69.42; H, 6.87; Cl, 8.65.

PART B: Preparation of 2-n-Butyl-4-chloro-1-[(2'-(cyanomethyl)biphenyl-4-yl)methyl]-5-(methoxymethyl)-imidazole

2-n-Butyl-4-chloro-1-[(2'-(hydroxymethyl)biphenyl-4-yl)methyl]-5-(methoxymethyl)imidazole (4.68 g) was converted to the title cyanomethyl compound by the procedure described in Example 1, Part B. Work up yielded 5.20 g of a brown oil which was further reacted with purification. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.54 (m, 1H); 7.40 (m, 2H); 7.28 (m, 3H); 7.08 (d, 2H, J = 10Hz); 5.23 (s, 2H); 4.33 (s, 2H); 3.63 (s, 2H); 3.30 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.70 (t of t, 2H, J = 7,7Hz); 1.37 (t of q, 2H, J = 7,7Hz); 0.90 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>24</sub>H<sub>25</sub>ClN<sub>3</sub>O: 407.1764. Found: 407.1778.

PART C: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[(2'-(tetrazol-5-yl)methyl)biphenyl-4-yl]-methylimidazole

2-n-Butyl-4-chloro-1-[(2'-(cyanomethyl)biphenyl-4-yl)methyl]-5-(methoxymethyl)imidazole (5.20 g) was converted to the above tetrazole in 2 days using the procedure of Example 90, Part C. Work-up and flash chromatography over silica gel eluting with a gradient solvent system of 1:1 hexane/ethyl acetate to 1:1 ethyl acetate/isopropanol yielded 3.13 g of a light yellow solid; m.p. 149.0-152.5°. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.37-7.15 (m, 6H); 6.96 (d, 2H, J = 9Hz); 5.18 (s, 2H); 4.30 (s, 2H); 4.24 (s, 2H); 3.27 (s, 3H); 2.57 (t, 2H, J = 7Hz); 1.56 (t of t, 2H, J = 7,7Hz); 1.28 (t of q, 2H, J = 7,7Hz); 0.77 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>24</sub>H<sub>27</sub>ClN<sub>6</sub>O: C, 63.97, H, 6.03; Cl, 7.86. Found: C, 63.79; H, 6.04; Cl, 7.70.

Example 175

Preparation of 2-n-Butyl-1-[(2'-(carboxymethyl)biphenyl-4-yl)methyl]-4-chloro-5-(hydroxymethyl)-imidazole•dicyclohexylamine salt

2-n-Butyl-4-chloro-1-[(2'-(cyanomethyl)biphenyl-4-yl)methyl]-5-(methoxymethyl)imidazole (2.60 g) and a 1:1 mixture of concentrated aqueous HCl and glacial acetic acid (50 mL) were mixed together and then refluxed for 6 hours. The solvents were removed in vacuo and water (200 mL) was added to the residue. The pH was adjusted to 3 with concentrated NH<sub>4</sub>OH and this aqueous mixture was extracted with ethyl acetate (3 x 200 mL). The organic layers were combined, dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to yield an oil. Subsequent flash chromatography in 60:40 ethyl acetate/hexane to 100% isopropanol yielded 1.07 g of a glass. This product was dissolved in acetone and dicyclohexylamine was added (1 eq). A gum precipitated which was redissolved with more acetone (total of 75 mL) and heat. Upon cooling, solid precipitate was obtained (291 mg); m.p. 135.0-137.0. NMR shows -OCH<sub>3</sub> to be missing. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.43-7.13 (m, 6H); 6.95 (d, 2H, J = 8Hz); 5.20 (s, 2H); 4.46 (s, 2H); 3.45 (s, 2H); 2.76 (m, 2H); 2.60 (t, 2H, J = 7Hz); 2.00-1.03 (m, 24H); 0.87 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>23</sub>H<sub>25</sub>ClN<sub>2</sub>O<sub>3</sub>: 412.1554. Found: 412.1544.

Example 176

PART A: Preparation of 2-n-Butyl-4-chloro-1-[(2'-(hydrazido)biphenyl-4-yl)methyl]-5-(methoxymethyl)-imidazole

2-n-Butyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(methoxymethyl)imidazole (2.00 g, 4.7 mmol, 1 eq), hydrazine (1.5 mL, 46.8 mmol, 10 eq) and methanol (30 mL) were mixed together and then refluxed for 3 days after which 1.5 mL more of hydrazine was added and the reaction refluxed for another day. More hydrazine (1.5 mL) was again added and the reaction was refluxed for an additional day. The reaction was worked up by first removing the hydrazine and methanol in vacuo, following by taking up the residue in ethyl acetate (200 mL) and washing it with water (3 x 100 mL). The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to yield 1.37 g of a white glass. NMR (CDCl<sub>3</sub>, 200 MHz,) δ 7.67-

7.31 (m, 4H); 7.40 (d, 2H, J = 9Hz); 7.03 (d, 2H, J = 9Hz); 7.56 (bs, 1H); 5.17 (s, 2H); 4.27 (s, 2H); 3.25 (s, 3H); 2.57 (t, 2H, J = 7Hz); 1.70 (t of t, 2H, 7,7Hz); 1.34 (t of q, 2H), J = 7,7Hz); 0.86 (t, 3H, J = 7Hz). Anal. Calcd. for  $C_{23}H_{27}ClN_4O_2$ : C, 64.70; H, 6.37; N, 13.12. Found: C, 64.47; H, 6.35; N, 12.85.

5 PART B: Preparation of 2-n-Butyl-4-chloro-5-methoxymethyl-1-[4-(2-(trifluoromethylsulfonylhydrazido)-biphenyl-4-yl)methyl]imidazole

A solution of triflic anhydride (0.42 mL, 2.5 mmol, 1.5 eq) in methylene chloride (2 mL) was slowly dripped into a stirred solution at  $-78^{\circ}\text{C}$  of 2-n-butyl-4-chloro-1-[(2'-(hydrazido)biphenyl-4-yl)methyl]-5-(methoxymethyl)imidazole (0.71 g, 1.7 mmol, 1.0 eq) and triethylamine (0.35 mL, 2.5 mmol, 1.5 eq) in methylene chloride (5 mL). The solution was stirred at  $-78^{\circ}\text{C}$  for 1 hour and then allowed to warm to room temperature. After 2 hours at room temperature, water (100 mL) was added, the pH adjusted to 5 and the aqueous layer extracted with ethyl acetate (3 x 100 mL). The organic layers were dried ( $\text{MgSO}_4$ ), the solvent removed in vacuo, and the residue flash chromatographed over silica gel beginning in 1:1 hexane/ethyl acetate and finishing in 100% ethyl acetate to yield 380 mg of a light yellow glass. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.82-7.15 (m, 8H); 6.94 (d, 2H, J = 8Hz); 5.13 (s, 2H); 4.25 (s, 2H); 3.17 (s, 3H); 2.53 (t, 2H, J = 7Hz); 1.69 (t of t, 2H, J = 7,7Hz); 1.27 (t of q, 2H, J = 7,7Hz); 0.81 (t, 3H, J = 7Hz). Fast Atom Bombardment Mass Spectrum: Mass Calcd. for  $C_{24}H_{26}ClF_3N_4O_4S$ : 559.15. Found: 559.12.

20 Example 177

PART A: Preparation of 4'-Methylbiphenyl-2-carboxaldehyde

Methyl 4'-methylbiphenyl-2-carboxylate (20.00 g, 88 mmol, 1 eq) was dissolved in dry toluene (250 mL) and cooled to  $-78^{\circ}$ : Diisobutylaluminum hydride (1.0 M in toluene, 220.0 mL, 220 mmol, 2.2 eq) was then dripped in slowly over 25 minutes keeping the temperature under  $-70^{\circ}$ . When the addition was complete, the mixture was stirred at  $-78^{\circ}$  for 15 minutes and then methanol (10 mL) was added cautiously. When gas evolution was complete, the mixture was poured into a solution of Rochelle salt (100 mL of saturated solution plus 600 mL water). The mixture was stirred or shaken until an extractable solution was obtained. The layers were separated and the aqueous layer extracted with ether (2 x 200 mL). The organic layers were combined, dried ( $\text{MgSO}_4$ ) and the solvent removed in vacuo to yield 16.7 g of a light yellow oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.56-7.16 (m, 8H); 4.59 (s, 2H); 2.40 (s, 3H); 1.74 (s, 1H). This oil (16.7 g, 84 mmol, 1 eq) was subsequently oxidized by dissolving in methylene chloride (100 mL) and stirring with manganese dioxide (7.34 g, 84 mmol, 1 eq). After stirring for one day at room temperature, more manganese dioxide (14.68 g, 168 mmol, 2 eq) was added. The next day, 14.68 g more of manganese dioxide was again added. After another day of stirring, the reaction was filtered through Celite® and the filtrate evaporated to an oil. The oil was chromatographed in 9:1 hexane/ethyl acetate over silica gel to yield 13.4 g of a light yellow opaque oil. The above oxidation can also be performed using pyridinium chlorochromate. NMR ( $\text{CDCl}_3$ , 200 MHz)  $\delta$  9.98 (s, 1H); 8.01 (d, 1H, J = 7Hz); 7.64 (t, 1H, J = 7Hz); 7.53-7.38 (m, 2H); 7.28-7.17 (m, 4H); 2.43 (s, 3H). Mass Calcd. for  $C_{14}H_{12}O$ : 196.0888. Found: 196.0881.

PART B: Preparation of 4'-Methyl-2-(2-nitroethen-1-yl)biphenyl

4'-Methylbiphenyl-2-carboxaldehyde (13.21 g, 67.3 mmol (1.0 eq), nitromethane (4.74 mL, 87.5 mmol, 1.3 eq), ammonium acetate (2.07 g, 26.0 mmol, 0.4 eq) and glacial acetic acid (30 mL) were mixed and refluxed for 2 days, at which time more nitromethane (4.74 mL) and ammonium acetate (2.07 g) were added and the reaction was refluxed for an additional 5 hours. The reaction mixture was poured into ice water (300 mL) and extracted with ethyl acetate (300 mL). The ethyl acetate layer was washed with water (3 x 200 mL), the organic layer dried ( $\text{MgSO}_4$ ), the solvent removed in vacuo and the residue chromatographed in 1:1 hexane/toluene to yield 11.22 g of a light yellow oil which crystallized. The product was recrystallized from methylcyclohexane to yield 8.47 g of yellow crystals; m.p.  $64.0-65.0^{\circ}$ . NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  8.04 (d, 1H, J = 13Hz); 7.69 (d, 1H, J = 9Hz) 7.59-7.37 (m, 4H); 7.50 (d, 1H, J = 13 Hz); 7.27 (d, 2H, J = 7Hz); 7.19 (d, 2H, J = 7Hz); 2.41 (s, 3H). Anal. Calcd. for  $C_{15}H_{13}NO_2$ : C, 75.30; H, 5.48; N, 5.85. Found: C, 75.32; H, 5.56; N, 5.58.

PART C: Preparation of 4'-methyl-2-(1,2,3-triazol-4-yl)biphenyl

4'-Methyl-2-(2-nitroethen-1-yl)biphenyl (6.58 g, 27.5 mmol, 1 eq), sodium azide (5.40 g, 82.3 mmol, 3 eq), and dimethylsulfoxide (minimum to dissolve everything) were mixed together and stirred at room temperature for 4.5 hours. Ethyl acetate (500 mL) was then added and the organic phase washed with water (3 x 400 mL). The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to yield 6.54 g of an orange glass. Chromatography in 75:25 hexane/ethyl acetate yielded 2.87 g of a yellow glass. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.83 (m, 1H); 7.51-7.32 (m, 3H); 7.18 (d, 2H, J = 8Hz); 7.13 (d, 2H, J = 8Hz); 7.03 (s, 1H); 2.38 (s, 3H). Mass Calcd. for C<sub>15</sub>H<sub>13</sub>N<sub>3</sub>: 235.1110. Found: 235.1111.

PART D: Preparation of 4'-Methyl-2-(N-(triphenylmethyl)-1,2,3-triazol-4-yl)biphenyl

4'-Methyl-2-(1,2,3-triazol-4-yl)biphenyl (2.61 g, 11 mmol, 1.0 eq), triethylamine (1.69 mL, 12 mmol, 1 eq), tritylbromide (3.88 g, 12 mmol, 1 eq) and methylene chloride (30 mL) were mixed and stirred at 0 °C and then allowed to warm to room temperature. After 1 hour, ethyl acetate was added (200 mL) and the organic phase was washed with water (3 x 200 mL). The organic layer was dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to yield 5.15 g of a yellow solid. This product was recrystallized from methylcyclohexane to give 3.26 g of off-white crystals; m.p. 181.0-182.5 °. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  8.18 (d, 1H, J = 7Hz); 7.50-7.16 (m, 12H); 7.05-6.89 (m, 10 Hz); 6.47 (s, 1H); 2.54 (s, 3H). Anal. Calcd. for C<sub>34</sub>H<sub>27</sub>N<sub>3</sub>: C, 85.50; H, 5.70; N, 8.80. Found: C, 86.60; H, 5.80; N, 8.94.

PART E: Preparation of 2-n-Butyl-4-chloro-5-hydroxymethyl-1-[(2'-(N-(triphenylmethyl)-1,2,3-triazol-4-yl)-biphenyl-4-yl)methyl]imidazole

4'-Methyl-2-(N-(triphenylmethyl)-1,2,3-triazol-4-yl)biphenyl (3.14 g, 6.57 mmoles) was brominated in the benzylic position by the procedure in Example 85, Part B, using benzoylperoxide instead of AIBN as radical initiator. Filtration of succinimide and evaporation yielded 4.45 g of a crude oil which was used as is. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  CH<sub>2</sub>Br, 4.41. This bromide (4.33 g, approx. 7.8 mmol, 1 eq) was alkylated onto 2-n-butyl-4-chloro-5-(hydroxymethyl)imidazole by the procedure described in Example 1, Part A. Flash chromatography in 75:25 hexane/ethyl acetate over silica gel yielded a yellow solid (0.67 g) which was recrystallized from carbon tetrachloride to yield 447 mg of white crystals; m.p. 173.0-176.5 °. NMR (CDCl<sub>3</sub>, 200 MHz)  $\delta$  8.03 (d, 1H, J = 9Hz); 7.51-7.14 (m, 14H); 6.98 (m, 6H); 6.86 (d, 2H, J = 9Hz); 6.63 (s, 1H); 5.15 (s, 2H); 4.33 (s, 2H); 2.53 (t, 2H, J = 7Hz); 1.15 (t of t, 2H, J = 7.7Hz); 1.32 (t of q, 2H, J = 7.7Hz); 0.87 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>42</sub>H<sub>38</sub>ClN<sub>5</sub>O: 663.2765. Found: 663.2762.

PART F: Preparation of 2-n-Butyl-4-chloro-5-hydroxymethyl-1-[(2'-(1,2,3-triazol-4-yl)biphenyl-4-yl)methyl]imidazole

2-n-Butyl-4-chloro-5-hydroxymethyl-1-[(2'-(N-(triphenylmethyl)triazol-4-yl)biphenyl-4-yl)methyl]imidazole (408 mg, 0.6 mmol, 1 eq), 1,4-dioxane (5 mL), water (1 mL) and 4.0 N HCl in dioxane (0.46 mL, 1.8 mmol, 3 eq) were mixed and stirred at room temperature. After 2 hours, water was added (200 mL), and the aqueous layer extracted with ethyl acetate (3 x 200 mL). The organic layers were dried (MgSO<sub>4</sub>) and the solvent removed in vacuo to yield 260 mg of an off-white glass. Flash chromatography of the product in 100% ethyl acetate over silica gel yielded 140 mg of a white glass. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.82 (m, 1H); 7.50-7.25 (m, 3H); 7.17 (d, 2H, J = 9Hz); 6.98 (d, 2H, J = 9Hz); 6.95 (s, 1H); 5.23 (s, 2H); 4.52 (s, 2H); 2.58 (t, 2H, J = 7Hz); 1.63 (t of t, 2H, J = 7.7Hz); 1.30 (t of q, 2H, J = 7.7Hz); 0.82 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>23</sub>H<sub>24</sub>ClN<sub>5</sub>O: 421.1669. Found: 421.1670.

Examples 178 and 179PART A: Preparation of Ethyl 3-(4-methylphenyl)-3-oxo-2-(allyl)propanoate

Ethyl 3-(4-methylphenyl)-3-oxopropanoate (prepared as described in W. Wierenga and H. I. Skulnick, J. Org. Chem. (1979), **44**, 310) (63.66 g, 309 mmol, 1 eq) was added to a freshly prepared sodium ethoxide solution (Na, 7.43 g, 323 mmol, 1.05 eq; EtOH, 250 mL). The ethanol was removed in vacuo and the residue was dissolved in DMF (250 mL). Allyl bromide (29.3 mL, 338 mmol, 1.1 eq) followed by sodium iodide (4.56 g, 304 mmol, 1 eq) were then added and the contents stirred overnight at room temperature. The DMF was removed in vacuo, water (250 mL) was added and the aqueous layer extracted with ethyl



acetate (3 x 200 mL). The organic layers were dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to yield 74.21 g of an amber oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.81 (d, 2H,  $J = 10\text{ Hz}$ ); 7.30 (d, 2H,  $J = 10\text{ Hz}$ ); 5.96-5.72 (m, 1H); 5.21-5.00 (m, 2H); 4.41 (t, 1H,  $J = 7\text{ Hz}$ ); 4.16 (q, 2H,  $J = 7\text{ Hz}$ ); 2.78 (t, 2H,  $J = 7\text{ Hz}$ ); 2.42 (s, 3H); 1.18 (t, 3H,  $J = 7\text{ Hz}$ ). Anal. Calcd. for  $\text{C}_{15}\text{H}_{18}\text{O}_3$ : C, 73.15; H, 7.37. Found: C, 73.10; H, 7.38.

#### PART B: Preparation of 3-Carboethoxy-4-(4-methylphenyl)-4-(oxo)butanal

Ethyl 3-(4-methylphenyl)-3-oxo-2-(allyl)propanoate (74.21 g, 301 mmol, 1.0 eq), osmium tetroxide (100 mg, cat.), sodium metaperiodate (141.8 g, 663 mmol, 2.2 eq), ether (500 mL) and water (1 L) were mixed and stirred at room temperature. After 24 hours, an additional 110 mg of  $\text{OsO}_4$  was added and after another 24 hours, 200 mg more of  $\text{OsO}_4$  was added together with sodium metaperiodate (190 g, 888 mmol, 3.0 eq). After 4 days, the layers were separated and the ether layer washed with aqueous sodium bisulfite (1 x 500 mL) followed by brine (1 x 300 mL). The ether layer was dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to yield 64.99 g of a dark brown oil. This oil was flash chromatographed over silica gel in 4:1 hexane/ethyl acetate to yield 37.5 g of an amber oil. NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  9.79 (s, 1H); 7.93 (d, 2H,  $J = 9\text{ Hz}$ ); 7.27 (d, 2H,  $J = 9\text{ Hz}$ ); 4.87 (t, 1H,  $J = 7\text{ Hz}$ ); 4.13 (q, 2H,  $J = 7\text{ Hz}$ ); 3.37-3.08 (AB multiplet, 2H); 2.40 (s, 3H); 1.14 (t, 3H,  $J = 7\text{ Hz}$ ). Anal. Calcd. for  $\text{C}_{14}\text{H}_{16}\text{O}_4$ : C, 67.73; H, 6.50. Found: C, 67.53; H, 6.54.

#### PART C: Preparation of 3-Carboethoxy-2-(4-methylphenyl)furan

Ethyl 3-Carboethoxy-4-(4-methylphenyl)-4-(oxo)butanal (10.00 g), trifluoroacetic anhydride (50 mL) and trifluoroacetic acid (2 drops) were mixed and stirred at  $0^\circ$  over ice and allowed to warm to room temperature. After 3 hours, more trifluoroacetic anhydride (50 mL) together with trifluoroacetic acid (2 drops) were added at room temperature. The next day, the solvent was removed *in vacuo* and the residue partitioned between 1 N NaOH (200 mL) and ethyl acetate (200 mL). The layers were separated and the organic layer washed with 1 N NaOH (2 x 200 mL). The organic layer was dried ( $\text{MgSO}_4$ ) and the solvent removed *in vacuo* to yield a brown oil (9.95 g) which was flash chromatographed in 99:1 hexane/ethyl acetate to yield 2.57 g of an off-white solid; m.p.  $79.0\text{--}80.5^\circ$ . NMR (200 MHz,  $\text{CDCl}_3$ )  $\delta$  7.88 (d, 2H,  $J = 9\text{ Hz}$ ); 7.42 (d, 1H,  $J = 2\text{ Hz}$ ); 7.26 (d, 2H,  $J = 9\text{ Hz}$ ); 6.83 (d, 1H,  $J = 2\text{ Hz}$ ); 4.34 (q, 2H,  $J = 7\text{ Hz}$ ); 2.40 (s, 3H); 1.34 (t, 3H,  $J = 7\text{ Hz}$ ). Anal. Calcd. for  $\text{C}_{14}\text{H}_{14}\text{O}_3$ : C, 73.03; H, 6.13. Found: C, 73.52; H, 6.30.

#### PART D: Preparation of 2-n-Butyl-1-[4-(3-carboxyfuran-2-yl)benzyl]-4-chloro-5-(hydroxymethyl)imidazole (isomer A) and 2-n-butyl-1-[4-(3-carboxyfuran-2-yl)benzyl]-5-chloro-4-(hydroxymethyl)imidazole (isomer B)

3-Carboethoxy-2-(4-methylphenyl)furan was brominated, alkylated, and saponified by the procedures described in Example 85, Parts B, C, and E.

Isomer A, the faster eluting isomer, was recrystallized from acetonitrile; m.p.  $158.5\text{--}160.0^\circ$ . NMR (200 MHz,  $\text{DMSO}-d_6$ )  $\delta$  12.80 (bm, 1H); 7.92 (d, 2H,  $J = 9\text{ Hz}$ ); 7.82 (d, 1H,  $J = 2\text{ Hz}$ ); 7.17 (d, 2H,  $J = 9\text{ Hz}$ ); 6.84 (d, 1H,  $J = 2\text{ Hz}$ ); 5.30 (s, 2H), 5.30 (m, 1H); 4.34 (s, 2H); 2.47 (t, 2H,  $J = 7\text{ Hz}$ ); 1.47 (t of t, 2H,  $J = 7, 7\text{ Hz}$ ); 1.24 (t of q, 2H,  $J = 7, 7\text{ Hz}$ ); 0.74 (t, 3H,  $J = 7\text{ Hz}$ ). Anal. Calcd. for  $\text{C}_{20}\text{H}_{21}\text{ClN}_2\text{O}_4$ : C, 61.78; H, 5.44; N, 9.12. Found: C, 61.66; H, 5.39; N, 9.09.

Isomer B was recrystallized from nitromethane/acetonitrile; m.p.  $118.5\text{--}120.5^\circ$ . NMR (200 MHz,  $\text{DMSO}-d_6$ )  $\delta$  12.89 (bm, 1H); 7.92 (d, 2H,  $J = 9\text{ Hz}$ ); 7.82 (d, 1H,  $J = 2\text{ Hz}$ ); 7.13 (d, 2H,  $J = 9\text{ Hz}$ ); 6.83 (d, 1H,  $J = 2\text{ Hz}$ ); 5.23 (s, 2H); 4.93 (m, 1H); 4.29 (d, 2H,  $J = 7\text{ Hz}$ ); 2.57 (t, 2H,  $J = 7\text{ Hz}$ ); 1.53 (t of t, 2H,  $J = 7, 7\text{ Hz}$ ); 1.27 (t of q, 2H,  $J = 7, 7\text{ Hz}$ ); 0.77 (t, 3H,  $J = 7\text{ Hz}$ ). Mass Calcd. for  $\text{C}_{20}\text{H}_{21}\text{ClN}_2\text{O}_4$ : 388.1190. Found: 388.1171.

#### Example 180

#### PART A: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethyl)imidazole

To a solution of 7.50 mL of 1.6 M n-butyllithium/hexane in 50 mL of tetrahydrofuran at  $0^\circ$  was added dropwise 1.50 mL of t-butanol. To this solution was added 4.52 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole followed by 1.50 mL of 2-methoxyethoxymethyl chloride. The resulting solution was stirred at  $25^\circ$  for 16 hours. The mixture was diluted with diethyl ether, washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography afforded 3.50 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethyl)imidazole.

ymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.83 (d, 1H); 7.52 (t, 1H); 7.40 (t, 1H), 7.28 (m, 3H); 7.00 (d, 1H); 5.19 (s, 2H); 4.68 (s, 2H); 4.48 (s, 2H); 3.67 (m, 2H); 3.64 (s, 3H); 3.54 (m, 2H); 3.37 (s, 3H); 2.58 (t, 2H); 1.67 (quint., 2H); 1.34 (sext., 2H); 0.88 (t, 3H).

5 PART B: Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole

A solution of 3.15 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole and 2.77 g of potassium methanethiolate in 125 mL of dimethylformamide was stirred at 125° for 4 hours. After cooling the solvent was removed in vacuo, and the residue was dissolved in water. The resulting aqueous solution was washed with diethyl ether, adjusted to pH 3 employing 10% hydrochloric acid, and extracted with methylene chloride. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product was recrystallized from chlorobutane to afford 2.45 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.95 (d, 1H); 7.57 (t, 1H); 7.46 (t, 1H); 7.38 (m, 3H); 7.05 (d, 2H); 5.22 (s, 2H); 4.64 (s, 2H); 4.48 (s, 2H); 3.58 (m, 4H); 3.40 (s, 3H); 2.54 (t, 2H); 1.60 (quint., 2H); 1.32 (sext., 2H); 0.84 (t, 3H).

20 PART C: Preparation of 1-[(2'-Methoxyaminocarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole

A solution of 0.24 ml of oxalyl chloride in 5 mL of chloroform was added dropwise to a solution of 1 mL of dimethylformamide in 4 mL of chloroform at -20°. After this solution had been stirred at -20° for 20 minutes, 0.28 mL of N-methylmorpholine was added followed by 1.21 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole. After another 20 minutes at -20°, 0.55 mL of N-methylmorpholine and 1.35 mL of methoxylamine were added to the mixture. The reaction mixture was warmed slowly to 25°, stirred at 25° for 4 hours, and finally refluxed for 40 hours. After cooling the mixture was diluted with ethyl acetate. The resulting solution was washed with 10% hydrochloric acid, water, 10% sodium bicarbonate solution and brine. Finally the solution was dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography (elution: methanol/chloroform) furnished 0.21 g of 1-[(2'-methoxyaminocarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  7.85 (s, 1H); 7.63 (d, 1H); 7.53-7.33 (m, 5H); 7.05 (d, 2H); 5.20 (s, 2H); 4.67 (s, 2H); 4.47 (s, 2H); 3.63 (m, 5H); 3.55 (m, 2H); 3.36 (s, 3H); 2.56 (t, 2H); 1.67 (m, 2H); 1.32 (m, 2H); 0.87 (t, 3H).

35 PART D: Preparation of 1-[(2'-Methoxyaminocarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole

A solution of 0.20 g of 1-[(2'-methoxyaminocarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole in 60 ml of 1.5 M aqueous tetrafluoroboric acid/acetonitrile was stirred for 20 hours at 25°. The reaction mixture was poured into dilute sodium bicarbonate solution, and the resulting mixture was extracted with diethyl ether. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography (elution: methanol/chloroform) provided 0.11 g of 1-[(2'-methoxyaminocarbonylbiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$  11.31 (br s, 1H); 7.48 (m, 1H); 7.41-7.33 (m, 5H); 7.09 (d, 2H); 5.27 (br s, 3H); 4.32 (d, 2H); 3.44 (s, 3H); 2.49 (t, 2H); 1.48 (quint., 2H); 1.25 (sext., 2H); 0.80 (t, 3H).


50

55

The following compounds were prepared according to the procedures described in the above example.


NMR (200 MHz, DMSO-d<sub>6</sub>)

**Example 181**



δ 11.29 (br s, 1H),  
7.48 (m, 1H), 7.33 (m,  
10H), 7.09 (d, 2H),  
5.27 (d, 2H), 4.67 (s,  
2H), 4.31 (s, 2H), 2.47  
(t, 2H), 1.46 (quint.,  
2H), 1.21 (sext., 2H),  
0.76 (t, 3H).

**Example 182**



δ 10.81 (br s, 1H),  
9.02 (br s, 1H), 7.55-  
7.35 (m, 6H), 7.11 (d,  
2H), 5.28 (br s, 3H),  
4.34 (d, 2H), 2.50 (t,  
2H), 1.49 (quint., 2H),  
1.25 (sext., 2H), 0.78  
(t, 3H).

**Example 183**

**PART A: Preparation of 1-[(2'-Aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole**

This compound was prepared according to the procedure described in Example 141, Part A. From 3.30 g of 1-[(2'-nitrobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole, 1.60 g of iron powder, 3.20 ml of acetic acid, and 160 mL of methanol there was obtained 2.05 g of 1-[(2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.45 (d, 2H); 7.23-7.08 (m, 4H); 6.89-6.77 (m, 2H); 5.27 (s, 2H); 4.55 (br s, 2H); 2.62 (t, 2H); 1.69 (quint., 2H); 1.37 (sext., 2H); 0.88 (t, 3H).

**PART B: Preparation of 1-[(2'-Aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole**

This compound was prepared according to the procedure described in Example 180, Part A. From 2.03 g of 1-[(2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole, 3.75 mL of 1.6 M n-butyllithium/hexane, 0.75 ml of t-butanol, 0.75 ml of 2-methoxyethoxymethyl chloride, and 25 mL of tetrahydrofuran there was obtained 0.84 g of 1-[(2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.42 (d, 2H); 7.19-7.03 (m, 4H); 6.86 (m, 2H); 5.20 (s, 2H); 4.69 (m, 2H); 4.49 (m, 2H); 3.67 (m, 2H), 3.54 (m, 2H); 3.37 (s, 3H); 2.59 (t, 2H); 1.67 (quint., 2H); 1.34 (sext., 2H); 0.87 (t, 3H).



PART C: Preparation of 1-[(2'-Trifluoroacetamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole

To a solution of 0.84 g of 1-[(2'-aminobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole, 0.23 g of 4-dimethylaminopyridine, 1.28 mL of triethylamine, and 10 mL of tetrahydrofuran at 25 ° was added dropwise 1.30 mL of trifluoroacetic anhydride. The reaction mixture was stirred at 25 ° for 4 hours and then was poured into water. The resulting solution was adjusted to pH 4 using 10% hydrochloric acid and extracted with diethyl ether. The combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography afforded 0.96 g of 1-[(2'-trifluoroacetamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.22 (d, 1H); 7.89 (br s, 1H); 7.44 (m, 1H); 7.36-7.29 (m, 4H); 7.12 (d, 2H); 5.23 (s, 2H); 4.68 (s, 2H); 4.49 (s, 2H); 3.65 (m, 2H); 3.54 (m, 2H); 3.37 (s, 3H); 2.56 (t, 2H); 1.67 (quint., 2H); 1.34 (sext., 2H); 0.87 (t, 3H).

PART D: Preparation of 1-[(2'-Trifluoroacetamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole

This compound was prepared according to the procedure described in Example 180, Part D. From 0.96 g of 1-[(2'-trifluoroacetamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-(2-methoxyethoxymethoxymethyl)imidazole there was obtained 0.35 g of 1-[(2'-trifluoroacetamidobiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.24 (d, 1H); 7.89 (br s, 1H); 7.46 (m, 1H); 7.32 (m, 4H); 7.15 (d, 2H); 5.30 (s, 2H); 4.55 (d, 2H); 2.60 (t, 2H); 1.67 (br t, 1H); 1.70 (quint., 2H); 1.36 (sext., 2H); 0.88 (t, 3H).

Example 184

PART A: Preparation of 2-(4-Methylphenoxy)benzoic acid

To a solution of 5.95 g of *p*-cresol and 7.83 g of 2-chlorobenzoic in 50 mL of dimethylformamide at 25 ° was added, in portions, 14.50 g of anhydrous potassium carbonate. The resulting mixture was heated to 80 °, and 0.10 g of copper(I) iodide was added. The reaction mixture then was refluxed for 16 hours. While still hot the mixture was poured onto water-ice. The resulting suspension was filtered, and the filtrate was adjusted to pH 3.0 using aqueous hydrochloric acid. The precipitate was recovered by filtration. The crude solid was dissolved in an aqueous sodium hydroxide solution. This solution was acidified to pH 6.0 using hydrochloric acid, filtered, and then acidified to pH 3.0. Filtration provided 5.67 g of 2-(4-methylphenoxy)benzoic acid which was employed in the following reaction without further purification. NMR (200 MHz, CDCl<sub>3</sub>): δ 8.15 (d of d, 1H); 7.42 (d of d of d, 1H); 7.23-7.12 (m, 3H); 6.97 (d, 2H); 6.80 (d, 1H); 2.37 (s, 3H).

PART B: Preparation of Methyl 2-(4-methylphenoxy)benzoate

A solution of 37.70 g of 2-(4-methylphenoxy)benzoic acid was 12.0 mL of concentrated sulfuric acid in 500 mL of methanol was refluxed for 14 hours. After cooling, the reaction mixture was concentrated in vacuo and the residue was added to a mixture of methylene chloride and water. The organic phase was separated, washed with saturated sodium bicarbonate solution and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product was kugelrohr distilled (120-135 °/0.025 torr) to furnish 35.08 g of methyl 2-(4-methylphenoxy)benzoate, m.p. 31-34 °. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.87 (d of d, 1H); 7.39 (t of d, 1H); 7.11 (m, 3H); 6.88 (m, 3H); 3.81 (s, 3H); 2.30 (s, 3H).

PART C: Preparation of Methyl 2-(4-bromomethylphenoxy)benzoate

A solution of 35.08 g of methyl 2-(4-methylphenoxy)benzoate, 25.7 g of N-bromosuccinimide, 0.57 g of azobisisobutyronitrile, and 1200 mL of carbon tetrachloride was refluxed for 3 hours. After cooling to room temperature the resulting suspension was filtered and then concentrated in vacuo to provide 4.51 g of crude methyl 2-(4-bromomethylphenoxy)benzoate which was used in a subsequent reaction without further purification; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.92 (d of d, 1H); 7.45 (t of d, 1H); 7.16 (m, 3H); 6.90 (m, 3H); 4.49 (s, 2H); 3.83 (s, 3H).

PART D: Preparation of 2-Butyl-4-chloro-1-[4-(2-carbomethoxyphenoxy)benzyl]-5-hydroxymethylimidazole

To a suspension of 7.51 g of sodium methoxide in 100 mL of dimethylformamide at 25 ° was added a solution of 26.50 g of 2-butyl-4(5)-chloro-5(4)-hydroxymethylimidazole in 100 mL of DMF. The resulting mixture was stirred at 25 ° for 0.25 hours; to this mixture was added dropwise a solution of 45.1 g of methyl 2-(4-bromomethylphenoxy)benzoate in 100 mL of DMF. Finally, the reaction mixture was stirred at 40 ° for 4 hours. After cooling to 25 °, the solvent was removed in vacuo. The residue was dissolved in ethyl acetate, and this solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 10-25% ethyl acetate/benzene) afforded 7.80 g of 2-butyl-4-chloro-1-[4-(2-carbomethoxyphenoxy)benzyl]-5-hydroxymethylimidazole. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.92 (d, 1H); 7.48 (t, 1H); 7.21 (t, 1H); 6.93 (m, 5H); 5.21 (s, 2H); 4.48 (s, 2H); 3.79 (s, 3H); 2.56 (t, 2H); 1.65 (quint., 2H); 1.34 (sext., 2H); 0.88 (t, 3H).

PART E: Preparation of 2-Butyl-4-chloro-1-[4-(2-carboxyphenoxy)benzyl]-5-hydroxymethylimidazole

A solution of 7.70 g of 1-[4-(2-carbomethoxyphenoxy)benzyl]-2-butyl-4-chloro-5-hydroxymethyl imidazole in 250 mL of ethanol and 125 mL of 10% aqueous sodium hydroxide was refluxed for 5 hours. After cooling, the reaction mixture was filtered, and the solvent was removed in vacuo. The residue was dissolved in water, and the solution was acidified to pH 3.5 using hydrochloric acid. The precipitated solid was recovered by filtration and recrystallized from acetone to furnish 6.52 g of 2-butyl-4-chloro-1-[4-(2-carboxyphenoxy)benzyl]-5-hydroxymethylimidazole, m.p. 178-180 °. NMR (200 MHz, DMSO) δ 7.79 (d, 1H); 7.53 (t, 1H); 7.23 (t, 1H); 7.07 (d, 2H); 6.94 (d, 1H); 6.87 (d, 2H); 5.18 (s, 2H); 4.32 (s, 2H); 2.47 (t, 2H); 1.46 (quint., 2H); 1.23 (sext., 2H); 0.78 (t, 3H).

The following compounds have been or could be prepared by the above procedures.

Table 12

5

10

15

20

25

30










35

40

45

50

55

Ex. No.	R <sup>6</sup>				MP (°C)
		R <sup>7</sup>	R <sup>8</sup>		
185	n-butyl	Cl	CH <sub>2</sub> OH		166-167
186	n-butyl	Cl	CH <sub>2</sub> OH		
187	n-butyl	Cl	CH <sub>2</sub> OH		
188	n-propyl	H	CH <sub>2</sub> OH		
189	n-propyl	Cl	CH <sub>2</sub> OH		
190	CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	Cl	CH <sub>2</sub> OH		
191	n-butyl	Cl	CH <sub>2</sub> OH		

Example 192PART A: Preparation of 1-(4-Benzyloxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole

5 To a suspension of 1.43 g of sodium methoxide in 20 mL of dimethylformamide at 25 ° was added a solution of 5.00 g of 2-butyl-4(5)-chloro-5(4)-hydroxymethylimidazole in 15 mL of dimethylformamide (DMF). The resulting mixture was stirred at 25 ° for 0.25 hours, and then to this mixture was added dropwise a solution of 4-benzyloxybenzyl chloride in 15 mL of DMF. Finally, the reaction mixture was stirred at 40 °, the solvent was removed in vacuo. The residue was dissolved in ethyl acetate, and this solution was  
 10 washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 10-25% ethyl acetate/benzene) afforded 3.27 g of 1-(4-benzyloxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole; m.p. 115-116 °; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.39 (m, 5H); 6.94 (s, 4H); 5.15 (s, 2H); 5.04 (s, 2H); 4.47 (bs, 2H); 2.56 (t, 2H); 2.07 (bs, 1H); 1.63 (quint., 2H); 1.32 (sext., 2H); 0.87 (t, 3H).

15

PART B: Preparation of 1-(4-Hydroxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole

A mixture of 0.50 g of 1-(4-benzyloxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole, 0.50 g of 10% palladium/carbon and 40 mL of tetrahydrofuran was stirred at room temperature under hydrogen gas (1  
 20 atm.) for 6 hours. The mixture was filtered through Celite® under nitrogen, and the resulting solution was concentrated in vacuo. The crude product was extracted with hot chloroform. After cooling, the chloroform mixture was concentrated in vacuo, and the resulting solid was washed with hexane to afford 0.16 g of 1-(4-hydroxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole; NMR (200 MHz, DMSO-d<sub>6</sub>): δ 9.43 (s, 1H); 6.81 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.21 (t, 1H); 5.10 (s, 2H); 4.33 (d, 2H); 2.47 (t, 2H); 1.44 (quint 2H); 1.23 (sext., 2H); 0.79 (t, 3H).

25

PART C: Preparation of 1-[4-(2-Cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-hydroxymethylimidazole

To a solution of 1.00 g of 1-(4-hydroxybenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole in 15 mL of DMF at 25 ° was added 0.185 g of sodium methylate, and the resulting mixture was stirred at 25 ° for 0.25  
 30 hours. To this mixture was then added a solution of 0.80 g of α-bromo-o-tolunitrile in 5 mL of DMF. The reaction mixture was stirred at 25 ° for 16 hours. The solvent was removed in vacuo, and the residue dissolved in ethyl acetate. This solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography on silica gel (elution: 10-25% ethyl acetate/benzene) provided 0.76 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-hydroxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.73-7.59 (m, 3H); 7.44 (m, 1H); 6.96 (s, 4H); 5.23 (s, 2H); 5.14 (s, 2H); 4.50 (d, 2H); 2.57 (t, 2H); 1.66 (quint., 2H); 1.33 (sext., 2H); 0.87 (t, 3H).

35

PART D: 1-[4-(2-Cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-cyanomethylimidazole

40 To a solution of 0.76 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-hydroxymethylimidazole in 20 mL of chloroform at 25 ° was added dropwise 0.95 mL of thionyl chloride and the mixture was stirred at 25 ° for 2 hours. The solvent was removed in vacuo. The residue was dissolved in 20 mL of toluene, and then the toluene was removed in vacuo. Finally, the residue was dissolved in 10 mL of dimethyl sulfoxide, and the resulting solution was added to a solution of 0.71 g of sodium cyanide in 10 mL of dimethylsulfoxide. The mixture was stirred at 25 ° for 1 hour and then poured into water. This emulsion was extracted with  
 45 ethyl acetate; and the combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution 0-25% ethyl acetate/benzene) afforded 0.67 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-cyanomethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.79-7.60 (m, 3H); 7.47 (m, 1H); 7.00 (s, 4H); 5.24 (s, 2H);  
 50 5.14 (s, 2H); 3.46 (s, 2H); 2.66 (t, 2H); 1.71 (quint., 2H); 1.40 (sext., 2H); 0.92 (t, 3H).

PART E: 1-[4-(2-Carboxybenzyloxy)benzyl]-2-butyl-4-chloroimidazole-5-acetic acid

A solution of 0.65 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-cyanomethylimidazole in 20  
 55 mL of ethylene glycol and 10 mL of 10% aqueous sodium hydroxide was refluxed for 14 hours. After cooling, the reaction mixture was filtered, and the solvent was removed in vacuo. The residue was dissolved in water, and the solution was acidified to pH 3.5 using hydrochloric acid. The precipitated solid was recovered by filtration and recrystallized from aqueous ethanol to furnish 0.21 g of 1-[4-(2-carboxyben-

zyloxy)benzyl]-2-butyl-4-chloroimidazole-5-acetic acid, m.p. 170-172 °; NMR (200 MHz, DMSO-d<sub>6</sub>): δ 12.9 (bs, 2H); 7.94 (d, 1H); 7.61 (d, 1H); 7.60 (t, 1H); 7.46 (t, 1H); 6.99 (s, 4H); 5.45 (s, 2H); 5.11 (s, 2H); 3.49 (s, 2H); 2.52 (t, 2H); 1.48 (quint., 2H); 1.24 (sext., 2H); 0.82 (t, 3H).

#### 5 Example 193

##### PART A: Preparation of 1-(4-Hydroxybenzyl)-2-butyl-5-hydroxymethylimidazole

A mixture of 1.00 g of 10% palladium/carbon and 1.00 g of 1-(4-benzyloxybenzyl)-2-butyl-4-chloro-5-  
10 hydroxymethyl imidazole in 20 mL of methanol was stirred at 25 ° for five minutes. Hydrogen gas was bubbled into the solution, and the mixture was stirred under hydrogen gas (1 atm.) at 25 ° for 2 hours. The mixture was filtered, and the resulting solution concentrated in vacuo to furnish 0.75 g of 1-(4-hydroxybenzyl)-2-butyl-5-hydroxymethylimidazole; NMR (200 MHz, DMSO-d<sub>6</sub>): δ 9.75 (bs, 1H); 7.55 (s, 1H); 6.91 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.80 (bs, 1H); 5.35 (s, 2H); 4.45 (s, 2H); 2.89 (t, 2H); 1.44 (quint, 2H); 1.21 (sext., 2H); 0.80 (t, 3H).

##### PART B: Preparation of 1-[4-(2-Carboxybenzyloxy)benzyl]-2-butyl-5-hydroxymethylimidazole

The title compound was prepared from 1-(4-hydroxybenzyl)-2-butyl-5-hydroxymethylimidazole using the alkylation and hydrolysis procedures described in Example 192, Parts C and E, m.p. 115-116 °; NMR (200  
20 MHz, DMSO-d<sub>6</sub>): δ 7.92 (d, 1H); 7.59 (m, 2H); 7.43 (m, 1H); 6.95 (A<sub>2</sub>B<sub>2</sub>, 4H); 6.74 (s, 1H); 5.40 (s, 2H); 5.11 (s, 2H); 4.31 (s, 2H); 2.48 (t, 2H); 1.47 (quint., 2H); 1.23 (sext., 2H); 0.77 (t, 3H).

#### Example 194

##### PART A: Preparation of 1-[4-(2-Cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-methoxymethylimidazole

To a solution of 0.29 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-hydroxymethylimidazole in 8.0 mL of dimethyl sulfoxide at 25 ° was added 0.93 g of potassium t-butoxide followed by 0.060 mL of methyl iodide. The reaction mixture was stirred at 25 ° for 2.5 hours and then was poured into water. The  
30 aqueous emulsion was extracted with ethyl acetate; the organic phases were combined and washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated in vacuo. Column chromatography on silica gel (elution: 5-25% ethyl acetate/benzene) furnished 0.17 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-methoxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.72-7.57 (m, 3H); 7.43 (m, 1H); 6.94 (s, 4H); 5.22 (s, 2H); 5.04 (s, 2H); 4.27 (s, 2H); 3.26 (s, 3H); 2.56 (t, 2H); 1.65 (quint., 2H);  
35 1.33 (sext., 2H); 0.88 (t, 3H).

##### PART B: Preparation of 1-[4-(2-Carboxybenzyloxy)benzyl]-2-butyl-4-chloro-5-methoxymethylimidazole

The title compound was prepared from 1-[4-(2-cyanobenzyloxy)benzyl]-2-butyl-4-chloro-5-methoxymethylimidazole via the hydrolysis procedure described in Example 192, Part E; NMR (200 MHz, DMSO-d<sub>6</sub>): δ 7.91 (d, 1H); 7.57 (m, 2H); 7.42 (m, 1H); 6.97 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.41 (s, 2H); 5.09 (s, 2H); 4.27 (s, 2H); 3.17 (s, 3H); 2.49 (t, 2H); 1.44 (quint, 2H); 1.21 (sext., 2H); 0.79 (t, 3H).

The compounds shown in Table 13 where X = -OCH<sub>2</sub>- were prepared or could be prepared employing the above procedures of Examples 192-194 and procedures previously described.

Table 13

Ex.  
No. R<sup>6</sup>

R<sup>7</sup>

R<sup>8</sup>

MP (°C)

195 n-butyl

Cl

CH<sub>2</sub>OH

(oil)<sup>a</sup>

196 n-butyl

Cl

CH<sub>2</sub>OH

197 n-butyl

Cl

CH<sub>2</sub>OCH<sub>2</sub>CH<sub>3</sub>

198 n-butyl

Cl

CH<sub>2</sub>OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>

199 n-butyl

Cl

CH<sub>2</sub>OC(=O)CH<sub>3</sub>

(oil)<sup>b</sup>

200 CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>-  
F F

Cl

CH<sub>2</sub>OH

201 n-propyl

CF<sub>3</sub>

CH<sub>2</sub>OH



- a NMR (200 MHz, DMSO- $d_6$ ):  $\delta$  7.91 (d, 1H);  
 7.58 (m, 2H); 7.42 (m, 1H); 6.98 ( $A_2B_2$ ,  
 4H); 5.42 (s, 2H); 5.15 (s, 2H); 4.32 (s,  
 2H); 2.48 (t, 2H); 1.44 (quint., 2H); 1.23  
 (sext., 2H); 0.79 (t, 3H).
- b NMR (200 MHz,  $CDCl_3$ ):  $\delta$  8.13 (d, 1H);  
 7.75 (d, 1H); 7.58 (t, 1H); 7.39 (t, 1H);  
 6.88 ( $A_2B_2$ , 4H); 5.51 (s, 2H); 5.04 (s,  
 2H); 4.95 (s, 2H); 2.60 (t, 2H); 1.83 (s,  
 3H); 1.65 (quint., 2H); 1.32 (sext., 2H);  
 0.85 (t, 3H).

## 20 Example 202

### PART A: Methyl 2-[4-(Bromomethyl)benzoyl]benzoate

Methyl 2-toluybenzoate (CA reg. # 6424-25-5; available by simple esterification of commercially  
 25 available 2-toluybenzoic acid) (10.00 g, 39.3 mmol, 1 eq), N-bromosuccinimide (7.00 g, 39.3 mmol, 1 eq),  
 benzoyl peroxide (1.0 g) and 100 mL carbon tetrachloride were mixed and refluxed overnight (peroxide  
 added last). The mixture was filtered and 250 mL of a 100 g/l aqueous solution of sodium bisulfite solution  
 was added. The layers were separated and the organic layer was dried ( $MgSO_4$ ) and concentrated. The  
 brown solid residue was recrystallized from ether/hexane to give 6.47 g of product; m.p. 88.2-91.5°. NMR  
 30 (200 MHz,  $CDCl_3$ )  $\delta$  8.07 (d, 1H,  $J$  = 7Hz); 7.82-7.07 (m, 7H); 4.50 (s, 2H); 3.67 (s, 3H). Anal. Calcd. for  
 $C_{16}H_{13}O_3Br$ : C, 57.68; H, 3.93; Br, 23.98. Found: C, 57.84; H, 4.04; Br 23.99. Mass Calcd. for  $C_{16}H_{13}O_3Br$ :  
 332.0048. Found: 332.0033.

### PART B: Preparation of 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-hydroxymethylimidazole

35 To a solution of 2-butyl-4-chloro-5-(hydroxymethyl)imidazole (11.12 g, 54 mmol, 1 eq) in 200 mL  
 methanol was added dropwise a freshly prepared sodium methoxide solution (1.36 g Na, 59 mmol, 1.1 eq in  
 50 mL MeOH). After stirring for 0.5 hours, the methanol was removed in vacuo and the resultant glass was  
 dissolved in 200 mL DMF. To this mixture was added a solution of methyl 2-[4-(bromomethyl)benzoyl]-  
 40 benzoate (18.00 g, 59 mmol, 1.1 eq) in DMF and the entire contents was stirred overnight under  $N_2$  at room  
 temperature. The solvent was then removed in vacuo and the residue dissolved in 500 mL ethyl acetate and  
 500 mL  $H_2O$ . The layers were separated and the aqueous layer was extracted twice with 500 mL portions of  
 ethyl acetate. The organic layers were dried and concentrated and the crude product flash chromatog-  
 45 raphed to separate the two regioisomers in 60:40 hexane/ethyl acetate over silica gel. The faster moving  
 isomer was isolated to yield 14.72 g of a glassy solid. NMR (200 MHz,  $CDCl_3$ )  $\delta$  8.03 (d, 1H,  $J$  = 7Hz); 7.67  
 (m, 4H); 7.36 (d, 1H,  $J$  = 7Hz); 7.05 (d, 2H,  $J$  = 7Hz); 5.28 (s, 2H); 4.43 (s, 2H); 3.63 (s, 3H); 2.53 (t, 2H,  $J$  =  
 7Hz); 1.60 (t of t, 2H,  $J$  = 7,7Hz); 1.30 (t of q, 2H,  $J$  = 7,7Hz); 0.87 (t, 3H,  $J$  = 7Hz). Mass Calcd. for  
 $C_{25}H_{26}ClF_3N_4O_5S$ : 586.1264. Found: 586.1285.

### 50 PART C: 2-Butyl-1-[4-(2-Carboxybenzoyl)benzyl]-4-chloro-5-(hydroxymethyl)imidazole

2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-hydroxymethylimidazole (500 mg, 1.13 mmol, 1  
 eq), 0.5 N KOH in methanol (2.27 mL, 1.14 mmol, 1 eq), and 0.5 mL of  $H_2O$  were mixed and stirred. After 6  
 hours, water (50 mL) was added and the pH was lowered to 3-5 with conc. HCl. The aqueous mixture was  
 55 extracted with ethyl acetate (3 x 50 mL) and the organic layers were dried ( $MgSO_4$ ) and concentrated to  
 give 200 mg of product; m.p. 90.0-95.0°. NMR (200 MHz,  $CDCl_3$ )  $\delta$  8.05 (d, 1H,  $J$  = 7Hz); 7.48-7.75 (m,  
 4H); 7.37 (d, 1H,  $J$  = 7Hz); 7.00 (d, 2H,  $J$  = 7Hz); 5.20 (s, 2H); 4.40 (s, 2H); 2.45 (t, 2H,  $J$  = 7Hz); 1.50 (t of t,  
 2H,  $J$  = 7Hz); 1.25 (t of q, 2H,  $J$  = 7Hz); 0.79 (t, 3H,  $J$  = 7Hz). Anal. Calcd. for  $C_{23}H_{23}ClN_2O_4 \cdot (CH_3OH)$ : C,

62.81; H, 5.93; Found: C, 62.95; H, 5.99. Mass spectrum shows M-H<sub>2</sub>O. Mass Calcd. for C<sub>23</sub>H<sub>23</sub>ClN<sub>2</sub>O<sub>4</sub>-H<sub>2</sub>O: 408.1235. Found: 408.1228.

### Example 203

5

#### Preparation of 2-n-Butyl-1-[4-(2-carboxybenzoyl)benzyl]-4-hydroxymethyl-5-chlorimidazole

Using the procedure of Example 202, 2-n-butyl-1-[4-(2-carboxybenzoyl)benzyl]-4-hydroxymethyl-5-chlorimidazole was prepared from 2-n-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-hydroxymethyl-5-chlorimidazole, m.p. 214.0-216.0°. NMR (200 MHz, CDCl<sub>3</sub> + DMSO-d<sub>6</sub>) δ 8.07 (d, 1H, J = 7.7Hz); 7.32 (d, 1H, J = 7Hz); 7.10 (d, 2H, J = 7Hz); 5.19 (s, 2H); 4.50 (s, 2H); 2.61 (t, 2H, J = 7Hz); 1.63 (t of t, 2H, J = 7.7Hz); 1.33 (t of q, 2H, J = 7.7Hz); 0.87 (t, 3H, J = 7Hz). Titration of the product with 1.000 N NaOH showed the presence of exactly one acidic functionality. Anal. Calcd. for C<sub>23</sub>H<sub>23</sub>ClN<sub>2</sub>O<sub>4</sub>: C, 64.71; H, 5.43; N, 6.56. Found: C, 64.75; H, 5.30; N, 6.65.

15

### Example 204

#### PART A: Preparation of 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-(chloromethyl)imidazole, hydrochloride salt

20

2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-hydroxymethylimidazole (5.00 g, 11.3 mmol, 1 eq) was dissolved in 50 mL chloroform and to this solution was dropwise added thionyl chloride (4.13 mL, 56.6 mmol, 5 eq) with stirring at room temperature. After 4 hours, the solvent and excess thionyl chloride were removed by rotary evaporation. Toluene (100 mL) was added to the residue and the solvent again removed by rotary evaporation. Toluene was again added and while evaporating the second time, product crystallized from solution yielding 2.91 g of a white solid; m.p. 139.0-143.5°. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.07 (d, 1H, J = 7Hz); 7.80 (d, 2H, J = 10Hz); 7.68 (t, 1H, J = 7Hz); 7.58 (t, 1H, J = 7Hz); 7.35 (d, 1H, J = 7Hz); 7.13 (d, 2H, J = 10Hz); 5.43 (s, 2H); 4.42 (s, 2H); 3.67 (s, 3H); 2.96 (m, 2H); 1.75 (m, 2H); 1.39 (m, 2H); 0.88 (t, 2H, J = 7Hz). Mass Calcd. for C<sub>24</sub>H<sub>24</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>3</sub>: 458.1162. Found: 458.1160.

30

#### PART B: 2-Butyl-1-[4-(2-Carbomethoxybenzoyl)benzyl]-4-chloro-5-((1,2,4-triazol-1-yl)methyl)imidazole


2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-chloromethylimidazole•HCl salt (1.00 g, 2.06 mmol, 1.0 eq), potassium triazolidine (0.26 g, 2.39 mmol, 1.1 eq) and DMF (50 mL) were mixed and heated at 90° under N<sub>2</sub> overnight. The reaction was worked up by removing the solvent in vacuo, taking up the residue in water (200 mL) and ethyl acetate (200 mL), separating the layers and extracting the aqueous with ethyl acetate (2 x 200 mL). The organic layers were dried (MgSO<sub>4</sub>) and concentrated; the residue was flash chromatographed over silica gel in 100% ethyl acetate to give 780 mg of a white glassy solid. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.05 (s, 1H); 8.05 (d, 1H, J = 7Hz); 7.83 (s, 1H); 7.74 (d, 2H, J = 10Hz); 7.66 (t, 1H, J = 7Hz); 7.58 (t, 1H, J = 7Hz); 7.33 (d, 1H, J = 7Hz); 6.98 (d, 2H, J = 7Hz); 5.37 (s, 2H); 5.15 (s, 2H); 3.69 (s, 3H); 2.56 (t, 2H, J = 7Hz); 1.73 (m, 2H); 1.36 (t of q, 2H, J = 7.7Hz); 0.87 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>26</sub>H<sub>26</sub>ClN<sub>5</sub>O<sub>3</sub>: 491.1722. Found: 491.1816.

The following intermediates were prepared by the above procedure using the appropriate nucleophile, imidazole starting material, and solvent.






45

50

55






<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>R</u>	<u>MP(°C)</u>
n-butyl	Cl	CH <sub>2</sub> -N 		(oil) <sup>a</sup>
n-butyl	Cl	CH <sub>2</sub> N <sub>3</sub>		127.0-129.5
n-butyl	Cl	CH <sub>2</sub> CN		(oil) <sup>b</sup>
n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>		(solid) <sup>c</sup>
<p>a NMR (200 MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H, J= 7Hz); 7.72 (d, 2H, J= 8Hz); 7.65 (t, 1H, J= 7Hz); 7.56 (t, 1H, J= 7Hz); 7.36 (d, 1H, J= 7Hz); 7.33 (bs, 1H); 7.00 (bs, 1H); 6.89 (d, 2H, J= 8Hz); 6.78 (bs, 1H); 4.91 (s, 2H); 4.88 (s, 2H); 3.67 (s, 3H); 2.54 (t, 2H, J= 7Hz); 1.65 (t of t, 2H, J= 7,7Hz); 1.33 (t of q, 2H, J= 7,7Hz); 0.85 (t, 3H, J= 7Hz).</p> <p>b NMR (200 MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H, J= 7Hz); 7.76 (d, 2H, J= 10Hz); 7.64 (t, 1H, J= 7Hz); 7.56 (t, 1H, J= 7Hz); 7.36 (d, 1H, J= 7Hz); 7.06 (d, 2H, J= 10Hz); 5.24 (s, 2H); 3.66 (s, 3H); 3.47 (s, 2H); 2.63 (t, 2H, J= 7Hz); 1.70 (t of t, 2H, J= 7,7Hz); 1.37 (t of q, 2H, J= 7,7Hz); 0.89 (t, 3H, J= 7Hz).</p> <p>c NMR (200 MHz, CDCl<sub>3</sub>) δ 8.05 (d, 1H, J= 8Hz); 7.72 (d, 2H, J= 8Hz); 7.61 (m, 2H); 7.38 (d, 1H, J= 7Hz); 7.04 (d, 2H, J= 7Hz); 5.20 (s, 2H); 4.26 (s, 2H); 3.63 (s, 3H); 3.21 (s, 3H); 2.50 (t, 2H, J= 7Hz); 1.65 (m, 2H); 1.29 (m, 2H); 0.84 (t, 3H, J= 7Hz).</p>				


## PART C: 2-Butyl-1-[4-(2-Carboxybenzoyl)benzyl]-4-chloro-5-((1,2,4-triazol-1-yl)methyl)imidazole

2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-((1,2,4-triazol-1-yl)methyl)imidazole (780 mg, 1.59 mmol, 1 eq), 0.5 N KOH in MeOH (6.34 mL, 3.17 mmol, 2 eq) and methanol (20 mL) were mixed and stirred at 20° under N<sub>2</sub>. After 2.5 hours, one more equivalent of 0.5 N KOH in MeOH was added. After seven hours, the solution was acidified to a pH of 4 with 1 N HCl, and 200 mL each of ethyl acetate and water was added. The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 200 mL). The organic layers were dried (MgSO<sub>4</sub>) and concentrated to give 640 mg of a white glassy solid; m.p. 180.0-188.0°. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.94 (d, 1H, J = 7Hz); 7.74 (s, 1H); 7.65 (s, 1H); 7.55 (d, 2H, J = 7Hz); 7.70-7.50 (m, 3H); 6.67 (d, 2H, J = 7Hz); 5.34 (s, 2H); 5.14 (s, 2H); 2.64 (t, 2H, J = 7Hz); 1.74 (t of t, 2H, J = 7,7Hz); 1.36 (t of q, 2H, J = 7,7Hz); 0.89 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>25</sub>H<sub>24</sub>ClN<sub>5</sub>O<sub>3</sub>•EtOAc: C, 61.53; H, 5.70; N, 12.37. Found: C, 61.72; H, 5.19, N, 12.27.

Examples 205-207 in Table 14 were prepared by the procedure described in Example 203, Part C using the appropriate imidazole starting materials.

Table 14



Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
205	n-butyl	Cl	CH <sub>2</sub> -N 	CO <sub>2</sub> H	(oil) <sup>a</sup>
206	n-butyl	Cl	CH <sub>2</sub> N <sub>3</sub>	CO <sub>2</sub> H	188.0-190.0
207	n-butyl	Cl	CH <sub>2</sub> OCH <sub>3</sub>	CO <sub>2</sub> H	210.0-211.5

<sup>a</sup> NMR (200 MHz, CDCl<sub>3</sub>/D<sub>2</sub>O exchange) δ

9.67 (s, 1H); 7.98 (d, 1H, J = 7Hz); 7.63 (t, 1H, J = 7Hz); 7.55 (t, 2H, J = 7Hz); 7.41 (d, 2H, J = 10Hz); 7.41 (d, 1H, J = 7Hz); 7.09 (s, 1H); 7.08 (s, 1H); 6.70 (d, 2H, J = 10Hz); 5.65 (s, 2H); 5.58 (s, 2H); 2.59 (t, 2H, J = 7Hz); 1.71 (t of t, 2H, J = 7,7Hz); 1.36 (t of q, 2H, J = 7,7Hz); 0.87 (t, 3H, J = 7Hz).

Example 208PART A: Preparation of 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-[(1H-tetrazol-5-yl)methyl]-imidazole

5 The title compound was prepared from 2-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-(cyanomethyl)imidazole by the procedure described in Example 26; NMR (200 MHz, DMSO-d<sub>6</sub>) δ 8.00 (d, 1H, J = 7Hz); 7.78 (t, 1H, J = 7Hz); 7.70 (t, 1H, J = 7Hz); 7.50 (d, 2H, J = 8Hz); 7.46 (d, 1H, J = 7Hz); 7.05 (d, 2H, J = 8Hz); 5.35 (s, 2H); 4.20 (s, 2H); 3.57 (s, 3H); 2.52 (t, 2H, J = 7Hz); 1.52 (t of t, 2H, J = 7,7Hz);  
 10 1.27 (t of q, 2H, J = 7,7Hz); 0.70 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>25</sub>H<sub>25</sub>ClN<sub>5</sub>O<sub>3</sub>: C, 60.91; H, 5.11; N, 17.05. Found: C, 60.84; H, 5.12; N, 16.71. Mass Calcd. for C<sub>25</sub>H<sub>25</sub>ClN<sub>5</sub>O<sub>3</sub>: 492.1686. Found: 492.1614.

PART B: Preparation of 2-Butyl-1-[4-(2-carboxybenzoyl)benzyl]-4-chloro-5-[(1H-tetrazol-5-yl)methyl]-imidazole

15 The title compound was prepared from 2-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-[(1H-tetrazol-5-yl)methyl]imidazole by the procedure described in Example 202, Part C; m.p. 228.0-229.5°. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.98 (d, 1H, J = 7Hz); 7.73 (t, 1H, J = 7Hz); 7.69 (t, 1H, J = 7Hz); 7.55 (d, 2H, J = 8Hz); 7.38 (d, 1H, J = 7Hz); 7.05 (d, 2H, J = 8Hz); 5.32 (s, 2H); 4.16 (s, 2H); 2.50 (t, 2H, J = 7Hz); 1.50 (t of t, 2H, J = 7,7Hz); 1.24 (t of q, 2H, J = 7,7Hz); 0.80 (t, 3H, J = 7Hz). Anal. Calcd. for C<sub>24</sub>H<sub>23</sub>ClN<sub>5</sub>O<sub>3</sub>: C, 60.19; H, 4.84; N, 17.55. Found: C, 59.73; H, 4.61; N, 17.82.


Example 20925 PART A: Preparation of 5-Aminomethyl-2-n-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloroimidazole, chromium salt

5-Azidomethyl-2-n-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloroimidazole (4.24 g, 9.1 mmol, 1 eq), chromium (II) chloride (6.75 g, 54.7 mmol, 6 eq), acetone (40 mL) and water (13 mL) were mixed and  
 30 stirred (the chromium (II) chloride being added last). After N<sub>2</sub> evolution had stopped, the reaction mixture was diluted with saturated aqueous sodium bicarbonate (250 mL) and extracted with ethyl acetate (3 x 250 mL). The organic layers were dried (MgSO<sub>4</sub>) and concentrated to give solids which after washing with ether gave 2.92 g of white solid (chromium salt of the product); m.p. 178.5-181.0°. NMR (200 MHz, CDCl<sub>3</sub>/DMSO-d<sub>6</sub>) δ 8.85 (bs, 1H); 8.05 (d, 1H, J = 7Hz); 7.57-7.25 (m, 4H); 7.36 (d, 1H, J = 7Hz); 7.06 (bd, 2H, J = 7Hz);  
 35 5.67 (bs, 2H); 3.85 (bs, 2H); 3.67 (s, 3H); 2.60 (t, 2H, J = 7Hz); 1.68 (m, 2H); 1.37 (t of q, 2H, J = 7,7Hz); 0.89 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>24</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>3</sub>: 439.1663. Found: 439.1663. Anal. Calcd. for Cr-(C<sub>24</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>3</sub>)<sub>2</sub>: C, 61.87; H, 5.62; N, 9.02. Found: C, 61.46; H, 5.59; N, 8.54.

40 PART B: Preparation of 2-Butyl-4-chloro-1-[4-(2-carbomethoxybenzoyl)benzyl]-5-(methoxycarbonylaminomethyl)imidazole

5-Aminomethyl-2-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloroimidazole (chromium salt) (500 mg, 1.14 mmol, 1 eq) was dissolved in a mixture of 1.00 N NaOH (1.14 mL, 1.14 mmol, 1 eq) and H<sub>2</sub>O (10 mL). Tetrahydrofuran may be added to assist solvation. The solution was cooled to 0° when methyl  
 45 chloroformate (0.176 mL, 2.28 mmol, 2 eq) in THF (5 mL) was slowly dripped in, in five equal portions, alternating with five portions of 1.00 N NaOH (total of 1.14 mL, 1.14 mmol, 1 eq). When the addition was complete, the mixture was stirred at room temperature for 4 hours. Water (100 mL) was added and the pH adjusted to 5 with 1N HCl. The aqueous was extracted with ethyl acetate (3 x 100 mL), the organic layers dried (MgSO<sub>4</sub>) and stripped to give a white glass (560 mg). Flash chromatography in 100% ethyl acetate to  
 50 100% isopropanol yielded 280 mg of product as an oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H, J = 7Hz); 7.75 (d, 2H, J = 7Hz); 7.75-7.56 (m, 2H); 7.39 (d, 1H, J = 7Hz); 7.02 (d, 2H, J = 7Hz); 5.32 (s, 2H); 4.83 (m, 1H); 4.28 (d, 2H, J = 7Hz); 3.70 (s, 3H); 3.57 (s, 3H); 2.58 (t, 2H, J = 7Hz); 1.72 (t of t, 2H, J = 7,7Hz); 1.37 (t of q, 2H, J = 7,7Hz); 0.92 (t, 3H, J = 7Hz). Mass Calcd. for C<sub>26</sub>H<sub>28</sub>ClN<sub>3</sub>O<sub>5</sub>: 497.1717. Found: 497.1699.

55 The following intermediates were prepared or could be prepared by the procedure described in Example 209, Part B from the corresponding 5-(aminoalkyl)imidazole intermediate and the appropriate chloroformate or sulfonyl chloride.



5

10

15

20

25

30

35

40

45

50

55

<u>R<sup>1</sup></u>	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>MP (°C)</u>
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}_2\text{CH}_3$	
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}_2\text{CH}_2\text{CH}_3$	
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}(\text{CH}_3)_2$	
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}_2\text{C}_6\text{H}_5$	
	n-butyl	Cl	$\text{CH}_2\text{NHCOCH}_2\text{C}_6\text{H}_5$	
	n-butyl	Cl	$\text{CH}_2\text{-NH-SO}_2\text{-CH}_3$	163.0-168.0

PART C: Preparation of 2-Butyl-4-chloro-1-[4-(2-carboxybenzoyl)benzyl]-5-(methoxycarbonylaminomethyl)-imidazole

Using the procedure of Example 202, Part C (with or without refluxing), 2-butyl-1-[4-(2-carboxybenzoyl)-  
5 benzyl]-4-chloro-5-(methoxycarbonylaminomethyl)imidazole was prepared from 2-butyl-1-[4-(2-carbomethox-  
ybenzoyl)benzyl]-4-chloro-5-(methoxycarbonylaminomethyl)imidazole; mp = sublimes. NMR (200 MHz,  
DMSO-d<sub>6</sub>)  $\delta$  13.17 (bm, 1H); 7.97 (d, 1H, J = 7Hz); 7.71 (t, 1H, J = 7Hz); 7.63 (t, 1H, J = 7Hz); 7.56 (d, 2H,  
J = 10Hz); 7.50 (m, 1H); 7.36 (d, 1H, J = 7Hz); 7.03 (d, 2H, J = 10Hz); 5.31 (s, 2H); 4.06 (d, 2H, J = 7Hz);  
2.46 (t, 2H, J = 7Hz); 1.48 (t of t, 2H, J = 7,7Hz); 1.22 (t of q, 2H, J = 7,7Hz); 0.78 (t, 3H, J = 7Hz). Anal.  
10 Calcd. for C<sub>25</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>5</sub>: C, 62.05; H, 5.42; N, 8.68. Found: C, 61.97; H, 5.58; N, 8.40. Mass Calcd. for  
C<sub>25</sub>H<sub>26</sub>ClN<sub>3</sub>O<sub>5</sub>: 483.1561. Found: 483.1560.

Examples 210-216 in Table 15 were prepared or could be prepared by the procedure described in  
Example 209, Part C using the appropriate starting material.

15

20

25

30

35


40

45

50

55

Table 15



Ex. No.	<u>R<sup>13</sup></u>	<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>MP (°C)</u>
210	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHCOCH <sub>2</sub> CH <sub>3</sub>	
211	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHCOCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	
212	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHC(=O)OCH(CH <sub>3</sub> ) <sub>2</sub>	
213	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHCOCH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	
214	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHCOC <sub>6</sub> H <sub>5</sub>	
215	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHSC(=O)CH <sub>3</sub>	(oil) <sup>a</sup>
216	CO <sub>2</sub> H	n-butyl	Cl	CH <sub>2</sub> NHCOCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>	

a NMR (200 MHz, CDCl<sub>3</sub>) δ 7.97 (d, 1H, J= 7Hz); 7.71-7.50 (m, 4H); 7.45 (d, 1H, J= 7Hz); 6.95 (d, 2H, J= 8Hz); 5.23 (s, 2H); 4.15 (s, 2H); 2.57 (t, 2H, J= 7Hz); 1.67 (t of t, 2H, J= 7.7Hz); 1.36 (t of q, 2H, J= 7.7Hz); 0.87 (t, 3H, J= 7Hz).

Example 217PART A: Preparation of 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-[(trifluoromethylsulfonamido)methyl]imidazole

5 Triflic anhydride (0.21 mL, 1.25 mmol, 1.1 eq) was slowly added to a pyridine (20 mL) solution of the chromium salt of 5-aminomethyl-2-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloroimidazole (0.50 g, 1.1 mmol, 1.0 eq) at 0°C. The solution was allowed to warm to room temperature. After 1.5 hour, 1.5 equivalents of triflic anhydride were added at 0°. After an additional 4 hours at room temperature, water  
 10 (200 mL) was added and the pH adjusted to 5. The aqueous was extracted with ethyl acetate (3 x 100 mL) and the organic layers dried (MgSO<sub>4</sub>) and concentrated to yield 150 mg of a yellow oil which was used as is for the subsequent hydrolysis step. NMR (200 MHz, CDCl<sub>3</sub>) δ 8.33 (bm, 1H); 7.96 (d, 1H, J = 7Hz); 7.64 (d, 2H, J = 10Hz); 7.56 (t, 1H, J = 7Hz); 7.48 (t, 1H, J = 7Hz); 7.28 (d, 1H, J = 7Hz); 6.92 (d, 2H, J = 10Hz); 5.21 (s, 2H); 4.14 (s, 2H); 3.17 (s, 3H); 2.48 (t, 2H, J = 7Hz); 1.55 (t of t, 2H, J = 7,7Hz); 1.24 (m, 2H); 0.79  
 15 (t, 3H, J = 7Hz).

PART B: Preparation of 2-Butyl-1-[4-(2-carboxybenzoyl)benzyl]-4-chloro-5-[(trifluoromethylsulfonamido)methyl]imidazole

20 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-4-chloro-5-[(trifluoromethylsulfonamido)methyl]imidazole (150 mg, 0.26 mmol, 1 eq), 1.000 N NaOH (0.55 mL, 0.55 mmol, 2.1 eq), methanol (20 mL), and water (0.5 mL) were mixed and stirred for 5 hours at room temperature under N<sub>2</sub>. The solvent was removed in vacuo. Water (50 mL) was added and the pH was adjusted to 4 with 1 N HCl. Tan solids precipitated. These were collected and dried to yield 89 mg. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.98 (d, 1H, J = 7Hz); 7.70 (t, 1H, J =  
 25 7Hz); 7.68 (t, 1H, J = 7Hz); 7.63 (d, 2H, J = 10Hz); 7.37 (d, 1H, J = 7Hz); 7.10 (d, 2H, J = 10Hz); 5.34 (s, 2H); 4.20 (s, 2H); 2.50 (t, 2H, J = 7Hz); 1.49 (t of t, 2H, J = 7,7Hz); 1.27 (t of q, 2H, J = 7,7Hz); 0.80 (t, 3H, J = 7Hz). Mass calcd. for C<sub>24</sub>H<sub>23</sub>ClF<sub>3</sub>N<sub>3</sub>O<sub>5</sub>S: 557.0999. Found: 557.0988

Example 218


30 PART A: Preparation of 2-Butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-5-[(4-carbomethoxy-1,2,3-triazol-1-yl)methyl]-4-chloroimidazole and 2-butyl-1-[4-(2-carbomethoxybenzoyl)benzyl]-5-[(5-carbomethoxy-1,2,3-triazol-1-yl)methyl]-4-chloroimidazole

35 5-Azidomethyl-2-butyl-4-chloro-1-[4-(2-carbomethoxybenzoyl)benzyl]imidazole (0.50 g, 1.07 mmol, 1 eq), methyl propiolate (0.95 mL, 10.7 mmol, 10 eq) and toluene (20 mL) were mixed and refluxed under N<sub>2</sub> for 3 hours. The reaction mixture was concentrated and the residue flash chromatographed over silica gel in 75:25 hexane/ethyl acetate. The two regioisomers were separated to give 10 mg of the faster eluting isomer as a glass and 330 mg of the slower as a solid. The slower isomer could be further purified by washing with  
 40 ethyl acetate to give 190 mg of white crystalline solid. Faster eluting isomer: NMR (200 MHz, CDCl<sub>3</sub>) δ 8.06 (d, 1H, J = 8Hz); 7.96 (s, 1H); 7.73-7.54 (m, 4H); 7.37 (d, 1H, J = 8Hz); 6.86 (d, 2H, J = 8Hz); 5.76 (s, 2H); 5.41 (s, 2H); 3.90 (s, 3H); 3.68 (s, 3H); 2.56 (t, 2H, J = 7Hz); 1.67 (t of t, 2H, J = 7,7Hz); 1.35 (t of q, 2H, J = 7,7Hz); 0.86 (t, 2H, J = 7Hz). Mass calcd. for C<sub>28</sub>H<sub>28</sub>N<sub>5</sub>O<sub>5</sub>Cl: 549.1778. Found: 549.1860. Slower eluting isomer: m.p. 163.5-167.0°; NMR (200 MHz, CDCl<sub>3</sub>) δ 8.06 (d, 1H, J = 8Hz); 8.00 (s, 1H); 7.72 (d, 2H, J =  
 45 8Hz); 7.72-7.55 (m, 2H); 7.41 (d, 1H, J = 7Hz); 6.96 (d, 2H, J = 8Hz); 5.40 (s, 2H); 5.23 (s, 2H); 3.95 (s, 3H); 3.69 (s, 3H); 2.58 (t, 2H, J = 7Hz); 1.70 (t of t, 2H, J = 7,7Hz); 1.38 (t of q, 2H, J = 7,7Hz); 0.89 (t, 3H, J = 7Hz). Mass calcd. for C<sub>28</sub>H<sub>28</sub>N<sub>5</sub>O<sub>5</sub>Cl: 549.1778. Found: 549.1763.

The intermediates shown below were prepared or could be prepared by the procedure described in Example 218, Part A using the appropriate starting materials.

50

55



<u>R<sup>6</sup></u>	<u>R<sup>7</sup></u>	<u>R<sup>8</sup></u>	<u>R<sup>13</sup></u>	<u>MP(°C)</u>
n-butyl	Cl		CO <sub>2</sub> CH <sub>3</sub>	(oil) <sup>a</sup> (mixture of 2 regio- isomers)
n-butyl	Cl		CO <sub>2</sub> CH <sub>3</sub>	
n-butyl	Cl		NHSO <sub>2</sub> CF <sub>3</sub>	
n-butyl	Cl		NHSO <sub>2</sub> CF <sub>3</sub>	
n-butyl	Cl		NHSO <sub>2</sub> CF <sub>3</sub>	
n-propyl	H		NHSO <sub>2</sub> CF <sub>3</sub>	
n-propyl	H		NHSO <sub>2</sub> CF <sub>3</sub>	



a NMR (200 MHz,  $\text{CDCl}_3$ ) shows a mixture of  
 2 regioisomers:  $\delta$  8.08 (d, 1H,  $J = 8\text{Hz}$ );  
 7.80–7.55 (m, 4H); 7.44–7.34 (m, 1H); 7.28  
 (s, 1H); 7.00–6.88 (m, 2H); 5.40 (s, 0.5 x  
 2H); 5.32 (s, 0.5 x 4H); 5.29 (s, 0.5 x 2H);  
 3.71 (s, 0.5 x 3H); 3.69 (s, 0.5 x 3H);  
 2.75–2.48 (m, 4H); 1.80–1.21 (m, 8H);  
 1.00–0.81 (m, 6H).


PART B: Preparation of 2-Butyl-1-[4-(2-carboxybenzoyl)benzyl]-5-[(4-carboxy-1,2,3-triazol-1-yl)methyl]-4-chloroimidazole and 2-butyl-1-[4-(2-carboxybenzoyl)benzyl]-5-[(5-carboxy-1,2,3-triazol-1-yl)methyl]-4-chloroimidazole

The slower eluting isomer in Example 218, Part A (190 mg, 0.35 mmol, 1 eq), 0.5 N KOH in methanol (2.76 mL, 1.39 mmol, 4 eq) and 5 mL of water were mixed and refluxed overnight under  $\text{N}_2$ . Water (50 mL) was added and the pH adjusted to 5. The aqueous mixture was extracted with ethyl acetate (3 x 50 mL), the organic fractions dried ( $\text{MgSO}_4$ ) and concentrated to give a residue which was triturated with ether yielding 160 mg of solid product. NMR (200 MHz,  $\text{DMSO-d}_6$  +  $\text{py-d}_5$ )  $\delta$  8.20 (d, 1H,  $J = 8\text{Hz}$ ); 7.86–7.63 (m, 4H); 7.57 (d, 1H,  $J = 8\text{Hz}$ ); 7.43 (s, 1H); 7.04 (d, 2H,  $J = 10\text{Hz}$ ); 6.84 (s, 2H); 6.63 (s, 2H); 2.62 (t, 2H,  $J = 7\text{Hz}$ ); 1.65 (t of t, 2H,  $J = 7,7\text{Hz}$ ); 1.30 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.81 (t, 3H,  $J = 7\text{Hz}$ ). Mass calcd. for  $\text{C}_{26}\text{H}_{24}\text{N}_5\text{O}_5\text{Cl-CO}_2$ : 477.1567. Found: 477.1593.

The faster eluting isomer in Example 218, Part A was hydrolyzed in a similar fashion except that upon acidification in the work-up, solid product precipitated, m.p. 149.0–152.5°. NMR (200 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.02 (s, 1H); 8.02 (d, 2H,  $J = 7\text{Hz}$ ); 7.74 (t, 1H,  $J = 7\text{Hz}$ ); 7.66 (t, 1H,  $J = 7\text{Hz}$ ); 7.50 (d, 2H,  $J = 7\text{Hz}$ ); 7.37 (d, 1H,  $J = 7\text{Hz}$ ); 6.92 (d, 2H,  $J = 7\text{Hz}$ ); 5.83 (s, 2H); 5.42 (s, 2H); 2.52 (t, 2H,  $J = 7\text{Hz}$ ); 1.55 (t of t, 2H,  $J = 7\text{Hz}$ ); 1.28 (t of q, 2H,  $J = 7,7\text{Hz}$ ); 0.78 (t, 3H,  $J = 7\text{Hz}$ ). Mass calcd. for  $\text{C}_{26}\text{H}_{24}\text{N}_5\text{O}_5\text{Cl-CO}_2$ : 477.1567. Found: 477.1479.

Examples in Table 16 were prepared or could be prepared by the procedure described in Example 218, Part B.

Table 16



Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
219	n-butyl	Cl		CO <sub>2</sub> H	(oil) <sup>a</sup> (2 regio-isomers)
220	n-butyl	Cl		CO <sub>2</sub> H	
221	n-butyl	Cl		NHSO <sub>2</sub> CF <sub>3</sub>	
222	n-butyl	Cl		NHSO <sub>2</sub> CF <sub>3</sub>	

a NMR (200 MHz, CDCl<sub>3</sub>) δ 8.03 (m, 1H);  
 7.77-7.42 (m, 5H); 7.33 (s, 1H); 5.36 (s,  
 2H); 5.26 (s, 2H); 2.68-2.45 (m, 4H);  
 1.82-1.48 (m, 4H); 1.42-1.20 (m, 4H);  
 1.00-0.80 (m, 6H).

#### Example 223

##### PART A: Preparation of 1-(4-Formylbenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole

To a solution of 5.05 g of 1-(4-cyanobenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole in 350 mL of benzene at 25° was added dropwise 22.8 mL of diisobutylaluminum hydride (0.15 M in toluene). The mixture was warmed to 45° and stirred for 16 hours. After cooling, the reaction mixture was poured in ice-cold 20% aqueous sulfuric acid. This solution was allowed to warm to 25° and then stirred for 2 hours. The solution was cooled to 0°, neutralized using aqueous sodium hydroxide and extracted with ethyl acetate.

The combined organic phases were washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 0-20% ethyl acetate/benzene) provided 3.60 g of 1-(4-formylbenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$ : 9.96 (s, 1H); 7.47 (A<sub>2</sub>M<sub>2</sub>, 4H); 5.26 (s, 2H); 4.42 (s, 2H); 2.54 (t, 2H); 1.64 (quint., 2H); 1.32 (sext., 2H); 0.86 (t, 3H).

**PART B: Preparation of 1-[(2'-Cyano-trans-stilben-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole**

To a solution of 0.98 g of  $\alpha$ -bromo-o-tolunitrile in 25 mL of dimethylformamide at 25° was added 1.40 g of triphenylphosphine. The mixture was stirred at 80° for 3 hours, then treated with 1.53 g of 1-(4-formylbenzyl)-2-butyl-4-chloro-5-hydroxymethylimidazole, followed immediately by 0.54 g of sodium methoxide, and the mixture was diluted with water and extracted with benzene. The organic phases were combined and washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 0-20% ethyl acetate/benzene) afforded 0.45 g of 1-[(2'-cyano-trans-stilben-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$ : 8.01 (d, 1H); 7.85 (d, 1H); 7.73 (t, 1H); 7.47 (t, 1H); 7.44 (AB, 2H, J = 16.3); 7.38 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.28 (s, 2H); 5.24 (t, 1H); 4.34 (d, 2H); 2.49 (t, 2H); 1.47 (quint., 2H); 1.24 (sext., 2H); 0.79 (t, 3H).

**PART C: 1-[(2'-Carboxy-trans-stilben-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole**

A solution of 0.40 g of 1-[(2'-cyano-trans-stilben-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole in 20 mL of ethylene glycol and 12 mL of 10% aqueous sodium hydroxide was refluxed for 5.5 hours. After cooling, the reaction mixture was filtered, and the solvent was removed *in vacuo*. The residue was dissolved in water, and the solution was acidified to pH 3.5 using hydrochloric acid and the resulting emulsion was extracted with chloroform. The combined organic phases were washed with saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography on silica gel (elution: 5% methanol/chloroform) afforded 0.12 g of 1-[(2'-carboxy-trans-stilben-4-yl)methyl]-2-butyl-4-chloro-5-hydroxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$ : 8.08-8.00 (m, 2H); 7.71 (d, 1H); 7.57-7.47 (m, 3H); 7.34 (t, 1H); 7.01-6.92 (m, 3H); 5.21 (s, 2H); 4.50 (s, 2H); 2.60 (t, 2H); 1.62 (quint, 2H); 1.31 (sext., 2H); 0.03 (t, 3H).

Example 224

PART A: Preparation of N-(4-Benzyloxybenzyl)glycine ethyl ester

To a suspension of 11.0 g of glycine ethyl ester hydrochloride in 100 mL of dimethylformamide at 25° was added 22.0 mL of triethylamine. To the resulting milky suspension was added 9.08 g of 4-benzyloxybenzyl chloride in 50 mL of DMF dropwise over 0.5 hour. The mixture was stirred for 16 hours at 25°. The reaction mixture was diluted with diethyl ether and then filtered to remove the precipitated triethylamine hydrochloride. The resulting solution was concentrated *in vacuo*, and the residue was dissolved in ethyl acetate. The solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Kugelrohr distillation provided 5.90 g of N-(4-benzyloxybenzyl)glycine ethyl ester [bp 160-180° (0.015 torr.)]; NMR (200 MHz, CDCl<sub>3</sub>)  $\delta$ : 7.43-7.27 (m, 5H); 7.06 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.01 (s, 2H); 4.14 (quart., 2H); 3.71 (s, 2H); 3.36 (s, 3H); 2.01 (bs, 1H); 1.24 (t, 3H).

PART B: Preparation of N-(4-Benzyloxybenzyl)-N-formylglycine ethyl ester

A solution of 5.83 g of N-(4-benzyloxybenzyl)glycine ethyl ester, 0.86 mL of formic acid, and 20 mL of xylene was refluxed for 2 hours using a Dean-Stark trap to remove the water produced in the reaction. After cooling, the reaction mixture was washed with 20% aqueous formic acid, water, saturated sodium bicarbonate solution, water, and brine. Finally the mixture was dried over anhydrous sodium sulfate, filtered,

and the filtrate was concentrated to furnish 6.23 g of crude N-(4-benzyloxybenzyl)-N-formyl glycine ethyl ester, used in the following reaction without further purification.


#### PART C: Preparation of 1-(4-Benzyloxybenzyl)-5-carbomethoxy-2-(3H)-imidazolethione

To a suspension of 1.10 g of sodium methoxide in 35 mL of tetrahydrofuran at 10° there was added in one portion, a solution of 6.23 g of N-(4-benzyloxybenzyl)-N-formyl glycine ethyl ester and 3.46 mL of methyl formate in 15 mL of THF. The mixture was stirred at 10° for 1 hour and then at 25° for 16 hours. The solvent was removed in vacuo and the residue dissolved in 36 mL of methanol. To this solution was added 3.57 mL of conc. hydrochloric acid, and the mixture was stirred at 40° for 0.5 hour. A solution of 2.80 g of potassium thiocyanate in 6 mL of water was added, and the resulting mixture was stirred for 16 hours at 40°. Finally, 40 mL of water was added, and the mixture was allowed to cool to 25°. The precipitated solid was recovered by filtration to afford 3.60 g of 1-(4-benzyloxybenzyl)-5-carbomethoxy-2-(3H)-imidazolethione; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  11.25 (bs, 1H); 8.05 (s, 1H); 7.39 (m, 5H); 7.03 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.06 (s, 2H); 4.56 (s, 2H); 3.81 (s, 3H).

#### PART D: Preparation of 1-(4-Benzyloxybenzyl)-2-propylthio-5-carboethoxyimidazole

To 60 mL of ethanol at 25° was added portionwise 0.30 g of sodium metal. After the sodium metal has reacted 3.54 g of 1-(4-benzyloxybenzyl)-5-carbomethoxy-2-(3H)-imidazolethione was added followed immediately by 2.24 mL of 1-iodopropane, and the mixture was stirred at 24° for 3 hours. At this point, the solvent was removed in vacuo, and the residue was dissolved in methylene chloride. This solution was washed with water and brine, dried over anhydrous sodium sulfate, filtered, and concentrated to furnish 3.46 g of crude 1-(4-benzyloxybenzyl)-2-propylthio-5-carboethoxyimidazole, used in a subsequent reaction without further purification; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.77 (s, 1H); 7.45-7.32 (m, 5H); 7.03 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.49 (s, 2H); 5.03 (s, 2H); 4.28 (quart., 2H); 3.20 (t, 2H); 1.32 (t, 3H); 1.02 (t, 3H).

The following intermediates were prepared or could be prepared employing the above procedure.




R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>
n-C <sub>6</sub> H <sub>13</sub> S-	H	CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>
n-C <sub>4</sub> H <sub>9</sub> S-	H	CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>

#### PART E: Preparation of 1-(4-Benzyloxybenzyl)-2-propylthio-5-hydroxymethylimidazole

A solution of 2.05 g of 1-(4-benzyloxybenzyl)-2-propylthio-5-carboethoxyimidazole in 10 mL of tetrahydrofuran was added dropwise to 10 mL of 1M lithium aluminum hydride in THF at 0° such that the reaction temperature remained below 5°. The resulting solution then was stirred at 0° for 1 hour. At this point, the reaction mixture was quenched by sequential dropwise addition of 0.40 mL of water, 0.40 mL of 15% aqueous sodium hydride, and 1.20 mL of water. The resulting suspension was filtered employing

diethyl ether, and the filtrate was concentrated to furnish 1.55 g of 1-(4-benzyloxybenzyl)-2-propylthio-5-hydroxymethylimidazole; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.41-7.29 (m, 5H); 7.03-6.86 (m, 5H); 5.22 (s, 2H); 5.01 (s, 2H); 4.45 (s, 2H); 3.01 (t, 2H); 2.32 (bs, 1H); 1.66 (sext., 2H); 0.97 (t, 3H).

The intermediates shown below were prepared or could be prepared employing the above procedure.




R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>
n-C <sub>6</sub> H <sub>13</sub> S-	H	CH <sub>2</sub> OH
n-C <sub>4</sub> H <sub>9</sub> S-	H	CH <sub>2</sub> OH

#### PART F: Preparation of 1-(4-Hydroxybenzyl)-2-propylthio-5-hydroxymethylimidazole

A solution of 1.40 g of 1-(4-benzyloxybenzyl)-2-propylthio-5-hydroxymethylimidazole in 15 mL of trifluoroacetic acid was refluxed for 0.25 hour. After cooling, the reaction was poured into water containing an excess of sodium bicarbonate, and the resulting emulsion was extracted with ethyl acetate. The combined organic phases were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. Column chromatography on silica gel (elution: 0-5% methanol/chloroform) afforded 0.28 g of 1-(4-hydroxybenzyl)-2-propylthio-5-hydroxymethylimidazole; NMR (200 MHz, DMSO-d<sub>6</sub>):  $\delta$  9.41 (s, 1H); 6.88 (s, 1H); 6.79 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.14 (t, 1H); 5.07 (s, 2H); 4.33 (d, 2H); 2.89 (t, 2H); 1.54 (sext., 2H); 0.88 (t, 3H).

These intermediates were prepared or could be prepared employing the above procedure.





R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>
n-C <sub>6</sub> H <sub>13</sub> S-	H	CH <sub>2</sub> OH
n-C <sub>4</sub> H <sub>9</sub> S-	H	CH <sub>2</sub> OH

STEP G: Preparation of 1-[4-(2-Cyanobenzyloxy)benzyl]-2-propylthio-5-hydroxymethylimidazole

The title compound was prepared from 1-(4-hydroxybenzyl)-2-propylthio-5-hydroxymethylimidazole using the procedure described in Example 192, Part C; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  7.66 (m, 3H); 7.43 (m, 1H); 7.03 (s, 1H); 6.99 (A<sub>2</sub>B<sub>2</sub>, 4H); 5.23 (s, 2H); 5.22 (s, 2H); 4.47 (s, 2H); 3.04 (t, 2H); 1.69 (sext., 2H); 0.98 (t, 3H).



The following 2-mercaptoimidazoles shown below were prepared by the procedure illustrated above.

STEP H: Preparation of 1-[4-(2-Carboxybenzyloxy)benzyl]-2-propylthio-5-hydroxymethylimidazole

A solution of 0.23 g of 1-[4-(2-cyanobenzyloxy)benzyl]-2-propylthio-5-hydroxymethylimidazole in 17 mL of ethylene glycol and 7 mL of 10% aqueous sodium hydroxide was refluxed for 14 hours. After cooling, the reaction mixture was filtered, and the solvent was removed in vacuo. The residue was dissolved in water, and the solution was acidified to pH 3.5 using hydrochloric acid. The precipitated solid was recovered by filtration and recrystallized from aqueous ethanol to furnish 0.094 g of 1-[4-(2-carboxybenzyloxy)benzyl]-2-propylthio-5-hydroxymethylimidazole; NMR (200 MHz, DMSO-d<sub>6</sub>):  $\delta$  13.12 (bs, 1H); 7.93 (d, 1H); 7.58 (m, 2H); 7.45 (m, 1H); 6.99 (A<sub>2</sub>B<sub>2</sub>, 4H); 6.98 (s, 1H); 5.42 (s, 2H); 5.25 (bs, 1H); 5.17 (s, 2H); 4.35 (s, 2H); 2.92 (t, 2H); 1.54 (sext., 2H); 0.89 (t, 3H).

The following 2-mercaptoimidazoles of Table 17 were prepared or could be prepared by the procedure illustrated above.

Table 17

5						
10						
15						
	Ex. NO.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	X	R <sup>13</sup>
20	225	n-C <sub>6</sub> H <sub>13</sub> S-	H	CH <sub>2</sub> OH	4-OCH	
25	226	n-C <sub>4</sub> H <sub>9</sub> S-	H	CH <sub>2</sub> OH	4-OCH <sub>2</sub>	

## Example 227

## PART A: Preparation of 1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazole-5-aldehyde

A mixture of 1 g of 1-(4-nitrobenzyl)-2-butyl-4-chloro-5-hydroxymethyl imidazole and 5 g of activated MnO<sub>2</sub> in CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature for 16 hours. The reaction mixture was filtered through celite and the filtrate was concentrated to give a thick oil which was purified by flash column chromatography on silica gel (Hexane:ethyl acetate=1.5:1 elution). The desired compound was obtained as a colorless solid, 0.76 g; m.p. 88-89°; NMR (200 MHz, CDCl<sub>3</sub>): δ 9.74 (2, 1H); 5.64 (s, 2H); 2.63 (t, 3H, J=7.4 Hz); 1.68 (m, 2H); 1.34 (m, 2H); 0.89 (t, 3H, J=7.3 Hz).

## PART B: Preparation of 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propenoic acid, ethyl ester, E and Z isomers

A mixture of 1.2 g of 1-(4-nitrobenzyl)-2-butyl-4-chloroimidazole-5-aldehyde and 1.5 g of (carboxymethylene)triphenylphosphorane in 50 mL of benzene was refluxed for 2 hours. The reaction mixture was concentrated and the residue was purified by flash column chromatography on silica gel (Hexane:EtOAc=3:1 elution). The major product, the E isomer, was eluted first and was obtained as a thick oil initially which solidified to give an amorphous solid, 1.2 g. The minor product, the Z isomer was eluted next and was isolated as a thick liquid, 85 mg. E isomer: NMR (200 MHz, CDCl<sub>3</sub>): 7.3 and 6.53 (d, 2H, J=16 Hz); 5.3 (s, 2H); 2.62 (t, 2H, J=7.3 Hz); 1.69 (m, 2H); 1.28 (m, 5H); 0.89 (t, 3H, J=7.3 Hz). Z isomer: NMR (200 MHz, CDCl<sub>3</sub>): (key peaks only) δ 6.45 and 6.02 (d, 2H, J=11.8 Hz); 5.17 (s, 2H).

## PART C: Preparation of 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol, E isomer

A solution of 0.5 g of 3-[1-(4-nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propenoic acid, ethyl ester, E isomer in 20 mL of THF was cooled with an ice bath, 1.7 mL of 1.5 M diisopropylaluminum hydride (in toluene) was added slowly. The cooling bath was removed and the reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was then quenched with 3 mL of conc. NH<sub>4</sub>Cl solution and the



mixture was stirred for an additional 30 minutes. During this period an extensive gel-like material formed. The reaction mixture was further diluted with ether and filtered through celite. The filtrate was concentrated and the crude product was purified by flash column chromatography on silica gel (Hexane: EtOAc = 1:1 elution). The desired compound was obtained as a thick liquid; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  6.5-6.15 (m, 2H); 5.21 (s, 2H); 4.25 (d, 2H, J = 4.5 Hz); 2.35 (t, 3H, J = 7.4 Hz); 1.68 (m, 2H); 1.34 (m, 2H); 0.86 (t, 3H, J = 7.4 Hz).

PART D: Preparation of 3-[1-(4-Aminobenzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol, E isomer

A mixture of 0.2 g of 3-[1-(4-nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol, 0.15 g of iron and 0.3 mL of glacial acetic acid in 10 mL of absolute ethanol was refluxed for 1 hour. The reaction mixture was concentrated to dryness and the residue was dissolved in 20 mL of water and the solution was made basic to pH 8 by adding K<sub>2</sub>CO<sub>3</sub>. The mixture was then extracted with ethyl acetate and the ethyl acetate layer was washed with water. The organic layer was concentrated to give a crude product which was purified by flash silica gel column chromatography (ethyl acetate elution). A pure product was obtained as an amorphous solid; NMR (200 MHz, CDCl<sub>3</sub>):  $\delta$  6.76 and 6.62 (dd, 4H, J = 8.5 Hz); 6.42-6.22 (m, 2H); 2.57 (t, 2H, J = 7.3 Hz); 1.65 (m, 2H); 1.33 (m, 2H); 0.87 (t, 2H, J = 7.3 Hz).

PART E: Preparation of 3-[1-(4-(2-Carboxybenzamido)benzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol, E isomer

To a solution of 95 mg of 3-[1-(4-aminobenzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol in 2 mL of CHCl<sub>3</sub> was added 45 mg of phthalic anhydride and the mixture was stirred at room temperature for 1 hour. During this period of time the initially clear solution became turbid and produced solid. The reaction mixture was diluted with 2 mL of ether and the solid was collected by filtration and washed with ether. The desired product was obtained as a tan solid, 115 mg, m.p. 150-151 °; NMR (10% DMSO-d<sub>6</sub>/CDCl<sub>3</sub>):  $\delta$  9.94 (s, 1H); 7.71 and 6.93 (d, 4H, J = 8.3 Hz); 6.36 (m, 2H); 5.1 (s, 2H); 4.18 (d, 2H, J = 3.9 Hz); 2.6 (t, 3H, J = 7.4 Hz); 1.68 (m, 2H); 1.34 (m, 2H); 0.89 (t, 3H, J = 7.4 Hz).

Example 228

PART A: Preparation of 3-[2-Butyl-4-chloro-1-(4-aminobenzyl)imidazol-5-yl]propenoic acid ethyl ester, E isomer

A mixture of 0.5 g of 3-[2-butyl-4-chloro-1-(4-nitrobenzyl)imidazol-5-yl]propenoic acid ethyl ester (E isomer) prepared from Part B of Example 227, 1 g of iron and 2 mL of glacial acetic acid in 30 mL of absolute ethanol was refluxed for 1 hour. The reaction mixture was concentrated to dryness and the residue was dissolved in 50 mL of H<sub>2</sub>O. The aqueous solution was adjusted to pH 8 by K<sub>2</sub>CO<sub>3</sub> and was extracted with ethyl acetate. The crude product obtained upon concentration of the ethyl acetate extract was purified by flash silica gel column chromatography (hexane:ethyl acetate = 1:1 elution). The desired compound was obtained as a thick colorless oil, 0.35 g.

PART B: Preparation of 3-[2-Butyl-4-chloro-1-(4-(2-carboxybenzamido)benzyl)imidazol-5-yl]propenoic acid ethyl ester, E isomer

A mixture of 361 mg of the aniline derivative obtained from Part A and 150 mg of phthalic anhydride in 3 mL of chloroform was stirred at room temperature for 1 hour. The reaction mixture was concentrated and the residue was triturated in ethyl ether. The resulting solid was collected and dried to give a colorless solid, 450 mg, m.p. 180-181 °. NMR (CDCl<sub>3</sub>, 5% DMSO-d<sub>6</sub>):  $\delta$  0.91 (t, 3H, J = 7.1 Hz); 1.1-1.4 (m, 5H); 1.60 (q, 2H, J = 7.3 Hz); 2.71 (t, 2H, J = 8.4 Hz); 4.17 (q, 2H, J = 7.3 Hz); 5.23 (s, 2H); 6.46 + 7.38 (d each, 2H, J = 16.1 Hz); 6.0-8.0 (m, 8H), 10.2 (s, 1H).



Example 229PART A: Preparation of 1-(2'-Carbomethoxybiphenyl-4-yl)methyl-2-butyl-4-chloroimidazole-5-carboxaldehyde

5 A mixture of 0.68 g of the hydroxymethyl precursor prepared in Example 85, Part C and 3.4 g of activated MnO<sub>2</sub> in 30 mL of CHCl<sub>3</sub> was stirred at room temperature for 4 hours. The reaction mixture was then filtered through celite and the filtrate was concentrated to give a thick oily residue which was purified by flash chromatography on silica gel (hexane:ethyl acetate=2:1 elution). The desired aldehyde was  
 10 obtained as a thick colorless oil, 0.5 g; NMR (CDCl<sub>3</sub>): 9.78 (s, 1H); 5.6 (s, 2H); 3.63 (s, 3H); 2.63 (t, 3H, J=7.4 Hz); 1.68 (m, 2H); 1.34 (m, 2H); 0.89 (t, 3H, J=7.4 Hz).

PART B: 4-[1-(2'-Carbomethoxybiphenyl-4-yl)methyl-2-butyl-4-chloroimidazol-5-yl]-3-buten-2-one, E isomer

15 A mixture of 0.5 g of 1-(2'-carbomethoxybiphenyl-4-yl)methyl-2-butyl-4-chloroimidazole-5-carboxaldehyde and .04 g of 1-triphenylphosphoranylidene-2-propanone in 20 mL of benzene was refluxed for 16 hours. The reaction mixture was concentrated to give an oily residue which was purified by flash chromatography on silica gel (hexane:ethyl acetate=1:1 elution). The desired compound was obtained as a thick yellowish liquid, 0.46 g; NMR (200 MHz, CDCl<sub>3</sub>): δ 7.9-6.8 (m, 10H); 5.24 (s, 2H); 3.62 (s, 3H); 3.62 (s,  
 20 3H); 2.69 (t, 2H, J=7.4 Hz); 2.26 (s, 3H); 1.72 (m, 2H); 1.38 (m, 2H); 0.91 (t, 3H, J=7.4 Hz).

PART C: Preparation of 4-[1-(2'-Carbomethoxybiphenyl-4-yl)methyl-2-butyl-4-chloroimidazol-5-yl]-3-buten-2-ol, E isomer

25 A solution of 0.45 g of the compound prepared in Part B in 5 mL of methanol was cooled with ice and 0.2 g of NaBH<sub>4</sub> was added portionwise. After all the NaBH<sub>4</sub> was added the reaction mixture was stirred for 10 minutes. The reaction mixture was concentrated to dryness and the residue was treated with 3 mL of satd. NH<sub>4</sub>Cl and the mixture was stirred at room temperature for 10 min. The mixture was then extracted with ethyl acetate and the ethyl acetate extract was concentrated to give a thick liquid, 0.45 g; NMR (200  
 30 MHz, CDCl<sub>3</sub>): 6.45-6.15 (m, 2H); 5.16 (s, 2H); 4.34 (m, 1H, ); 3.67 (s, 3H).

Example 230PART A: Preparation of 1-(4-nitrobenzyl)-2-butyl-4-chloro-5-(2-phenylethen-1-yl)imidazole, E isomer

35 A solution of 0.4 g of benzyltriphenylphosphonium chloride in 20 mL of dried THF was cooled to -30°. To the above solution was added 0.65 mL of 1.6 M n-BuLi dropwise. As the BuLi was added the solution turned to deep orange color. After stirring for 10 min. at -30°, 0.32 g of 1-(4-nitrobenzyl)-2-butyl-4-chloroimidazole-5-aldehyde was added and the reaction mixture was allowed to warm up to room  
 40 temperature and stirred at room temperature for 2 hours. The reaction mixture was quenched with 2 mL of saturated NH<sub>4</sub>Cl solution and diluted with ethyl acetate, and the ethyl acetate solution was washed with water and a brine solution. Evaporation gave a thick oily residue which was purified by the flash silica gel column chromatography (hexane:ethyl acetate=3:1 elution) to give a thick yellow oil, 0.39 g.

PART B: Preparation of 1-[4-(2-Carboxybenzamido)benzyl]-2-butyl-4-chloro-5-(2-phenylethen-1-yl)imidazole, E isomer

The compound was prepared from the compound of Part A by the procedure described in Example 227, Parts D and E; m.p. 111-113° (dec).

50

Example 231PART A: Preparation of 3-[2-Butyl-4-chloro-1-(4-nitrobenzyl)imidazol-5-yl]-3-propen-1-ol acetate, E isomer

55 A mixture of 1 g of 3-[1-(4-nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propen-1-ol obtained from Part C of Example 227, 1 mL of acetic anhydride and 2 mL of pyridine in 20 mL of CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature for 16 hours. The reaction mixture was diluted with 100 mL of ethyl acetate and the organic layer was washed with H<sub>2</sub>O. The crude product obtained upon concentration of the organic layer was

purified by flash silica gel chromatography (hexane: ethyl acetate = 1:1 elution) to give the desired acetate as a thick colorless oil, 0.95 g.

PART B: Preparation of 3-[2-Butyl-4-chloro-1-(4-aminobenzyl)imidazol-5-yl]-3-propen-1-ol acetate, E isomer

The nitro compound obtained from Part A was reduced to the amino compound by the conditions described in Part D of Example 227. The desired compound was obtained as a colorless thick oil.

PART C: Preparation of 3-[2-Butyl-4-chloro-1-(4-(2-carboxybenzamido)benzyl)imidazol-5-yl]-3-propen-1-ol acetate, E isomer

The phthalamic acid derivative was obtained from the aniline derivative obtained from Part B and phthalic anhydride by the method described in Part E of Example 227. The desired compound was obtained as a colorless solid, m.p. 84-87°.

NMR (CDCl<sub>3</sub>) δ 0.91 (t, 3H, J = 7.1Hz); 1.2 (m, 2H); 1.7 (m, 2H); 2.0 (s, 3H); 2.7 (t, 2H, J = 7.4Hz); 4.57 (d, 2H, J = 5.4Hz); 5.06 (s, 2H); 6.24 (m, 2H); 6.9-8.0 (m, 8H); 8.8 (s, 1H).

Example 232

Preparation of 3-[1-(4-((N-Trifluoromethanesulfonyl)anthranilamido)benzyl)-2-butyl-4-chloroimidazol-5-yl]-3-propen-1-ol acetate, E isomer

A mixture of 0.72 g of 3-[2-butyl-4-chloro-1-(4-aminobenzyl)imidazol-5-yl]-3-propen-1-ol acetate obtained from Example 231, Part B and 0.6 mL of triethylamine in 20 mL of CH<sub>2</sub>Cl<sub>2</sub> was cooled with an ice bath. To this solution was added 0.6 g of *o*-(trifluoromethanesulfonamido)benzoyl chloride dropwise and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was then diluted with 100 mL of ethyl acetate, and the ethyl acetate solution was washed with water, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give a crude product which was purified by a flash silica gel column chromatography (3% acetonitrile in ethyl acetate) to give the desired compound as a solid, 1.05 g, m.p. 156-158°; NMR (200 MHz, CDCl<sub>3</sub>): δ 12.9 (bs, 1H); 8.12-6.91 (m); 6.3 (s); 5.09 (s); 4.61 (d, 2H, J = 4.5 Hz); 2.04 (s, 3H).

Example 233

Preparation of 3-[1-(4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl)-2-butyl-4-chloroimidazol-5-yl]-3-propen-1-ol, E isomer

A mixture of 0.9 g of the compound of Example 232 and 3 mL of 1N NaOH in 6 mL of methanol was stirred at room temperature for 16 hours. The reaction mixture was diluted with 50 mL of water and the aqueous solution was acidified to a pH of 3 with 1N HCl to produce extensive solids which were collected and washed with water. The solids were then dried in vacuo to give 0.85 g of the desired product, m.p. 129-131°; NMR (200 MHz, 5% DMSO-d<sub>6</sub>/CDCl<sub>3</sub>): δ 11.15 (bs, 1H); 8.02-6.95 (m, 8H); 6.5-6.3 (m, 2H); 5.13 (s, 2H); 4.19 (d, 2H, J = 3.5 Hz).

Example 234

PART A: Preparation of 3-[2-Butyl-4-chloro-1-(4-nitrobenzyl)imidazol-5-yl]-2-(carboethoxy)propanoic acid, ethyl ester

The sodium salt of diethyl malonate was generated from 2.5 g of NaH (50% oil dispersion) and 8 mL of diethyl malonate in 100 mL of dried DMF with ice cooling. To the above solution was added 5 g of the chloromethyl compound and the mixture was stirred at room temperature for 3 hours. The reaction mixture was stirred at room temperature for 3 hours. The reaction mixture was concentrated and the residue was diluted with 100 mL of water. The aqueous layer was acidified to a pH of 6 by 1N HCl and the product was extracted with ethyl acetate. The crude product was purified by column chromatography (Hexane:EtOAc = 2:1 elution) which afforded the product as a thick yellow oil, 2.8 g.

PART B: Preparation of 3-[2-Butyl-4-chloro-1-(4-nitrobenzyl)imidazol-5-yl]propanoic acid methyl ester

A mixture of 0.5 g of the compound from Part A in 20 mL of 3N HCl was refluxed for 2 hours. The reaction mixture was cooled and neutralized to a pH of 6 with 4N NaOH solution. The resulting gummy solids were extracted into ethyl acetate and concentrated to give a thick yellow oil, 0.5 g. The propionic acid derivative was dissolved in ethyl ether and was treated with diazomethane in ethyl ether to give a crude methyl ester which was purified by column chromatography (hexane:ethyl acetate = 1:1) which afforded the product as a waxy solid, 0.34 g.

10 PART C: Preparation of 3-[2-Butyl-4-chloro-1-(4-(2-carboxybenzamido)benzyl)imidazol-5-yl]propanoic acid methyl ester

The nitro compound of Part B was reduced to the corresponding amino compound by methods previously described. A mixture of 17 mg of the amino compound and 7.5 g of phthalic anhydride in 1 mL of CHCl<sub>3</sub> was stirred at room temperature for 1 hour. The reaction mixture was concentrated to dryness and the residue was triturated with ether. The resulting solids were collected and washed with ether. The pure product was obtained as a colorless solid, 20 mg, m.p. 150.5-151.5° (dec.).

Example 235

20

Preparation of 3-[2-Butyl-4-chloro-1-(4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl)imidazol-5-yl]propanoic acid methyl ester

Reaction between the amino compound of Example 234, Part C and *o*-(trifluoromethanesulfonamido)-benzoyl chloride using the conditions described in Example 232 produced the title compound as a solid, m.p. 168-172°.

Example 236

30 PART A: Preparation of 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]propanoic acid, N,N-dimethylamide

To a solution of 0.7 g of propionic acid from Part B of Example 234 in 20 mL of methylene chloride was added 0.5 mL of pyridine, 0.16 g of dimethylamine HCl salt and 0.42 g of dicyclohexylcarbodiimide. The mixture was then stirred at room temperature for 16 hours. At the end of the reaction the mixture was filtered through celite and the filtrate was concentrated to give a thick oily product. Thus obtained crude product was purified by flash column chromatography (100% elution) to give a pure product as a thick colorless oil, 0.68 g; NMR (200 MHz, CDCl<sub>3</sub>) δ 2.89 (s, 3H); 2.93 (s, 3H); 5.43 (s, 2H).

40 PART B: Preparation of 3-[1-(4-Aminobenzyl)-2-butyl-4-chloroimidazol-5-yl]propanoic acid, N,N-dimethylamide

The nitro compound from Part A was reduced by the same method described in Part D of Example 227 to give the amino compound as a solid, m.p. 146-148°.

45

PART C: Preparation of 3-[2-Butyl-4-chloro-1-(4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl)imidazol-5-yl]propanoic acid, N,N-dimethylamine amide

The amino compound from Part B was treated with *o*-(trifluoromethanesulfonamido)benzoyl chloride as described in Example 232 to give the trifluoromethylsulfonamide product, m.p. 106-108°.

PART D: Preparation of 3-[2-Butyl-4-chloro-1-(4-(2-carboxybenzamido)benzyl)imidazol-5-yl]propanoic acid, N,N-dimethylamine amide

55 The amino compound from Part B was reacted with phthalic anhydride as described in Part E of Example 227 to give the phthalamic acid derivative, m.p. 139-142°.

Example 237PART A: Preparation of 3-[1-(4-Nitrobenzyl-2-butyl-4-chloroimidazol-5-yl)-2-carboethoxy-2-methylpropanoic acid, ethyl ester

5 A solution of 2 g of the malonate derivative obtained from Part A of Example 234 in 10 mL of dried DMF was cooled with ice. To the solution was added 0.22 g of NaH (50% oil dispersion) and the solution was stirred for 5 minutes before adding 0.3 mL of methyl iodide. The reaction mixture then stirred at room temperature for 2 hours. The reaction mixture was diluted with 400 mL of ethyl acetate and the organic layer was washed with H<sub>2</sub>O and brine. The crude product obtained upon concentration of the organic layer was purified by flash silica gel column chromatography (hexane:ethyl acetate = 1:1 elution) to give a pure compound as a thick colorless oil, 1.8 g.

PART B: Preparation of 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]-2-methylpropanoic acid

15 The malonate derivative from Part A was subjected to the hydrolysis-decarboxylation condition as described in Part B of Example 234. The desired compound was obtained as a thick yellowish liquid.

PART C: Preparation of 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]-2-methylpropanoic acid, isopropyl ester

20 A mixture of 0.38 g of the acid from Part B, 1 mL of isopropyl alcohol and 0.22 g of dicyclohexylcarbodiimide in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue was taken into ethyl acetate. Insoluble material was filtered off and the filtrate was concentrated to give a crude product which was purified by column chromatography (hexane:ethyl acetate = 2:1 elution) to give the desired compound as a thick colorless oil, 0.36 g.

PART D: Preparation of 3-[1-(4-((N-trifluoromethanesulfonyl)anthranilamido)benzyl)-2-methylpropanoic acid, isopropyl ester

30 The title compound was prepared from the ester of Part C by the methods described in Parts B and C of Example 236; m.p. 132-135°.

Examples 238 and 239PART A: Preparation of d and l 3-[1-(4-Nitrobenzyl)-2-butyl-4-chloroimidazol-5-yl]-2-methylpropanoic acid, d-(+)- $\alpha$ -methylbenzylamide

40 A mixture of 0.71 g of the propionic acid derivative from Part B of Example 237, 0.25 mL of d-(+)- $\alpha$ -methylbenzylamine and 0.4 g of dicyclohexylcarbodiimide in 50 mL of CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature for 16 hours. The reaction mixture was concentrated and residue was dissolved in 100 mL of ethyl acetate. Insoluble material was filtered off through celite and the filtrate was concentrated to give a crude product which was purified by silica gel column chromatography (hexane:ethyl acetate = 2:1 elution). Two diastereoisomers were separated as a thick colorless oil, 0.37 g each.

PART B: Preparation of d and l 3-[1-(4-Aminobenzyl)-2-butyl-4-chloroimidazol-5-yl]-2-methylpropanoic acid, d-(+)- $\alpha$ -methylbenzylamide

50 The nitro compound from Part A was reduced by the same method described in Part D of Example 227 to give the amino compound as a thick colorless oil.

PART C: Preparation of d and l 3-[1-(4-(2-Carboxybenzamido)benzyl)-2-butyl-4-chloroimidazol-5-yl]-2-methylpropanoic acid, d-(+)- $\alpha$ -methylbenzylamide

55 Each diastereoisomer of the amino compound from Part B was reacted with phthalic anhydride separately as described in Part E of Example 227, to give the phthalamic acid derivatives, m.p. 188-189.5° and 201-202°, respectively.

Example 240Preparation of 1-[(2'-Carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-carboxylic acid

5 To a solution of 1.03 g of 1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-2-butyl-4-chloro-5-hydrox-  
ymethylimidazole in 10 mL of anhydrous acetic acid at 25 ° was added a solution of 0.62 g of chromium  
trioxide in 10 mL of water. The mixture was stirred at 25 ° for 15 minutes and then poured into water. The  
precipitated solids were recovered by filtration and then dissolved in 50 mL of 1.0 N aqueous sodium  
hydroxide solution. The alkaline solution was allowed to stand at 25 ° overnight and then was acidified to pH  
10 3 with 10% aqueous hydrochloric acid. The precipitated solid was recovered by filtration and recrystallized  
from ethyl acetate to afford 0.10 g of 1-[(2'-carboxybiphenyl-4-yl)methyl]-2-butyl-4-chloroimidazole-5-car-  
boxylic acid (m.p. 186-187 ° (decomp.)). NMR (DMSO-d<sub>6</sub>) δ 12.97 (br s, 2H); 7.68 (d, 1H); 7.53 (t, 1H); 7.41  
(t, 1H); 7.34 (d, 1H); 7.28 (d, 2H); 7.02 (d, 2H); 5.61 (s, 2H); 2.60 (t, 2H); 1.53 (quint., 2H); 1.27 (sext., 2H);  
0.81 (t, 3H).

15

20

25

30

35

40

45

50

55

Examples 241-264 were prepared using procedures illustrated in Examples 227-240.

Table 18

5

10

15

20

25

30


35

40

45

50

55



Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP(°C)
241	n-butyl	Cl			115-120
242	n-butyl	Cl			171.5-172.5
243	n-butyl	Cl			160-162
244	n-butyl	Cl	(CH <sub>2</sub> ) <sub>2</sub> COCH <sub>3</sub>		164-162
245	n-propyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>		
246	n-butyl	Cl	CH <sub>2</sub> CH(CH <sub>3</sub> )CO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>		123-125
247	n-butyl	Cl	(CH <sub>2</sub> ) <sub>3</sub> OAc		124-127

Table 18 (continued)
















Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
248	n-butyl	Cl	(CH <sub>2</sub> ) <sub>3</sub> OAc		64-67
249	n-butyl	Cl			142-144
250	n-butyl	Cl			63-64.5
251	n-butyl	Cl			
252	n-butyl	Cl	CO <sub>2</sub> H		
253	n-pentyl	H	CO <sub>2</sub> H		
254	n-propyl	H			
255	n-propyl	Cl			
256	n-propyl	Cl			



Table 18 (continued)


















Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
257	n-butyl	Cl			
258	n-butyl	Cl			
259	n-butyl	Cl			
260	n-butyl	Cl			
261	n-butyl	Cl			
262	n-butyl	Cl			
263	n-butyl	Cl			
264	n-butyl	Cl			75-76.5



Table 18 (continued)

Ex. No.	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
265	n-butyl	Cl	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H		83-85

## Example 266

**PART A: Preparation of 2-(But-1-en-1-yl)-5-t-butyldimethylsilyloxymethyl-1-[(2'-carbomethoxybiphenyl-4-yl)-methyl]-4-chloroimidazole**

2-(But-1-en-1-yl)-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-(hydroxymethyl)imidazole (1.4 g), t-butyldimethylsilyl chloride (0.55 g), and imidazole (0.5 g) were mixed and stirred in DMF (5 mL) for 18 hours at room temperature. Dilution with ethyl acetate and washing the organic phase with water followed by drying (MgSO<sub>4</sub>), evaporation of the solvent in *vacuo*, and flash chromatography in 3:1 hexane/ethyl acetate yielded 1.5 g of a clear oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.83 (d, 1H); 7.52 (t, 1H); 7.40 (t, 1H); 7.33-7.24 (m, 3H); 7.08 (d, 2H); 6.83 (d of t, 1H); 6.13 (d, 1H); 5.30 (s, 2H); 4.57 (s, 2H); 3.64 (s, 3H); 2.21 (quint., 2H); 1.04 (t, 3H); 0.86 (s, 9H); 0.05 (s, 6H).

**PART B: Preparation of 5-t-Butyldimethylsilyloxymethyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole-2-carboxaldehyde**

2-(But-1-en-1-yl)-5-(t-butyldimethylsilyloxymethyl)-1-[(2-carbomethoxybiphenyl-4-yl)methyl]-4-chlorimidazole (262 mg) was reacted with osmium tetroxide and sodium periodate by the procedure described in Example 178, Part B for 1.5 hours at room temperature. Work-up and flash chromatography in 3:1 hexane/ethyl acetate yielded 200 mg of an amorphous solid. NMR (200 MHz, CDCl<sub>3</sub>) δ 9.74 (s, 1H); 7.84 (d, 1H), 7.54 (t, 1H), 7.43 (t, 1H), 7.34-7.25 (m, 3H), 7.16 (d, 2H) 5.83 (s, 2H), 4.65 (s, 2H), 3.64 (s, 3H), 0.90 (s, 9H), 0.09 (s, 6H).

**PART C: Preparation of 5-t-Butyldimethylsilyloxy-methyl-1-[(2'-carbomethoxybiphenyl-4-yl)-methyl]-4-chloro-2-(*cis*-pent-1-en-1-yl)-imidazole**

5-t-Butyldimethylsilyloxymethyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloroimidazole-2-carboxaldehyde (200 mg) was added all at once to a solution of n-butyldiphenylphosphonium bromide (0.26 g) and potassium t-butoxide (70 mg) in THF at 0°C. The reaction mixture was stirred at room temperature for 15 minutes when it was quenched with saturated aqueous ammonium chloride solution. The mixture was extracted with ethyl acetate, the organic layers washed with water, dried (MgSO<sub>4</sub>) and the solvent removed in *vacuo*. The residue was flash chromatographed in hexane/ethyl acetate (5:1) to yield 100 mg of an oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.85 (d, 1H), 7.54 (t, 1H), 7.42 (t, 1H), 7.35-7.24 (m, 3H), 7.07 (d, 2H), 6.07 (d, 1H), 5.87 (d of t, 1H), 5.28 (s, 2H), 4.59 (s, 2H), 3.64 (s, 3H), 2.69 (quart., 2H), 1.46 (sext., 2H), 0.91 (t, 3H), 0.86 (s, 9H), 0.05 (s, 6H).

**PART D: Preparation of 1-[(2'-Carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-hydroxymethyl-2-(*cis*-pent-1-en-1-yl)imidazole**

5-t-Butyldimethylsilyloxymethyl-1-[(2'-carbomethoxybiphenyl-4-yl)methyl]-4-chloro-2-(cis-pent-1-en-1-yl)-imidazole (100 mg) was desilylated with fluoride by procedures familiar to one skilled in the art. Flash chromatography in 1:1 hexane/ethyl acetate yielded 65 mg of a viscous, colorless oil. NMR (200 MHz, CDCl<sub>3</sub>) δ 7.85 (d, 1H), 7.55 (t, 1H), 7.42 (t, 1H), 7.28 (m, 3H), 7.05 (d, 2H), 6.11 (d, 1H), 5.92 (d of t, 1H), 5.30 (s, 2H), 4.57 (d, 2H), 3.64 (s, 3H), 2.69 (quart., 2H), 1.62 (t, 1H), 1.47 (sext., 2H), 0.92 (t, 1H).

**PART E: Preparation of 1-[(2-Carboxybiphenyl-4-yl)-methyl]-4-chloro-5-hydroxymethyl-2-(cis-pent-1-en-1-yl)imidazole**


1-[2'-Carbomethoxybiphenyl-4-yl)methyl]-4-chloro-5-hydroxymethyl-2-(cis-pent-1-en-1-yl)imidazole (65 mg) was hydrolyzed by a procedure similar to that found in Example 85, Part E. Work-up yielded 45 mg of colorless solids; m.p. 148-150°. NMR (200 MHz, DMSO-d<sub>6</sub>) δ 7.77 (d, 1H); 7.50 (t, 1H); 7.38 (t, 1H); 7.33 (m, 3H); 7.08 (d, 2H); 6.10 (d, 1H); 5.84 (d of t, 1H); 5.32 (s, 2H); 4.47 (s, 2H); 2.65 (quart., 2H), 1.45 (sext., 2H); 0.92 (t, 3H).

Table 19 further illustrates compounds which were made or could be made by the methods described in the specification.

Table 19

5

10



15

20

25

30

35

40

45

50

55

Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
267	1	n-butyl	Cl	CH <sub>2</sub> OH		
268	1	n-propyl	H	CH <sub>2</sub> OH		
269	1	n-butyl	Cl	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>		
270	1	n-pentyl	Cl	CH <sub>2</sub> OH		
271	1	n-butyl	Cl	CH <sub>2</sub> NHC(=O)OC <sub>3</sub> H <sub>7</sub>		
272	2	n-butyl	Cl	CH <sub>2</sub> OH		
273	1	n-propyl	H	CH <sub>2</sub> OH		

Table 19 (continued)









Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP(°C)
274	1	n-butyl	CF <sub>3</sub>	CH <sub>2</sub> OH		
275	1	n-butyl	Cl	CH <sub>2</sub> OH		
276	1	n-butyl	H	CH <sub>2</sub> OH		
276	1	n-hexyl	Cl	CH <sub>2</sub> NHCO <sub>2</sub> CH <sub>3</sub>		
278	1	n-butyl	Cl	CH <sub>2</sub> OH		
279	1	n-butyl	Cl	CH <sub>2</sub> OH		
280	0	n-butyl	Cl	CH <sub>2</sub> OH		
281	1	n-propyl	Cl	CH <sub>2</sub> OH		

Table 19 (continued)








Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
5						
282	1	n-butyl	Cl	CH <sub>2</sub> OH		
10						
283	1	n-butyl	Cl	CH <sub>2</sub> OH		
15						
284	1	n-hexyl	H	CH <sub>2</sub> OH		
20						
285	1	n-butyl	Cl	CH <sub>2</sub> OH		
25						
30						
286	1	n-propyl	H	CH <sub>2</sub> OH		
35						
287	1	n-butyl	Cl	(CH <sub>2</sub> ) <sub>2</sub> F		
40						
288	1	n butyl	Cl	CH <sub>2</sub> OC(=O)NHCH <sub>3</sub>		
45						
50						
55						

Table 19 (continued)











Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
289	1	n-butyl	Cl	$\text{CH}_2\overset{\text{S}}{\underset{\text{O}}{\text{C}}}\text{NHCH}_3$		
290	1	n-propyl	H	$\text{CH}_2\overset{\text{S}}{\underset{\text{O}}{\text{C}}}\text{NHCH}_2\text{CH}_2\text{CH}_3$		
291	1	n-pentyl	H	$\text{CH}_2\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{NHCH}_3$		
292	1	n-butyl	Cl	$(\text{CH}_2)_3\text{F}$		181-182.5
293	1	n-butyl	Cl	$\text{CH}_2\text{ONO}_2$		
293	1	n-butyl	Cl			
295	1	n-butyl	Cl	$\text{CH}_2\text{OH}$		
296	1	n-butyl	Cl	$\text{CH}_2\text{OH}$		
297	1	n butyl	Cl	$\text{CH}_2\text{OH}$		

Table 19 (continued)



















Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP(°C)
298	1	n-butyl	Cl	CH <sub>2</sub> OH	4-SCH <sub>2</sub> - 	
299	1	n-butyl	Cl	CH <sub>2</sub> OH	4-CONH- 	
300	1	n-butyl	Cl	CH <sub>2</sub> OH	4-NHCH <sub>2</sub> - 	
301	1	n-butyl	Cl	CH <sub>2</sub> OH	4-NHC(  )C <sub>6</sub> H <sub>5</sub> - 	
302	1	n-propyl	Cl	CH <sub>2</sub> OH	4-SO <sub>2</sub> NH <sub>2</sub> - 	
303	1	n-pentyl	Cl	CH <sub>2</sub> OH	4-CH <sub>2</sub> NH- 	
304	1	n-hexyl	Cl	CH <sub>2</sub> OH	4-CF=CF- 	
305	1	n-butyl	Cl	CH <sub>2</sub> OH	4-CH=CF- 	
306	1	n butyl	H	CH <sub>2</sub> OH	4-CH <sub>2</sub> CH <sub>2</sub> - 	

Table 19 (continued)

Ex. No.	r	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	R <sup>13</sup>	MP (°C)
307	1	n-butyl	Cl	CH <sub>2</sub> OH		
308	1	n-butyl	Cl	CH <sub>2</sub> OH		
309	1	n-butyl	Cl	CH <sub>2</sub> OH		
310	1	n-butyl	Cl	CH <sub>2</sub> OH		
311	1	n-butyl	Cl	CH <sub>2</sub> OH		
312	1	n-propyl	H	CH <sub>2</sub> OH		
313	1	n-pentyl	Cl	CH <sub>2</sub> OH		
314	1	n-butyl	Cl	CH=CHCH <sub>2</sub> OH		103-104.5

## Utility

The hormone angiotensin II (All) produces numerous biological responses (e.g. vasoconstriction) through stimulation of its receptors on cell membranes. For the purpose of identifying compounds such as



All antagonists which are capable of interacting with the All receptor, a ligand-receptor binding assay was utilized for the initial screen. The assay was carried out according to the method described by [Glossmann et al., *J. Biol. Chem.*, 249, 825 (1974)], but with some modifications. The reaction mixture contained rat adrenal cortical microsomes (source of All receptor) in Tris buffer and 2 nM of  $^3\text{H}$ -All with or without potential All antagonist. This mixture was incubated for 1 hour at room temperature and the reaction was subsequently terminated by rapid filtration and rinsing through glass micro-fibre filter. Receptor-bound  $^3\text{H}$ -All trapped in filter was quantitated by scintillation counting. The inhibitory concentration ( $\text{IC}_{50}$ ) of potential All antagonist which gives 50% displacement of the total specifically bound  $^3\text{H}$ -All is presented as a measure of the affinity of such compound for the All receptor (see Table 20).

The potential antihypertensive effects of the compounds of this invention may be demonstrated by administering the compounds to awake rats made hypertensive by ligation of the left renal artery [Cangiano et al., *J. Pharmacol. Exp. Ther.*, 208, 310 (1979)]. This procedure increases blood pressure by increasing renin production with consequent elevation of All levels. Compounds are administered orally at 100 mg/kg and/or intravenously via a cannula in the jugular vein at 10 mg/kg. Arterial blood pressure is continuously measured directly through a carotid artery cannula and recorded using a pressure transducer and a polygraph. Blood pressure levels after treatment are compared to pretreatment levels to determine the antihypertensive effects of the compounds (See Table 20).

20

25

30

35

40

45

50

55

Table 20

5	Ex. No.	Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
		IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
10	1	1.80	+	NA
	2 (sodium salt)	0.140	+	NA
	3 (sodium salt)	0.420		NA
15	4 (sodium salt)	0.280	+	NA
	5 (sodium salt)	0.190		NA
	6	5.70	NT	
20	7	0.420	+	NA
	8 (sodium salt)	0.790		NA
	9 (sodium salt)	5.80	NT	
	10 (sodium salt)	0.190	NT	
25	11 (sodium salt)	0.380	NA	NA
	12 (sodium salt)	0.030	+	NA
	13 (sodium salt)	6.90	+	NA
30	14	3.20	NT	
	15 (sodium salt)	9.4	+	NA
	16	0.018	+	NA
35	17 (sodium salt)	0.042	+	NA
	18	0.08	+	NA
	19 (sodium salt)	1.70	NT	
	20 (sodium salt)	5.30	NT	
40	21 (sodium salt)	2.10	+	NA
	25	3.90	NT	
	26 (sodium salt)	3.80		NA
45	27 (sodium salt)	1.20	+	+

50

55

Table 20 (continued)

		Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
		IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
5	Ex. No.			
10	28	8.00	NT	
	29	3.10	+	NA
	30 (sodium salt)	0.39	+	+
15	31	0.64		NT
	32 (sodium salt)	0.43		NT
	33	0.940		NT
20	35 (sodium salt)	3.40	+	+
	36 (sodium salt)	0.19	+	NA
	51	2.30	NA	NA
	52	1.10		NT
25	54	7.20	+	
	55	0.930	+	NA
	56	4.40		NT
30	57	4.90	+	NA
	58	8.30	+	NA
	59	3.00	NA	NA
35	60	1.20		NT
	61	5.00		NT
	62 (sodium salt)	9.20		NT
	63 (sodium salt)	3.70		NA
40	64	0.620	+	NA
	65	0.240	+	NA
	66	0.350	+	NA
45	67	1.10	+	NA
	70	2.50	+	NA
	71	2.80		NT
50	72	6.50	+	NA

Table 20 (continued)

Ex. No.	Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
	IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
74 ( <u>trans</u> compound)	3.90	+	NA
( <u>cis</u> compound)	4.50	+	NA
75 (sodium salt)	7.60	+	+
76 (sodium salt)	2.70	+	NA
77 (sodium salt)	5.70	NA	NA
78 (sodium salt)	8.00	+	+
79 (sodium salt)	0.50	+	NA
80 (sodium salt)	0.50	+	+
81 (sodium salt)	0.57	NA	NA
82	6.10		NT
83	6.40		NT
85	0.49	+	+
86	2.90	+	NA
87	2.50		NT
88	1.30		+
89	0.039	+	+
90 (sodium salt)	0.020	+	+
91	0.26	+	NA
92	0.062	+	
93	0.89	+	NA
94	0.280	+	+
95	1.20	+	NA
96	1.10		NT
97	0.270	+	NA
98 (sodium salt)	0.099	+	+

Table 20 (continued)

5	Ex. No.	Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
		IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
10	99	0.090	+	+
	100	0.090	+	+
	102	0.061	+	+
15	105	0.680	+	+
	106	1.90	+	+
	107	1.70	NT	.
	108	0.160	+	+
20	109	0.98	+	+
	110	1.30	+	+
	113	0.020	NT	
25	114	0.050	+	+
	115	0.43	+	+
	116	0.26	+	+
30	117	0.89	+	+
	118	0.089	+	+
	121	0.330	+	+
	123	5.60	+	NA
35	124	1.80	+	NA
	125	0.650	+	+
	126	0.340	+	+
40	127	0.150	+	+
	128	0.08	+	+
	129	0.330	+	+
45	130	0.470	+	+
	132	0.020	+	+
	134	0.180	+	+
	135	1.30	+	+
50	141	0.190	+	+

55

Table 20 (continued)

5	10	Ex. No.	Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
			IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
		144	0.083	+	+
		148 (sodium salt)	0.200	+	+
		149 (sodium salt)	0.450	+	+
	15	150 (sodium salt)	0.200	+	+
		151 (sodium salt)	0.560	+	+
		152 (sodium salt)	0.250	+	+
	20	153 (sodium salt)	0.200	+	+
		154 (sodium salt)	0.60	+	+
		156	0.060	+	
	25	160 (sodium salt)	0.120	+	+
		162 (sodium salt)	0.140	+	+
		165 (sodium salt)	3.00	+	NA
		166 (sodium salt)	0.240	+	NA
	30	171 (sodium salt)	0.600	+	NA
		173 (sodium salt)	0.700	+	
		174 (sodium salt)	0.300	+	NA
	35	175 (DCHA salt)	1.50	+	NA
		176	0.200	+	NA
		177	9.60	+	NA
	40	178	4.20	+	+
		179	4.40	+	NA
		180	2.90	+	NA
		181	4.90	+	NA
	45	182	4.10	+	NA
		183	6.30	+	NA
		184	0.40	+	NA

50

55

Table 20 (continued)

		Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
		IC <sub>50</sub> (μmolar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
5	Ex. No.			
10	185	0.400	+	NA
	192	2.30		NA
	193	0.31	+	NA
15	194	1.20		NT
	195	0.92	+	+
	199	1.80		NA
20	202 (sodium salt)	0.160	+	NA
	203 (sodium salt)	0.340	+	+
	204 (sodium salt)	1.90	+	NA
25	205 (sodium salt)	2.50		NT
	206 (sodium salt)	1.40		NT
	207 (sodium salt)	0.15	+	+
	208 (sodium salt)	0.330	+	NA
30	209 (sodium salt)	0.27		NT
	215 (sodium salt)	0.200	+	NA
	217 (sodium salt)	2.70		NT
35	218 (sodium salt)	2.0		NT
	219	0.68		NT
	223	5.40		NT
40	224	5.90		NT
	227	0.110	+	
	228	0.530		NT
	229	2.10	+	+
45	230	1.60	+	
	231	0.076		NT
	232	0.510	+	

Table 20 (continued)

5	Ex. No.	Angiotensin II Receptor Binding	Antihypertensive Effects in Renal Hypertensive Rats	
		IC <sub>50</sub> ( $\mu$ molar)	Intravenous Activity <sup>1</sup>	Oral Activity <sup>2</sup>
10	233	0.600	+	+
	234	0.064	+	NA
	235	0.160	+	NA
15	236	0.110	+	
	237	0.120	+	NA
	238	0.110	+	NA
20	239	0.092	+	
	241	0.170	+	
	242	0.270	+	
	243	0.200		NT
25	244	0.088	+	
	246	0.120	+	
	247	0.110		NT
30	248	0.250	+	
	249	0.072	+	NA
	250	0.120	+	NA
35	264	0.250	+	+
	265	0.270	+	+
	266	2.30	+	
	292	0.700	+	+
40	314	0.630	+	NA

45

50

55



1      Significant decrease in blood pressure at  
10 mg/kg or less

2      Significant decrease in blood pressure at  
100 mg/kg or less

NA - Not active at 100 mg/kg dosage administered.  
Although many of the compounds tested were not  
active orally, they were active  
intravenously. A few compounds (Examples 10,  
51, 59, 77 and 81) did not produce a  
significant decrease in blood pressure at 10  
mg/kg intravenously, but did produce some  
decrease at that level, and it is expected  
that they would be active intravenously at a  
higher dosage, e.g., 30 mg/kg.

NT - Not tested.

#### Dosage Forms

The compounds of this invention can be administered for the treatment of hypertension according to the invention by any means that effects contact of the active ingredient compound with the site of action in the body of a warm-blooded animal. For example, administration can be parenteral, i.e., subcutaneous, intravenous, intramuscular, or intra peritoneal. Alternatively, or concurrently, in some cases administration can be by the oral route.

The compounds can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but are generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

For the purpose of this disclosure, a warm-blooded animal is a member of the animal kingdom possessed of a homeostatic mechanism and includes mammals and birds.

The dosage administered will be dependent on the age, health and weight of the recipient, the extent of disease, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired. Usually, a daily dosage of active ingredient compound will be from about 1-500 milligrams per day. Ordinarily, from 10 to 100 milligrams per day in one or more applications is effective to obtain desired results. These dosages are the effective amounts both for treatment of hypertension and for treatment of congestive heart failure, i.e., for lowering blood pressure and for correcting the hemodynamic burden on the heart to relieve the congestion.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs, syrups, and suspensions. It can also be administered parenterally, in sterile liquid dosage forms.

Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastro-intestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain

preservatives, such as benzalkonium chloride, methyl- or propylparaben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in this field.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be  
5 illustrated as follows:

#### Capsules

A large number of unit capsules are prepared by filling standard two-piece hard gelatin capsules each  
10 with 100 milligrams of powdered active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

#### Soft Gelatin Capsules

15 A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

#### Tablets

20 A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

#### Injectable

25 A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol. The solution is made to volume with water for  
30 injection and sterilized.

#### Suspension

An aqueous suspension is prepared for oral administration so that each 5 milliliters contain 100  
35 milligrams of finely divided active ingredient, 100 milligrams of sodium carboxymethyl cellulose, 5 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin.

40

45


50

55

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. An antihypertensive compound of the formula:




(I)


wherein

R<sup>1</sup>


is -4-CO<sub>2</sub>H; -4-CO<sub>2</sub>R<sup>9</sup>;



-SO<sub>3</sub>H, -C(CF<sub>3</sub>)<sub>2</sub>OH;




-PO<sub>3</sub>H;




4-NHSO<sub>2</sub>CH<sub>3</sub> ; -4-NHSO<sub>2</sub>CF<sub>3</sub>; -CONHOR<sup>12</sup>; -SO<sub>2</sub>NH<sub>2</sub> ;

5




10




15

20



25

30




35

40

45

50

55




R<sup>2</sup> is H; Cl; Br; I; F; NO<sub>2</sub>; alkyl of 1 to 4 carbon atoms; acyloxy of 1 to 4 carbon atoms; alkoxy of 1 to 4 carbon atoms; CO<sub>2</sub>H; CO<sub>2</sub>R<sup>9</sup>; NHSO<sub>2</sub>CH<sub>3</sub>; NHSO<sub>2</sub>CF<sub>3</sub>; CONHOR<sup>12</sup>; SO<sub>2</sub>NH<sub>2</sub> ;








40


R<sup>3</sup> aryl; or furyl;  
 is H; Cl, Br, I or F; alkyl of 1 to 4 carbon atoms or alkoxy of 1 to 4 carbon atoms;  
 R<sup>4</sup> is CN, NO<sub>2</sub> or CO<sub>2</sub>R<sup>11</sup>;  
 R<sup>5</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, alkenyl or alkynyl of 2 to 4 carbon atoms;  
 R<sup>6</sup> is alkyl of 2 to 10 carbon atoms, alkenyl or alkynyl of 3 to 10 carbon atoms or the same groups substituted with F or CO<sub>2</sub>R<sup>14</sup>; cycloalkyl of 3 to 8 carbon atoms, cycloalkylalkyl, of 4 to 10 carbon atoms; cycloalkylalkenyl or cycloalkylalkynyl of 5 to 10 carbon atoms; (CH<sub>2</sub>)<sub>5</sub>Z(CH<sub>2</sub>)<sub>m</sub>R<sup>5</sup> optionally substituted with F or CO<sub>2</sub>R<sup>14</sup>; benzyl or benzyl substituted on the phenyl ring with 1 or 2 halogens, alkoxy of 1 to 4 carbon atoms, alkyl of 1 to 4 carbon atoms or nitro;  
 50 R<sup>7</sup> is H, F, Cl, Br, I, NO<sub>2</sub>, CF<sub>3</sub> or CN;  
 R<sup>8</sup> is H, CN, alkyl of 1 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, or the same groups substituted with F; phenylalkenyl wherein the aliphatic portion is 2 to 6 carbon atoms; -(CH<sub>2</sub>)<sub>m</sub>-imidazol-1-yl; -(CH<sub>2</sub>)<sub>m</sub>-1,2,3-triazolyl optionally substituted with one or two groups selected from CO<sub>2</sub>CH<sub>3</sub> or alkyl of 1 to 4 carbon atoms;  
 55 -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl;  
 -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>;




5




30



35




40



45

R<sup>9</sup> is

50









R<sup>10</sup> is alkyl of 1 to 6 carbon atoms or perfluoroalkyl of 1 to 6 carbon atoms, 1-adamantyl, 1-naphthyl, 1-(1-naphthyl)ethyl, or (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>;

R<sup>11</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;


55

R<sup>12</sup> is H, methyl or benzyl;

R<sup>13</sup> is -CO<sub>2</sub>H; -CO<sub>2</sub>R<sup>9</sup>; -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>R<sup>9</sup>;




- R<sup>14</sup> is H, alkyl or perfluoroalkyl of 1 to 8 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;
- R<sup>15</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl, benzyl, acyl of 1 to 4 carbon atoms, phenacyl;
- R<sup>16</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>, OR<sup>17</sup>, or NR<sup>18</sup>R<sup>19</sup>;
- R<sup>17</sup> is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;
- R<sup>18</sup> and R<sup>19</sup> independently are H, alkyl of 1 to 4 carbon atoms, phenyl, benzyl, α-methylbenzyl, or taken together form a ring of the formula



Q is  $\text{NR}^{20}$ , O or  $\text{CH}_2$ ;  
 $\text{R}^{20}$  is H, alkyl of 1-4 carbon atoms, or phenyl;  
 $\text{R}^{21}$  is alkyl of 1 to 6 carbon atoms,  $-\text{NR}^{22}\text{R}^{23}$ , or


5



10

$\text{R}^{22}$  and  $\text{R}^{23}$  independently are H, alkyl of 1 to 6 carbon atoms, benzyl, or are taken together as  $(\text{CH}_2)_u$  where u is 3-6;  
 $\text{R}^{24}$  is H,  $\text{CH}_3$  or  $-\text{C}_6\text{H}_5$ ;  
 $\text{R}^{25}$  is  $\text{NR}^{27}\text{R}^{28}$ ,  $\text{OR}^{28}$ ,  $\text{NHCONH}_2$ ,  $\text{NHCSNH}_2$ ,

15




20

$\text{R}^{26}$  is hydrogen, alkyl with from 1 to 6 carbon atoms, benzyl, or allyl;  
 $\text{R}^{27}$  and  $\text{R}^{28}$  are independently hydrogen, alkyl with from 1 to 5 carbon atoms, or phenyl;  
 $\text{R}^{29}$  and  $\text{R}^{30}$  are independently alkyl of 1-4 carbon atoms or taken together are  $-(\text{CH}_2)_q$ ;  
 $\text{R}^{31}$  is H, alkyl of 1 to 4 carbon atoms,  $-\text{CH}_2\text{CH}=\text{CH}_2$  or  $-\text{CH}_2\text{C}_6\text{H}_4\text{R}^{32}$ ;  
 $\text{R}^{32}$  is H,  $\text{NO}_2$ ,  $\text{NH}_2$ , OH or  $\text{OCH}_3$ ;

25

X is a carbon-carbon single bond,  $-\text{CO}-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{NH}-$ ,




30

$-\text{OCH}_2-$ ,  $-\text{CH}_2\text{O}-$ ,  $-\text{SCH}_2-$ ,  $-\text{CH}_2\text{S}-$ ,  $-\text{NHC}(\text{R}^{27})(\text{R}^{28})$ ,  $-\text{NR}^{23}\text{SO}_2-$ ,  $-\text{SO}_2\text{NR}^{23}-$ ,  $-\text{C}(\text{R}^{27})-$ ,  $(\text{R}^{28})\text{NH}-$ ,  $-\text{CH}=\text{CH}-$ ,  $-\text{CF}=\text{CF}-$ ,  $-\text{CH}=\text{CF}-$ ,  $-\text{CF}=\text{CH}-$ ,  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CF}_2\text{CF}_2-$ ,

35



40



45

Y is O or S;  
Z is O,  $\text{NR}^{11}$ , or S;  
m is 1 to 5;  
n is 1 to 10;  
p is 0 to 3;  
q is 2 to 3;  
r is 0 to 2;  
s is 0 to 5;  
t is 0 or 1;

50


and pharmaceutically acceptable salts of these compounds;  
provided that:

(1) the  $\text{R}^1$  group is not in the ortho position



(2) when  $R^1$  is

5



10

$X$  is a single bond, and  $R^{13}$  is  $CO_2H$ , or

15




20

then  $R^{13}$  must be in the ortho or meta position; or when  $R^1$  and  $X$  are as above and  $R^{13}$  is  $NHSO_2CF_3$  or  $NHSO_2CH_3$ ,  $R^{13}$  must be ortho;

(3) when  $R^1$  is

25



30

and  $X$  is other than a single bond, then  $R^{13}$  must be ortho except when  $X = NR^{23}CO$  and  $R^{13}$  is  $NHSO_2CF_3$  or  $NHSO_2CH_3$ , then  $R^{13}$  must be ortho or meta;


35

(4) when  $R^1$  is 4- $CO_2H$  or a salt thereof,  $R^6$  cannot be S-alkyl;

(5) when  $R^1$  is 4- $CO_2H$  or a salt thereof, the substituent on the 4-position of the imidazole cannot be  $CH_2OH$ ,  $CH_2OCOCH_3$ , or  $CH_2CO_2H$ ;

(6) when  $R^1$  is

40



45


$X$  is  $-OCH_2-$ ,

50

$R^{13}$  is 2- $CO_2H$ , and  $R^7$  is  $H$  then  $R^6$  is not  $C_2H_5S$ ;


(7) when  $R^1$  is

55




and  $R^6$  is n-hexyl then  $R^7$  and  $R^8$  are not both hydrogen;

(8) when R<sup>1</sup> is




R<sup>6</sup> is not methoxybenzyl.  
(9) the R<sup>6</sup> group is not





or CH<sub>2</sub>OH.

2. A compound of claim 1 having the formula:



Wherein

R<sup>1</sup> is -CO<sub>2</sub>H; -NHSO<sub>2</sub>CF<sub>3</sub>;




R<sup>6</sup> is alkyl of 3 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, alkynyl of 3 to 10 carbon atoms, cycloalkyl of 3 to 8 carbon atoms, benzyl substituted on the phenyl ring with up to two groups selected from alkoxy of 1 to 4 carbon atoms, halogen, alkyl of 1 to 4 carbon atoms, and nitro;


R<sup>8</sup> is phenylalkenyl wherein the aliphatic portion is 2 to 4 carbon atoms, -(CH<sub>2</sub>)<sub>m</sub>-imidazol-1-yl, -(CH<sub>2</sub>)<sub>m</sub>-1,2,3-triazolyl optionally substituted with one or two groups selected from CO<sub>2</sub>CH<sub>3</sub>

or alkyl of 1 to 4 carbon atoms,  
 $(\text{CH}_2)_m$ -tetrazolyl,  $-(\text{CH}_2)_n\text{OR}^{11}$ ;



5




10



15




20



25


$\text{R}^{13}$  is  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{R}^9$ ,  $\text{NHSO}_2\text{CF}_3$ ; and

30




$\text{R}^{16}$  is H, alkyl of 1 to 5 carbon atoms,  $\text{OR}^{17}$ , or  $\text{NR}^{18}\text{R}^{19}$ ;  
 X is carbon-carbon single bond,  $-\text{CO}-$ ,


35



40



45





$-\text{OCH}_2-$ ,  $-\text{CH}_2\text{O}-$ ,  $-\text{O}-$ ,  $-\text{SCH}_2-$ ,  $-\text{CH}_2\text{S}-$ ,  $-\text{NHCH}_2-$ ,  $-\text{CH}_2\text{NH}-$  or  $-\text{CH}=\text{CH}-$ ; and pharmaceuti-  
 cally acceptable salts of these compounds.

50 3. A compound of claim 2 wherein:

$\text{R}^2$  is H, alkyl of 1 to 4 carbon atoms, halogen, or alkoxy of 1 to 4 carbon atoms;  
 $\text{R}^6$  is alkyl, alkenyl or alkynyl of 3 to 7 carbon atoms;  
 $\text{R}^7$  is H, Cl, Br, I or  $\text{CF}_3$ ;  
 $\text{R}^8$  is  $-(\text{CH}_2)_m\text{OR}^{11}$ ;

55




or  $-\text{COR}^{16}$ ;

$\text{R}^{10}$  is  $\text{CF}_3$ , alkyl of 1 to 6 carbon atoms or phenyl;

$\text{R}^{11}$  is H, or alkyl of 1 to 4 carbon atoms;


$\text{R}^{13}$  is  $\text{CO}_2\text{H}$ ;  $\text{CO}_2\text{CH}_2\text{OCOC}(\text{CH}_3)_3$ ;  $\text{NHSO}_2\text{CF}_3$  and



$\text{R}^{14}$  is H, or alkyl of 1 to 4 carbon atoms;

$\text{R}^{15}$  is H, alkyl of 1 to 4 carbon atoms, or acyl of 1 to 4 carbon atoms;

$\text{R}^{16}$  is H, alkyl of 1 to 5 carbon atoms;  $\text{OR}^{17}$ ; or



$m$  is 1 to 5;

$X$  = single bond,  $-\text{O}-$ ;  $-\text{CO}-$ ;  $-\text{NHCO}-$ ; or  $-\text{OCH}_2-$ ; and pharmaceutically acceptable salts.

4. The compounds of claims 1 to 3, selected from 2-Butyl-4-chloro-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)-methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(methoxycarbonyl)aminomethyl]imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(propoxycarbonyl)aminomethyl]imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a

pharmaceutically acceptable salt thereof;


2-butyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-(1E-butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-hydroxymethylimidazole, or a pharmaceutically acceptable salt thereof; and

2-(1E-butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof.

5. A pharmaceutical composition comprising a pharmaceutically suitable carrier and at least one compound of Claims 1 to 4.


6. A process for the preparation of a compound of claims 1 to 4 wherein r is 1 which comprises contacting an imidazole derivative of Formula 1 with a benzyl derivative of Formula 2 in a solvent in the presence of a base for about 1 to about 10 hours at a temperature in the range of about 20 °C to the reflux temperature of the solvent to form a benzylimidazole of Formula 3:



wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> is stable under the reaction conditions and is a group as defined in claim 1 or an intermediate or protected form thereof which can be transformed to such a group and wherein X<sup>1</sup> is halogen, p-toluenesulfonyloxy or methylsulfonyloxy; and thereafter as necessary transforming said intermediate or protected forms of the R groups to R groups as defined in claim 1.

7. Process of claim 6 wherein compounds 1 and 2 are contacted in the presence of a base selected from the group consisting of a metal hydride, MH, a metal alkoxide, MOR, sodium carbonate, potassium carbonate, triethylamine and pyridine, in a dipolar aprotic solvent or, where the base is MOR, the solvent can be an alcohol, ROH, where M is lithium, sodium or potassium and R is methyl, ethyl or t-butyl.

8. Process of claim 6 wherein: R<sup>1</sup> is




X is a carbon-carbon single bond, -CO-, -O-, -S-, or -NH-;  
R<sup>2</sup> and R<sup>3</sup> are each independently H, Cl, Br, I, CO<sub>2</sub>R<sup>14</sup>, F, NO<sub>2</sub>, alkyl of 1 to 4 carbon atoms, alkoxy of 1 to 4 carbon atoms, aryl or furyl;  
R<sup>6</sup> and R<sup>7</sup> are as defined in claim 1;  
R<sup>8</sup> is alkyl of 1 to 10 carbon atoms or alkenyl of 3 to 10 carbon atoms, or the same groups substituted with F; phenylalkenyl wherein the aliphatic portion is 2 to 6 carbon atoms; -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>; -(CH<sub>2</sub>)<sub>n</sub>SR<sup>15</sup>; or -(CH<sub>2</sub>)<sub>n</sub>CN;

R<sup>11</sup> is as defined in Claim 1  
 R<sup>13</sup> is CO<sub>2</sub>R<sup>14</sup>, CN, NO<sub>2</sub>, trialkyltin tetrazole, or trityltetrazole; and  
 R<sup>14</sup> and R<sup>15</sup> are as defined in Claim 1

- 5 9. Process of Claim 8 wherein R<sup>13</sup> is -CO<sub>2</sub>R<sup>14</sup> and the product of Formula 3 is contacted with an alkali in an aqueous alcoholic solvent or with CF<sub>3</sub>CO<sub>2</sub>H at a temperature in the range of about 20 °C to the reflux temperature of the solvent for about 1-24 hours, followed by adjustment of the pH of the mixture to a value in the range of 3 to 7, to convert the product to the corresponding product wherein R<sup>13</sup> is -CO<sub>2</sub>H.
- 10 10. Process of Claim 9 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>13</sup> in Formula 1 is -CO<sub>2</sub>R<sup>14</sup> and is converted to -CO<sub>2</sub>H.
11. Process of Claim 9 wherein R<sup>14</sup> is t-butyl and the reaction is conducted in CF<sub>3</sub>CO<sub>2</sub>H.
- 15 12. Process of Claim 8 wherein R<sup>13</sup> is -CN and the product of Formula 3 is contacted with (i) a strong acid at reflux temperature of the solvent for about 2-96 hours or (ii) a strong alkali in an alcohol solvent at a temperature in the range of about 20 °C and the reflux temperature of the solvent for about 2-96 hours followed by adjustment of the pH to about 3-7, or (iii) sulfuric acid followed by acid or alkali, to convert the product to the corresponding compound wherein R<sup>13</sup> is -CO<sub>2</sub>H.
- 20 13. Process of Claim 12 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>13</sup> is -CO<sub>2</sub>R<sup>14</sup> and is converted to -CO<sub>2</sub>H.
14. Process of Claim 12 wherein R<sup>8</sup> is -(CH<sub>2</sub>)<sub>n</sub>CN and is converted to -(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>H, or is -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup> and is converted to (CH<sub>2</sub>)<sub>n</sub>OH when R<sup>13</sup> is converted to -CO<sub>2</sub>H.
- 25 15. Process of Claim 8 wherein R<sup>13</sup> is -CN and the product of Formula 3 is contacted with a mixture of equimolar amounts of sodium azide and ammonium chloride in a polar aprotic solvent at a temperature in the range of about 30 °C to the reflux temperature of the solvent, for about 1 hour to 10 days, to convert the product to the corresponding compound wherein R<sup>13</sup> is 5-tetrazolyl.
- 30 16. Process of Claim 15 wherein R<sup>8</sup> is -(CH<sub>2</sub>)<sub>m</sub>CN and is converted to -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl when R<sup>13</sup> is converted to 5-tetrazolyl.
- 35 17. Method of Claim 8 wherein R<sup>13</sup> is -CN and the product of Formula 3 is reacted with trialkyltin azide or triaryl tin azide followed by acidic or basic hydrolysis to convert the product to the corresponding compound wherein R<sup>13</sup> is 5-tetrazolyl.
- 40 18. Process of Claim 17 wherein R<sup>8</sup> is -(CH<sub>2</sub>)<sub>n</sub>CN and is converted to -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl when R<sup>13</sup> is converted to 5-tetrazolyl.
19. Process of Claim 8 wherein R<sup>13</sup> is -NO<sub>2</sub> and the product of Formula 3 is contacted with a reducing agent to form a second intermediate of Formula 3 in which R<sup>13</sup> is NH<sub>2</sub>, and the latter is contacted with an anhydride (CH<sub>3</sub>SO<sub>2</sub>)<sub>2</sub>O or (CF<sub>3</sub>SO<sub>2</sub>)<sub>2</sub>O or a chloride CH<sub>3</sub>SO<sub>2</sub>Cl or CF<sub>3</sub>SO<sub>2</sub>Cl of sulfonic acid in a solvent to produce a compound in which R<sup>13</sup> is -NHSO<sub>2</sub>CH<sub>3</sub> or -NHSO<sub>2</sub>CF<sub>3</sub>.
- 45 20. Process of Claim 19 wherein at least one of R<sup>2</sup>, R<sup>3</sup>, or R<sup>13</sup> is -NO<sub>2</sub> and is converted to -NHSO<sub>2</sub>CH<sub>3</sub> or -NHSO<sub>2</sub>CF<sub>3</sub>.
- 50 21. Process of Claim 9 or 12 wherein the compound of Formula 3 with R<sup>13</sup> = CO<sub>2</sub>H either
  - (a) is contacted with about 1-4 equivalents of thionyl chloride in excess thionyl chloride or another solvent at a temperature in the range of about 20 °C to the reflux temperature of the solvent for a period of about 5 minutes to about 2 hours to form an intermediate of Formula 3 wherein R<sup>13</sup> is COCl, and the latter is contacted with about 2-10 equivalents of hydroxylamine derivative H<sub>2</sub>NOR<sup>12</sup> in excess hydroxylamine derivative H<sub>2</sub>NOR<sup>12</sup> or other solvent, at a temperature in the range of about 25-80 °C for about 2-18 hours, or
  - (b) is contacted the hydroxylamine derivative H<sub>2</sub>NOR<sup>12</sup>, dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in a solvent at a temperature in the range of about 0-30 °C for about 1-24 hours;
- 55

to provide a compound in which  $R^{13}$  is  $\text{CONHOR}^{12}$ .

22. Process of Claim 6 wherein;  $R^1$  is



15

X is a carbon-carbon single bond, -CO-, -O-, -S-, or -NH-;  
 $R^2$ ,  $R^3$ ,  $R^6$  and  $R^7$  are as defined in Claim 1 and  
 $R^8$  is  $(\text{CH}_2)_n\text{OR}^{11}$ ,  $(\text{CH}_2)_n\text{OCOR}^{14}$ ,  
 $(\text{CH}_2)_n\text{CH}(\text{OH})\text{R}^{16}$ ,  $(\text{CH}_2)_n\text{COR}^{16}$   
 $(\text{CH}_2)_n\text{Cl}$ ,  $(\text{CH}_2)_n\text{CN}$ , CHO.

20 23. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OH}$  and the product of Formula 3 is contacted with an alcohol  $\text{R}^{11}\text{OH}$  in the anhydrous state in the presence of a strong acid or a Lewis acid, followed by saponification of any  $\text{CO}_2\text{R}^{14}$  groups concomitantly formed or present in intermediate 3, to form the corresponding compound of Formula 3 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OR}^{11}$  and  $\text{R}^{11}$  is not H.

25 24. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OR}^{11}$  and  $\text{R}^{11}$  is not H and the product of Formula 3 is contacted with an aqueous acidic medium at a temperature in the range of about  $25^\circ\text{C}$  and the reflux temperature of the solvent for a period of about 0.5-24 hours to form the corresponding compound of Formula 3 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OH}$ .

30 25. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OH}$  and the product of Formula 3 is contacted with  
 (a) a carboxylic acid anhydride  $(\text{R}^{14}\text{CO})_2\text{O}$  or chloride  $\text{R}^{14}\text{COCl}$  in a solvent in presence of a base at a temperature in the range of about  $0^\circ\text{C}$  and the reflux temperature of the solvent for about 0.5-24 hours or  
 (b) a carboxylic acid  $\text{R}^{14}\text{CO}_2\text{H}$  under anhydrous conditions in presence of a strong acid or Lewis  
 35 acid at about  $0^\circ$ - $100^\circ\text{C}$  for about 0.5 to 24 hours, to form the corresponding compound in which  $R^8$  is  $(\text{CH}_2)_n\text{OCOR}^{14}$ .

26. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OCOR}^{14}$  and the product of Formula 3 is contacted with aqueous acid or alkali to form the corresponding compound wherein  $R^8$  is  $(\text{CH}_2)_n\text{OH}$ .

40 27. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{OH}$  and the product of Formula 3 is contacted with an oxidizing agent at a temperature of about  $25$ - $45^\circ\text{C}$  for about 1-200 hours to produce a corresponding compound of Formula 3 in which  $R^8$  is  $(\text{CH}_2)_{n-1}\text{COR}^{16}$  and  $\text{R}^{16}$  is H.

45 28. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{COR}^{16}$  and  $\text{R}^{16}$  is H and the product of Formula 3 is contacted with an organometallic compound  $\text{R}^{15}\text{P}$  in which P is MgBr or Li in a solvent at a temperature in the range of about  $-78^\circ\text{C}$  to  $100^\circ\text{C}$  for about 0.5-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(\text{CH}_2)_n\text{CH}(\text{OH})\text{R}^{16}$  and  $\text{R}^{16}$  is not H.

50 29. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{CH}(\text{OH})\text{R}^{16}$  and  $\text{R}^{16}$  is not H and the product of Formula 3 is contacted with an oxidizing agent in a solvent to form a corresponding compound of Formula in which  $R^8$  is  $(\text{CH}_2)_n\text{COR}^{16}$  and  $\text{R}^{16}$  is not H.


55 30. Process of Claim 22 wherein  $R^8$  is  $(\text{CH}_2)_n\text{COR}^{16}$  and  $\text{R}^{16}$  is H and the compound of Formula 3 is contacted with an oxidizing agent in a solvent to form a corresponding compound of Formula 3 in which  $R^8$  is  $(\text{CH}_2)_n\text{COR}^{16}$  and  $\text{R}^{16}$  is OH.



31. Process of Claim 22 wherein  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is OH and the compound of Formula 3 is contacted with thionyl chloride in excess or in another solvent at a temperature in the range of about  $0^\circ C$  to the reflux temperature of the solvent for about 5 minutes to about 24 hours to form a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCOCl$  followed by contact of the latter with an amine  $NHR^{18}R^{19}$  in excess or in a solvent at temperatures in the range of about  $0^\circ C$  and reflux temperature of the solvent for about 5 minutes to about 24 hours to form a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCONR^{18}R^{19}$ .
32. Process of Claim 22 wherein  $R^8$  is  $(CH_2)_nOR^{11}$  and  $R^{11}$  is H and the product of Formula 3 is contacted with thionyl chloride in excess or in a solvent at a temperature in the range of about  $20^\circ C$  to the reflux temperature of the solvent for about 0.5-24 hours to form an intermediate compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCl$ .
33. Process of Claim 32 in which the compound of Formula 3 wherein  $R^8$  is  $(CH_2)_mCl$  is contacted with imidazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole or phthalimide in the presence of base in a solvent at temperatures in the range of about  $55^\circ C$  to the reflux temperature of the solvent for about 1-24 hours to produce a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_m$ -imidazole,  $(CH_2)_m$ -triazole,  $(CH_2)_m$ -tetrazole or  $(CH_2)_m$ -phthalimide.
34. Process of Claim 32 wherein the compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCl$  is contacted with sodium or potassium salt of a mercaptan  $R^{15}SH$  in a solvent at a temperature in the range of about  $25$ - $100^\circ C$  for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nSR^{15}$ .
35. Process of Claim 32 wherein the compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCl$  is contacted with an alkali metal cyanide in a solvent at a temperature in the range of about  $20$ - $100^\circ C$  for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCN$  and the latter compound is hydrolyzed to the corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is OH.
36. Process of Claim 32 wherein the compound of Formula 3 in which  $R^8$  is  $(CH_2)_{n-1}Cl$  is contacted with the sodium or potassium salt of a dialkyl malonate in a solvent at a temperature in the range of about  $20$ - $100^\circ C$  for about 0.5-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCH(CO_2alkyl)_2$  followed by saponification of the latter with aqueous alkali at a temperature in the range of about  $25^\circ C$  to the reflux temperature of the solvent followed by acidification with mineral acid to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCH(CO_2H)_2$  followed by heating the latter to about  $120^\circ C$  or in dilute mineral acid at reflux temperature to form a product of Formula 3 in which  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is OH.
37. Process of Claim 22 wherein  $R^8$  is  $(CH_2)_nCN$  and the compound of Formula 3 is contacted with sodium azide and ammonium chloride in a solvent at a temperature in the range of about  $30^\circ C$  and the reflux temperature of the solvent for about 1 hour to about 10 days to form a compound of the invention in which  $R^8$  is  $(CH_2)_n$ -tetrazole.
38. Process of Claim 22 wherein  $R^8$  is  $-CHO$  and the compound of Formula 3 is contacted with a methylene phosphorane  $(C_6H_5)_3P=CH(CH_2)_sCHR^{14}OR^{15}$  or  $(C_6H_5)_3P=CH(CH_2)_sCOR^{16}$  in a solvent at a temperature in the range of about  $25^\circ C$  to the reflux temperature of the solvent for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $-CH=CH(CH_2)_sCHR^{14}OR^{15}$  or  $-CH=CH(CH_2)_sCOR^{16}$ , except where  $R^{15}$  is H and  $R^{16}$  is OH, and optionally then contacting the compound of Formula 3 in which  $R^8$  is  $-CH=CH(CH_2)_sCOR^{16}$  with a reducing agent in a solvent at a temperature of about  $0^\circ$ - $25^\circ C$  for about 0.5-24 hours to form a product of Formula 3 in which  $R^8$  is  $-CH=CH(CH_2)_sCHR^{14}OH$ .
39. Process of Claim 22 wherein  $R^8$  is  $(CH_2)_mOH$  and the compound of Formula 3 is contacted with a fluorinating agent in a solvent at a temperature in the range of about  $-30^\circ C$  to  $25^\circ C$  for a period of about 0.5-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_mF$ .
40. Process of Claim 22 wherein the compound of Formula 3 in which  $R^8$  is  $(CH_2)_mCl$  is contacted with silver nitrate in a dipolar aprotic solvent at a temperature in the range of about  $25$ - $80^\circ C$  for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_mONO_2$ .





41. Process of Claim 22 wherein  $R^8$  is  $(CH_2)_nOH$  and the compound of Formula 3 is contacted with an isocyanate of Formula  $R^{10}NCO$  in a solvent at a temperature in the range of about  $25^\circ C$  to the reflux temperature of the solvent for a period of about 5 minutes to about 24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nOCONHR^{10}$ .
42. Process of Claim 22 wherein the compound in which  $R^8$  is  $(CH_2)_nCl$  is contacted with an amine  $R^{11}NH_2$  in excess amine or another solvent for a period of about 1-24 hours at a temperature in the range of about  $0^\circ C$  to the reflux temperature of the solvent to form an intermediate of Formula 3 in which  $R^8$  is  $(CH_2)_nNHR^{11}$ .
43. Process of Claim 22 in which  $R^8$  is  $(CH_2)_nCl$  and the compound of Formula 3 is contacted with an alkali metal azide in an aprotic solvent at a temperature in the range of about  $25-80^\circ C$  for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nN_3$  and the latter is contacted with a reducing agent to form an intermediate of Formula 3 in which  $R^8$  is  $(CH_2)_nNH_2$ .
44. Process of Claim 42 or 43 in which  $R^8$  is  $(CH_2)_nNHR^{11}$  or  $(CH_2)_nNH_2$  and the compound of Formula 3 is contacted with a chloroformate of Formula  $R^{10}OCOCI$  or a sulfonyl derivative of Formula  $R^{10}SO_2Cl$ , or  $(R^{10}SO_2)O$  in a solvent in the presence of a base at a temperature in the range of about  $0^\circ C$  to the reflux temperature of a solvent for about 5 minutes to about 24 hours to form a compound of Formula 3 in which  $R^8$  is  $-(CH_2)_nNR^{11}CO_2R^{10}$  or  $-(CH_2)_nNR^{11}SO_2R^{10}$ .
45. Process of Claim 42 or 43 in which the compound of Formula 3 with  $R^8$  equal to  $-(CH_2)_nNHR^{11}$  or  $(CH_2)_nNH_2$  is contacted with an isocyanate or isothiocyanate  $R^{10}NCY$  in a solvent at a temperature in the range of about  $25^\circ C$  to the reflux temperature of the solvent for about 5 minutes to about 24 hours to form a compound of the Formula 3 in which  $R^8$  is  $-(CH_2)_nNR^{11}CYNHR^{10}$ .
46. Process of Claim 6 wherein  $R^1$  is  $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$ , and  $R^8$  are as defined in Claim 1 in which the compound of Formula 3 wherein  $R^1$  is  $NO_2$  is reduced by means of iron and acetic acid, stannous chloride or hydrogen and palladium to a compound of Formula 3 wherein  $R^1$  is  $NH_2$  and the latter is reacted with an appropriate acid anhydride such as phthalic anhydride or a substituted phthalic anhydride in a solvent or with an appropriate acid chloride such as substituted anthranilic acid chloride in the presence of aqueous alkali or a base or with an appropriately substituted phthalic or anthranilic acid in the presence of dicyclohexylcarbodiimide in a solvent to produce a compound of the Formula 3 in which  $R^1$  is



and X is  $NHCO$ .

47. Process of Claim 6 wherein  $R^1$  is  $OCH_2C_6H_5$ ,  $R^2$  and  $R^3$  are H and  $R^6$ ,  $R^7$ , and  $R^8$  are as defined in Claim 1 and the resulting compound of Formula 3 with  $R^1$  equal to  $OCH_2C_6H_5$  is contacted with trifluoroacetic acid at reflux temperature for a period of about 0.2-1 hour or with hydrogen and palladium to form the corresponding compound of Formula 3 in which  $R^1$  is OH and the latter is contacted with a base at about  $25^\circ C$  and a suitable benzyl halide of the formula:




25 and X is -OCH<sub>2</sub>-.

48. Method of claim 6 wherein R<sup>8</sup> is -CHO, whereby the benzyl derivative of Formula 2 attaches to the imidazole derivative of Formula 1 preferentially at the nitrogen atom adjacent the carbon atom of the imidazole ring to which R<sup>8</sup> is attached.









# Claims for the following Contracting State : ES


1. A process for the preparation of an antihypertensive compound of the formula:



(I)


55 wherein  
R<sup>1</sup> is -4-CO<sub>2</sub>H; -4-CO<sub>2</sub>R<sup>9</sup>;






5


10



15



20




25

30

$\text{R}^2$  is H; Cl; Br; I; F;  $\text{NO}_2$ ; alkyl of 1 to 4 carbon atoms; acyloxy of 1 to 4 carbon atoms; alkoxy of 1 to 4 carbon atoms;  $\text{CO}_2\text{H}$ ;  $\text{CO}_2\text{R}^9$ ;  $\text{NHSO}_2\text{CH}_3$ ;  $\text{NHSO}_2\text{CF}_3$ ;  $\text{CONHOR}^{12}$ ;  $\text{SO}_2\text{NH}_2$  ;

35



aryl; or furyl;

$\text{R}^3$  is H; Cl, Br, I or F; alkyl of 1 to 4 carbon atoms or alkoxy of 1 to 4 carbon atoms;

$\text{R}^4$  is CN,  $\text{NO}_2$  or  $\text{CO}_2\text{R}^{11}$ ;


$\text{R}^5$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, alkenyl or alkynyl of 2 to 4 carbon atoms;

$\text{R}^6$  is alkyl of 2 to 10 carbon atoms, alkenyl or alkynyl of 3 to 10 carbon atoms or the same groups substituted with F or  $\text{CO}_2\text{R}^{14}$ ; cycloalkyl of 3 to 8 carbon atoms, cycloalkylalkyl, of 4 to 10 carbon atoms; cycloalkylalkenyl or cycloalkylalkynyl of 5 to 10 carbon atoms;  $(\text{CH}_2)_s\text{Z}(\text{CH}_2)_m\text{R}^5$  optionally substituted with F or  $\text{CO}_2\text{R}^{14}$ ; benzyl or benzyl substituted on the phenyl ring with 1 or 2 halogens, alkoxy of 1 to 4 carbon atoms, alkyl of 1 to 4 carbon atoms or nitro;






$\text{R}^7$  is H, F, Cl, Br, I,  $\text{NO}_2$ ,  $\text{CF}_3$  or CN;

$\text{R}^8$  is H, CN, alkyl of 1 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, or the same groups substituted with F; phenylalkenyl wherein the aliphatic portion is 2 to 6 carbon atoms;  $-(\text{CH}_2)_m\text{-imidazol-1-yl}$ ;  $-(\text{CH}_2)_m\text{-1,2,3-triazolyl}$  optionally substituted with one or two groups selected from  $\text{CO}_2\text{CH}_3$  or alkyl of 1 to 4 carbon atoms;  $-(\text{CH}_2)_m\text{-tetrazolyl}$ ;  $-(\text{CH}_2)_n\text{OR}^{11}$ ;


55




5




30



35




40



45

$R^9$  is

50









$R^{10}$  is alkyl of 1 to 6 carbon atoms or perfluoroalkyl of 1 to 6 carbon atoms, 1-adamantyl, 1-naphthyl, 1-(1-naphthyl)ethyl, or  $(CH_2)_p C_6 H_5$ ;

$R^{11}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;

55

$R^{12}$  is H, methyl or benzyl;

$R^{13}$  is  $-CO_2 H$ ;  $-CO_2 R^9$ ;  $-CH_2 CO_2 H$ ;  $-CH_2 CO_2 R^9$ ;




- $\text{R}^{14}$  is H, alkyl or perfluoroalkyl of 1 to 8 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;  
 $\text{R}^{15}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl, benzyl, acyl of 1 to 4 carbon atoms, phenacyl;  
 $\text{R}^{16}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms,  $(\text{CH}_2)_p\text{C}_6\text{H}_5$ ,  $\text{OR}^{17}$ , or  $\text{NR}^{18}\text{R}^{19}$ ;  
 $\text{R}^{17}$  is H, alkyl of 1 to 6 carbon atoms, cycloalkyl of 3 to 6 carbon atoms, phenyl or benzyl;  
 $\text{R}^{18}$  and  $\text{R}^{19}$  independently are H, alkyl of 1 to 4 carbon atoms, phenyl, benzyl,  $\alpha$ -methylbenzyl, or taken together form a ring of the formula




Q is NR<sup>20</sup>, O or CH<sub>2</sub>;  
 R<sup>20</sup> is H, alkyl of 1-4 carbon atoms, or phenyl;  
 R<sup>21</sup> is alkyl of 1 to 6 carbon atoms, -NR<sup>22</sup>R<sup>23</sup>, or

5




10 R<sup>22</sup> and R<sup>23</sup> independently are H, alkyl of 1 to 6 carbon atoms, benzyl, or are taken together as (CH<sub>2</sub>)<sub>u</sub> where u is 3-6;  
 R<sup>24</sup> is H, CH<sub>3</sub> or -C<sub>6</sub>H<sub>5</sub>;  
 R<sup>25</sup> is NR<sup>27</sup>R<sup>28</sup>, OR<sup>28</sup>, NHCONH<sub>2</sub>, NHCSNH<sub>2</sub>,

15



20 R<sup>26</sup> is hydrogen, alkyl with from 1 to 6 carbon atoms, benzyl, or allyl;  
 R<sup>27</sup> and R<sup>28</sup> are independently hydrogen, alkyl with from 1 to 5 carbon atoms, or phenyl;  
 R<sup>29</sup> and R<sup>30</sup> are independently alkyl of 1-4 carbon atoms or taken together are -(CH<sub>2</sub>)<sub>q</sub>;  
 R<sup>31</sup> is H, alkyl of 1 to 4 carbon atoms, -CH<sub>2</sub>CH=CH<sub>2</sub> or -CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>R<sup>32</sup>;  
 R<sup>32</sup> is H, NO<sub>2</sub>, NH<sub>2</sub>, OH or OCH<sub>3</sub>;  
 25 X is a carbon-carbon single bond, -CO-, -O-, -S-, -NH-,




30

-OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHC(R<sup>27</sup>)(R<sup>28</sup>), -NR<sup>23</sup>SO<sub>2</sub>-, -SO<sub>2</sub>NR<sup>23</sup>-, -C(R<sup>27</sup>)-  
 (R<sup>28</sup>)NH-, -CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, -CF<sub>2</sub>CF<sub>2</sub>-,

35



40



45

Y is O or S;  
 Z is O, NR<sup>11</sup>, or S;  
 m is 1 to 5;  
 n is 1 to 10;  
 p is 0 to 3;  
 50 q is 2 to 3;  
 r is 0 to 2;  
 s is 0 to 5;  
 t is 0 or 1;


and pharmaceutically acceptable salts of these compounds;

55


provided that:

(1) the R<sup>1</sup> group is not in the ortho position

(2) when  $R^1$  is




$X$  is a single bond, and  $R^{13}$  is  $CO_2H$ , or



then  $R^{13}$  must be in the ortho or meta position; or when  $R^1$  and  $X$  are as above and  $R^{13}$  is  $NHSO_2CF_3$  or  $NHSO_2CH_3$ ,  $R^{13}$  must be ortho;

(3) when  $R^1$  is




and  $X$  is other than a single bond, then  $R^{13}$  must be ortho except when  $X = NR^{23}CO$  and  $R^{13}$  is  $NHSO_2CF_3$  or  $NHSO_2CH_3$ , then  $R^{13}$  must be ortho or meta;

(4) when  $R^1$  is 4- $CO_2H$  or a salt thereof,  $R^6$  cannot be S-alkyl;

(5) when  $R^1$  is 4- $CO_2H$  or a salt thereof, the substituent on the 4-position of the imidazole cannot be  $CH_2OH$ ,  $CH_2OCOCH_3$ , or  $CH_2CO_2H$ ;


(6) when  $R^1$  is



$X$  is  $-OCH_2-$ ,

$R^{13}$  is 2- $CO_2H$ , and  $R^7$  is  $H$  then  $R^6$  is not  $C_2H_5S$ ;


(7) when  $R^1$  is



and  $R^6$  is n-hexyl then  $R^7$  and  $R^8$  are not both hydrogen;




(8) when R<sup>1</sup> is



R<sup>6</sup> is not methoxybenzyl.


(9) the R<sup>6</sup> group is not



or CH<sub>2</sub>OH,

which comprises contacting


an imidazole derivative of Formula 1 with a benzyl derivative of Formula 2 in a solvent in the presence of a base for about 1 to about 10 hours at a temperature in the range of about 20°C to the reflux temperature of the solvent to form a benzylimidazole of Formula 3:



wherein each of R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup>



is stable under the reaction conditions and is a group as defined above or an intermediate or protected form thereof which can be transformed to such a group and wherein X<sup>1</sup> is halogen, p-toluenesulfonyloxy or methylsulfonyloxy; and thereafter as necessary transforming said intermediate or protected forms of the R groups to R groups as defined above.

2. A process of claim 1 wherein the compounds prepared have the formula:







Wherein

R<sup>1</sup> is -CO<sub>2</sub>H; -NHSO<sub>2</sub>CF<sub>3</sub>;




$R^6$  is alkyl of 3 to 10 carbon atoms, alkenyl of 3 to 10 carbon atoms, alkynyl of 3 to 10 carbon atoms, cycloalkyl of 3 to 8 carbon atoms, benzyl substituted on the phenyl ring with up to two groups selected from alkoxy of 1 to 4 carbon atoms, halogen, alkyl of 1 to 4 carbon atoms, and nitro;


$R^8$  is phenylalkenyl wherein the aliphatic portion is 2 to 4 carbon atoms,  $-(CH_2)_m$ -imidazol-1-yl,  $-(CH_2)_m$ -1,2,3-triazolyl optionally substituted with one or two groups selected from  $CO_2CH_3$  or alkyl of 1 to 4 carbon atoms,  $(CH_2)_m$ -tetrazolyl,




$R^{13}$  is  $-CO_2H$ ,  $-CO_2R^9$ ,  $NHSO_2CF_3$ ; and



$R^{16}$  is H, alkyl of 1 to 5 carbon atoms,  $OR^{17}$ , or  $NR^{18}R^{19}$ ;  
 $X$  is carbon-carbon single bond,  $-CO-$ ,



-CH<sub>2</sub>CH<sub>2</sub>-,



-OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHCH<sub>2</sub>-, -CH<sub>2</sub>NH- or -CH=CH-; and pharmaceutically acceptable salts of these compounds.



3. A process of claim 2 wherein:

R<sup>2</sup> is H, alkyl of 1 to 4 carbon atoms, halogen, or alkoxy of 1 to 4 carbon atoms;


R<sup>6</sup> is alkyl, alkenyl or alkynyl of 3 to 7 carbon atoms;

R<sup>7</sup> is H, Cl, Br, I or CF<sub>3</sub>;

R<sup>8</sup> is -(CH<sub>2</sub>)<sub>m</sub>OR<sup>11</sup>;



-(CH<sub>2</sub>)<sub>m</sub>NHSO<sub>2</sub>R<sup>10</sup>;



or -COR<sup>16</sup>;

R<sup>10</sup> is CF<sub>3</sub>, alkyl of 1 to 6 carbon atoms or phenyl;

R<sup>11</sup> is H, or alkyl of 1 to 4 carbon atoms;

R<sup>13</sup> is CO<sub>2</sub>H; CO<sub>2</sub>CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>; NHSO<sub>2</sub>CF<sub>3</sub> and



R<sup>14</sup> is H, or alkyl of 1 to 4 carbon atoms;

R<sup>15</sup> is H, alkyl of 1 to 4 carbon atoms, or acyl of 1 to 4 carbon atoms;

R<sup>16</sup> is H, alkyl of 1 to 5 carbon atoms; OR<sup>17</sup>; or



m is 1 to 5;

X = single bond, -O-, -CO-, -NHCO-, or -OCH<sub>2</sub>-; and pharmaceutically acceptable salts.

4. A process of claims 1 to 3, wherein the compounds prepared are selected from 2-Butyl-4-chloro-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable

salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(methoxycarbonyl)aminomethyl]imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(propoxycarbonyl)aminomethyl]imidazole, or a pharmaceutically acceptable salt thereof;

2-Butyl-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-(1E-Butenyl)-4-chloro-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof;

2-propyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;


2-butyl-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof;

2-(1E-butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-hydroxymethyl)imidazole, or a pharmaceutically acceptable salt thereof; and

2-(1E-butenyl)-4-chloro-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxaldehyde, or a pharmaceutically acceptable salt thereof.

5. Process of claim 1 wherein compounds 1 and 2 are contacted in the presence of a base selected from the group consisting of a metal hydride, MH, a metal alkoxide, MOR, sodium carbonate, potassium carbonate, triethylamine and pyridine, in a dipolar aprotic solvent or, where the base is MOR, the solvent can be an alcohol, ROH, where M is lithium, sodium or potassium and R is methyl, ethyl or t-butyl.

6. Process of claim 1 wherein: R<sup>1</sup> is



X is a carbon-carbon single bond, -CO-, -O-, -S-, or -NH-;

R<sup>2</sup> and R<sup>3</sup> are each independently H, Cl, Br, I, CO<sub>2</sub>R<sup>14</sup>, F, NO<sub>2</sub>, alkyl of 1 to 4 carbon atoms, alkoxy of 1 to 4 carbon atoms, aryl or furyl;

R<sup>6</sup> and R<sup>7</sup> are as defined above;

R<sup>8</sup> is alkyl of 1 to 10 carbon atoms or alkenyl of 3 to 10 carbon atoms, or the same groups substituted with F; phenylalkenyl wherein the aliphatic portion is 2 to 6 carbon atoms; -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>; -(CH<sub>2</sub>)<sub>n</sub>SR<sup>15</sup>; or -(CH<sub>2</sub>)<sub>n</sub>CN;

R<sup>11</sup> is as defined above

R<sup>13</sup> is CO<sub>2</sub>R<sup>14</sup>, CN, NO<sub>2</sub>, trialkyltin tetrazole, or trityltetrazole; and


R<sup>14</sup> and R<sup>15</sup> are as defined above

7. Process of Claim 6 wherein R<sup>13</sup> is -CO<sub>2</sub>R<sup>14</sup> and the product of Formula 3 is contacted with an alkali in an aqueous alcoholic solvent or with CF<sub>3</sub>CO<sub>2</sub>H at a temperature in the range of about 20°C to the reflux temperature of the solvent for about 1-24 hours, followed by adjustment of the pH of the mixture

to a value in the range of 3 to 7, to convert the product to the corresponding product wherein  $R^{13}$  is  $-\text{CO}_2\text{H}$ .

- 5 8. Process of Claim 7 wherein at least one of  $R^2$ ,  $R^3$  or  $R^{13}$  in Formula 1 is  $-\text{CO}_2R^{14}$  and is converted to  $-\text{CO}_2\text{H}$ .
9. Process of Claim 8 wherein  $R^{14}$  is t-butyl and the reaction is conducted in  $\text{CF}_3\text{CO}_2\text{H}$ .
- 10 10. Process of Claim 7 wherein  $R^{13}$  is  $-\text{CN}$  and the product of Formula 3 is contacted with (i) a strong acid at reflux temperature of the solvent for about 2-96 hours or (ii) a strong alkali in an alcohol solvent at a temperature in the range of about  $20^\circ\text{C}$  and the reflux temperature of the solvent for about 2-96 hours followed by adjustment of the pH to about 3-7, or (iii) sulfuric acid followed by acid or alkali, to convert the product to the corresponding compound wherein  $R^{13}$  is  $-\text{CO}_2\text{H}$ .
- 15 11. Process of Claim 10 wherein at least one of  $R^2$ ,  $R^3$  or  $R^{13}$  is  $-\text{CO}_2R^{14}$  and is converted to  $-\text{CO}_2\text{H}$ .
12. Process of Claim 10 wherein  $R^8$  is  $-(\text{CH}_2)_n\text{CN}$  and is converted to  $-(\text{CH}_2)_n\text{CO}_2\text{H}$ , or is  $-(\text{CH}_2)_n\text{OR}^{11}$  and is converted to  $(\text{CH}_2)_n\text{OH}$  when  $R^{13}$  is converted to  $-\text{CO}_2\text{H}$ .
- 20 13. Process of Claim 6 wherein  $R^{13}$  is  $-\text{CN}$  and the product of Formula 3 is contacted with a mixture of equimolar amounts of sodium azide and ammonium chloride in a polar aprotic solvent at a temperature in the range of about  $30^\circ\text{C}$  to the reflux temperature of the solvent, for about 1 hour to 10 days, to convert the product to the corresponding compound wherein  $R^{13}$  is 5-tetrazolyl.
- 25 14. Process of Claim 13 wherein  $R^8$  is  $-(\text{CH}_2)_m\text{CN}$  and is converted to  $-(\text{CH}_2)_m$ -tetrazolyl when  $R^{13}$  is converted to 5-tetrazolyl.
15. Method of Claim 6 wherein  $R^{13}$  is  $-\text{CN}$  and the product of Formula 3 is reacted with trialkyltin azide or triaryltin azide followed by acidic or basic hydrolysis to convert the product to the corresponding compound wherein  $R^{13}$  is 5-tetrazolyl.
- 30 16. Process of Claim 15 wherein  $R^8$  is  $-(\text{CH}_2)_n\text{CN}$  and is converted to  $-(\text{CH}_2)_n$ -tetrazolyl when  $R^{13}$  is converted to 5-tetrazolyl.
- 35 17. Process of Claim 6 wherein  $R^{13}$  is  $-\text{NO}_2$  and the product of Formula 3 is contacted with a reducing agent to form a second intermediate of Formula 3 in which  $R^{13}$  is  $\text{NH}_2$ , and the latter is contacted with an anhydride  $(\text{CH}_3\text{SO}_2)_2\text{O}$  or  $(\text{CF}_3\text{SO}_2)_2\text{O}$  or a chloride  $\text{CH}_3\text{SO}_2\text{Cl}$  or  $\text{CF}_3\text{SO}_2\text{Cl}$  or sulfonic acid in a solvent to produce a compound in which  $R^{13}$  is  $-\text{NHSO}_2\text{CH}_3$  or  $-\text{NHSO}_2\text{CF}_3$ .
- 40 18. Process of Claim 17 wherein at least one of  $R^2$ ,  $R^3$ , or  $R^{13}$  is  $-\text{NO}_2$  and is converted to  $-\text{NHSO}_2\text{CH}_3$  or  $-\text{NHSO}_2\text{CF}_3$ .
19. Process of Claim 7 or 10 wherein the compound of Formula 3 with  $R^{13} = \text{CO}_2\text{H}$  either
  - 45 (a) is contacted with about 1-4 equivalents of thionyl chloride in excess thionyl chloride or another solvent at a temperature in the range of about  $20^\circ\text{C}$  to the reflux temperature of the solvent for a period of about 5 minutes to about 2 hours to form an intermediate of Formula 3 wherein  $R^{13}$  is  $\text{COCl}$ , and the latter is contacted with about 2-10 equivalents of hydroxylamine derivative  $\text{H}_2\text{NOR}^{12}$  in excess hydroxylamine derivative  $\text{H}_2\text{NOR}^{12}$  or other solvent, at a temperature in the range of about  $25$ - $80^\circ\text{C}$  for about 2-18 hours, or
  - 50 (b) is contacted the hydroxylamine derivative  $\text{H}_2\text{NOR}^{12}$ , dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in a solvent at a temperature in the range of about  $0$ - $30^\circ\text{C}$  for about 1-24 hours; to provide a compound in which  $R^{13}$  is  $\text{CONHOR}^{12}$ .

20. Process of Claim 1 wherein:  $R^1$  is



15

X is a carbon-carbon single bond, -CO-, -O-, -S-, or  
 $R^2$ ,  $R^3$ ,  $R^6$  and  $R^7$  are as defined in Claim 1 and  
 $R^8$  is  $(CH_2)_nOR^{11}$ ,  $(CH_2)_nOCOR^{14}$ ,  
 $(CH_2)_nCH(OH)R^{16}$ ,  $(CH_2)_nCOR^{16}$   
 $(CH_2)_nCl$ ,  $(CH_2)_nCN$ , CHO.

20

21. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nOH$  and the product of Formula 3 is contacted with an alcohol  $R^{11}OH$  in the anhydrous state in the presence of a strong acid or a Lewis acid, followed by saponification of any  $CO_2R^{14}$  groups concomitantly formed or present in intermediate 3, to form the corresponding compound of Formula 3 wherein  $R^8$  is  $(CH_2)_nOR^{11}$  and  $R^{11}$  is not H.

25

22. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nOR^{11}$  and  $R^{11}$  is not H and the product of Formula 3 is contacted with an aqueous acidic medium at a temperature in the range of about  $25^\circ C$  and the reflux temperature of the solvent for a period of about 0.5-24 hours to form the corresponding compound of Formula 3 wherein  $R^8$  is  $(CH_2)_nOH$ .

30

23. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nOH$  and the product of Formula 3 is contacted with  
 (a) a carboxylic acid anhydride  $(R^{14}CO)_2O$  or chloride  $R^{14}COCl$  in a solvent in presence of a base at a temperature in the range of about  $0^\circ C$  and the reflux temperature of the solvent for about 0.5-24 hours or  
 (b) a carboxylic acid  $R^{14}CO_2H$  under anhydrous conditions in presence of a strong acid or Lewis acid at about  $0^\circ - 100^\circ C$  for about 0.5 to 24 hours, to form the corresponding compound in which  $R^8$   
 35 is  $(CH_2)_nOCOR^{14}$ .

24. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nOCOR^{14}$  and the product of Formula 3 is contacted with aqueous acid or alkali to form the corresponding compound wherein  $R^8$  is  $(CH_2)_nOH$ .

40

25. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nOH$  and the product of Formula 3 is contacted with an oxidizing agent at a temperature of about  $25-45^\circ C$  for about 1-200 hours to produce a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_{n-1}COR^{16}$  and  $R^{16}$  is H.

45

26. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is H and the product of Formula 3 is contacted with an organometallic compound  $R^{16}P$  in which P is MgBr or Li in a solvent at a temperature in the range of about  $-78^\circ C$  to  $100^\circ C$  for about 0.5-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCH(OH)R^{16}$  and  $R^{16}$  is not H.

50

27. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nCH(OH)R^{16}$  and  $R^{16}$  is not H and the product of Formula 3 is contacted with an oxidizing agent in a solvent to form a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is not H.

55

28. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is H and the compound of Formula 3 is contacted with an oxidizing agent in a solvent to form a corresponding compound of Formula 3 in which  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is OH.


29. Process of Claim 20 wherein  $R^8$  is  $(CH_2)_nCOR^{16}$  and  $R^{16}$  is OH and the compound of Formula 3 is contacted with thionyl chloride in excess or in another solvent at a temperature in the range of about

- 0 °C to the reflux temperature of the solvent for about 5 minutes to about 24 hours to form a corresponding compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>COCl followed by contact of the latter with an amine NHR<sup>18</sup>R<sup>19</sup> in excess or in a solvent at temperatures in the range of about 0 °C and reflux temperature of the solvent for about 5 minutes to about 24 hours to form a corresponding compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>CONR<sup>18</sup>R<sup>19</sup>.
- 5
30. Process of Claim 20 wherein R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup> and R<sup>11</sup> is H and the product of Formula 3 is contacted with thionyl chloride in excess or in a solvent at a temperature in the range of about 20 °C to the reflux temperature of the solvent for about 0.5-24 hours to form an intermediate compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>Cl.
- 10
31. Process of Claim 30 in which the compound of Formula 3 wherein R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>Cl is contacted with imidazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole or phthalimide in the presence of base in a solvent at temperatures in the range of about 55 °C to the reflux temperature of the solvent for about 1-24 hours to produce a corresponding compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>-imidazole, (CH<sub>2</sub>)<sub>m</sub>-triazole, (CH<sub>2</sub>)<sub>m</sub>-tetrazole or (CH<sub>2</sub>)<sub>m</sub>-phthalimide.
- 15
32. Process of Claim 30 wherein the compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>Cl is contacted with sodium or potassium salt of a mercaptan R<sup>15</sup>SH in a solvent at a temperature in the range of about 25-100 °C for about 1-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>SR<sup>15</sup>.
- 20
33. Process of Claim 30 wherein the compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>Cl is contacted with an alkali metal cyanide in a solvent at a temperature in the range of about 20-100 °C for about 1-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>CN and the latter compound is hydrolyzed to the corresponding compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>COR<sup>16</sup> and R<sup>16</sup> is OH.
- 25
34. Process of Claim 30 wherein the compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n-1</sub>Cl is contacted with the sodium or potassium salt of a dialkyl malonate in a solvent at a temperature in the range of about 20-100 °C for about 0.5-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>CH(CO<sub>2</sub>alkyl)-<sub>2</sub> followed by saponification of the latter with aqueous alkali at a temperature in the range of about 25 °C to the reflux temperature of the solvent followed by acidification with mineral acid to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>CH(CO<sub>2</sub>H)<sub>2</sub> followed by heating the latter to about 120 °C or in dilute mineral acid at reflux temperature to form a product of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>COR<sup>16</sup> and R<sup>16</sup> is OH.
- 30
35. Process of Claim 20 wherein R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>CN and the compound of Formula 3 is contacted with sodium azide and ammonium chloride in a solvent at a temperature in the range of about 30 °C and the reflux temperature of the solvent for about 1 hour to about 10 days to form a compound of the invention in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>-tetrazole.
- 35
36. Process of Claim 20 wherein R<sup>8</sup> is -CHO and the compound of Formula 3 is contacted with a methylene phosphorane (C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>P=CH(CH<sub>2</sub>)<sub>s</sub>CHR<sup>14</sup>OR<sup>15</sup> or (C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>P=CH(CH<sub>2</sub>)<sub>s</sub>COR<sup>16</sup> in a solvent at a temperature in the range of about 25 °C to the reflux temperature of the solvent for about 1-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is -CH=CH(CH<sub>2</sub>)<sub>s</sub>CHR<sup>14</sup>OR<sup>15</sup> or -CH=CH(CH<sub>2</sub>)<sub>s</sub>COR<sup>16</sup>, except where R<sup>15</sup> is H and R<sup>16</sup> is OH, and optionally then contacting the compound of Formula 3 in which R<sup>8</sup> is -CH=CH(CH<sub>2</sub>)<sub>s</sub>COR<sup>16</sup> with a reducing agent in a solvent at a temperature of about 0 °-25 °C for about 0.5-24 hours to form a product of Formula 3 in which R<sup>8</sup> is -CH=CH(CH<sub>2</sub>)<sub>s</sub>CHR<sup>14</sup>OH.
- 40
37. Process of Claim 20 wherein R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>OH and the compound of Formula 3 is contacted with a fluorinating agent in a solvent at a temperature in the range of about -30 °C to 25 °C for a period of about 0.5-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>F.
- 45
38. Process of Claim 20 wherein the compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>Cl is contacted with silver nitrate in a dipolar aprotic solvent at a temperature in the range of about 25-80 °C for about 1-24 hours to form a compound of Formula 3 in which R<sup>8</sup> is (CH<sub>2</sub>)<sub>m</sub>ONO<sub>2</sub>.
- 50
39. Process of Claim 20 wherein R<sup>8</sup> is (CH<sub>2</sub>)<sub>n</sub>OH and the compound of Formula 3 is contacted with an isocyanate of Formula R<sup>10</sup>NCO in a solvent at a temperature in the range of about 25 °C to the reflux
- 55



temperature of the solvent for a period of about 5 minutes to about 24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nOCONHR^{10}$ .


40. Process of Claim 20 wherein the compound in which  $R^8$  is  $(CH_2)_nCl$  is contacted with an amine  $R^{11}NH_2$  in excess amine or another solvent for a period of about 1-24 hours at a temperature in the range of about  $0^\circ C$  to the reflux temperature of the solvent to form an intermediate of Formula 3 in which  $R^8$  is  $(CH_2)_nNHR^{11}$ .
41. Process of Claim 20 in which  $R^8$  is  $(CH_2)_nCl$  and the compound of Formula 3 is contacted with an alkali metal azide in an aprotic solvent at a temperature in the range of about  $25-80^\circ C$  for about 1-24 hours to form a compound of Formula 3 in which  $R^8$  is  $(CH_2)_nN_3$  and the latter is contacted with a reducing agent to form an intermediate of Formula 3 in which  $R^8$  is  $(CH_2)_nNH_2$ .
42. Process of Claim 40 or 41 in which  $R^8$  is  $(CH_2)_nNHR^{11}$  or  $(CH_2)_nNH_2$  and the compound of Formula 3 is contacted with a chloroformate of Formula  $R^{10}OCOCI$  or a sulfonyl derivative of Formula  $R^{10}SO_2Cl$ , or  $(R^{10}SO_2)O$  in a solvent in the presence of a base at a temperature in the range of about  $0^\circ C$  to the reflux temperature of a solvent for about 5 minutes to about 24 hours to form a compound of Formula 3 in which  $R^8$  is  $-(CH_2)_nNR^{11}CO_2R^{10}$  or  $-(CH_2)_nNR^{11}SO_2R^{10}$ .
43. Process of Claim 40 or 41 in which the compound of Formula 3 with  $R^8$  equal to  $-(CH_2)_nNHR^{11}$  or  $(CH_2)_nNH_2$  is contacted with an isocyanate or isothiocyanate  $R^{10}NCY$  in a solvent at a temperature in the range of about  $25^\circ C$  to the reflux temperature of the solvent for about 5 minutes to about 24 hours to form a compound of the Formula 3 in which  $R^8$  is  $-(CH_2)_nNR^{11}CYNHR^{10}$ .
44. Process of Claim 1 wherein  $R^1$  is  $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$ , and  $R^8$  are as defined in Claim 1 in which the compound of Formula 3 wherein  $R^1$  is  $NO_2$  is reduced by means of iron and acetic acid, stannous chloride or hydrogen and palladium to a compound of Formula 3 wherein  $R^1$  is  $NH_2$  and the latter is reacted with an appropriate acid anhydride such as phthalic anhydride or a substituted phthalic anhydride in a solvent or with an appropriate acid chloride such as substituted anthranilic acid chloride in the presence of aqueous alkali or a base or with an appropriately substituted phthalic or anthranilic acid in the presence of dicyclohexylcarbodiimide in a solvent to produce a compound of the Formula 3 in which  $R^1$  is




and X is  $NHCO$ .

45. Process of Claim 1 wherein  $R^1$  is  $OCH_2C_6H_5$ ,  $R^2$  and  $R^3$  are H and  $R^6$ ,  $R^7$ , and  $R^8$  are as defined in Claim 1 and the resulting compound of Formula 3 with  $R^1$  equal to  $OCH_2C_6H_5$  is contacted with trifluoroacetic acid at reflux temperature for a period of about 0.2-1 hour or with hydrogen and palladium to form the corresponding compound of Formula 3 in which  $R^1$  is OH and the latter is contacted with a base at about  $25^\circ C$  and a suitable benzyl halide of the formula:





15 to produce the corresponding compound of Formula 3 wherein R<sup>1</sup> is



25 and X is -OCH<sub>2</sub>-.

46. Method of claim 1 wherein R<sup>8</sup> is -CHO, whereby the benzyl derivative of Formula 2 attaches to the imidazole derivative of Formula 1 preferentially at the nitrogen atom adjacent the carbon atom of the imidazole ring to which R<sup>8</sup> is attached.

30

# Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. Blutdrucksenkende Verbindung der Formel


35



















50 worin  
R<sup>1</sup>

-4-CO<sub>2</sub>H, -4-CO<sub>2</sub>R<sup>9</sup>,

55







ist,  
 R<sup>2</sup> H, Cl, Br, I, F, NO<sub>2</sub>, Alkyl mit 1 bis 4 Kohlenstoffatomen, Acyloxy mit 1 bis 4 Kohlenstoffatomen, Alkoxy mit 1 bis 4 Kohlenstoffatomen, CO<sub>2</sub>H, CO<sub>2</sub>R<sup>9</sup>, NHSO<sub>2</sub>CH<sub>3</sub>, NHSO<sub>2</sub>CF<sub>3</sub>, CONHOR<sup>12</sup>, SO<sub>2</sub>NH<sub>2</sub>,




35

R<sup>3</sup> Aryl oder Furyl ist,  
 H, Cl, Br, I oder F, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Alkoxy mit 1 bis 4 Kohlenstoffatomen ist,  
 R<sup>4</sup> CN, NO<sub>2</sub> oder CO<sub>2</sub>R<sup>11</sup> ist,  
 R<sup>5</sup> H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Alkenyl oder Alkynyl mit 2 bis 4 Kohlenstoffatomen ist,  
 40 R<sup>6</sup> Alkyl mit 2 bis 10 Kohlenstoffatomen, Alkenyl oder Alkynyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F oder CO<sub>2</sub>R<sup>14</sup> substituierten Gruppen, Cycloalkyl mit 3 bis 8 Kohlenstoffatomen, Cycloalkylalkyl mit 4 bis 10 Kohlenstoffatomen, Cycloalkylalkenyl oder Cycloalkylalkynyl mit 5 bis 10 Kohlenstoffatomen, gegebenenfalls mit F oder CO<sub>2</sub>R<sup>14</sup> substituiertes (CH<sub>2</sub>)<sub>s</sub>Z(CH<sub>2</sub>)<sub>m</sub>R<sup>5</sup>, Benzyl oder am Phenylring mit 1 oder 2 Halogenen, Alkoxy mit 1 bis 4 Kohlenstoffatomen, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Nitro substituiertes Benzyl ist,  
 45 R<sup>7</sup> H, F, Cl, Br, I, NO<sub>2</sub>, CF<sub>3</sub> oder CN ist,  
 R<sup>8</sup> H, CN, Alkyl mit 1 bis 10 Kohlenstoffatomen, Alkenyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F substituierten Gruppen, Phenylalkenyl, in welchem der aliphatische Teil 2 bis 6 Kohlenstoffatome ist, -(CH<sub>2</sub>)<sub>m</sub>-imidazol-1-yl, gegebenenfalls mit einer oder zwei Gruppen, die aus CO<sub>2</sub>CH<sub>3</sub> oder Alkyl mit 1 bis 4 Kohlenstoffatomen ausgewählt sind, substituiertes -(CH<sub>2</sub>)<sub>m</sub>-1,2,3-triazolyl, -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl, -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>,



50







-PO<sub>3</sub>H, -C(CF<sub>3</sub>)<sub>2</sub>OH, -NHSO<sub>2</sub>CH<sub>3</sub>, -NHSO<sub>2</sub>CF<sub>3</sub>, -NHCOCF<sub>3</sub>, -CONHOR<sup>12</sup>, -SO<sub>2</sub>NH<sub>2</sub>,



-CONHNHSO<sub>2</sub>CF<sub>3</sub>,



ist,

R<sup>14</sup> H, Alkyl oder Perfluoralkyl mit 1 bis 8 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl oder Benzyl ist,

R<sup>15</sup> H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl, Benzyl, Acyl mit 1 bis 4 Kohlenstoffatomen, Phenacyl ist,

R<sup>16</sup> H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>, OR<sup>17</sup> oder NR<sup>18</sup>R<sup>19</sup> ist,

R<sup>17</sup> H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl oder Benzyl ist,

R<sup>18</sup> und R<sup>19</sup> unabhängig H, Alkyl mit 1 bis 4 Kohlenstoffatomen, Phenyl, Benzyl, α-Methylbenzyl sind oder zusammengekommen einen Ring der Formel




bilden,

Q NR<sup>20</sup>, O oder CH<sub>2</sub> ist,

R<sup>20</sup> H, Alkyl mit 1-4 Kohlenstoffatomen oder Phenyl ist,


R<sup>21</sup> Alkyl mit 1 bis 6 Kohlenstoffatomen, -NR<sup>22</sup>R<sup>23</sup> oder -CHCH<sub>2</sub>CO<sub>2</sub>-ist,



5

R<sup>22</sup> und R<sup>23</sup> unabhängig H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Benzyl oder zusammenge-  
 men (CH<sub>2</sub>)<sub>u</sub> sind, worin u 3-6 ist,  
 R<sup>24</sup> H, CH<sub>3</sub> oder -C<sub>6</sub>H<sub>5</sub> ist,  
 R<sup>25</sup> NR<sup>27</sup>R<sup>28</sup>, OR<sup>28</sup>, NHCONH<sub>2</sub>, NHCSNH<sub>2</sub>,


10



15

ist,  
 R<sup>26</sup> Wasserstoff, Alkyl mit 1 bis 6 Kohlenstoffatomen, Benzyl oder Allyl ist,  
 R<sup>27</sup> und R<sup>28</sup> unabhängig Wasserstoff, Alkyl mit 1 bis 5 Kohlenstoffatomen oder Phenyl sind,  
 R<sup>29</sup> und R<sup>30</sup> unabhängig Alkyl mit 1-4 Kohlenstoffatomen oder zusammenge-  
 sind,  
 R<sup>31</sup> H, Alkyl mit 1 bis 4 Kohlenstoffatomen, -CH<sub>2</sub>CH=CH<sub>2</sub> oder -CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>R<sup>32</sup> ist,  
 R<sup>32</sup> H, NO<sub>2</sub>, NH<sub>2</sub>, OH oder OCH<sub>3</sub> ist,  
 X eine Kohlenstoff-Kohlenstoff-Einfachbindung, -CO-, -O-, -S-, -NH-,


25




30

-OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHC(R<sup>27</sup>)(R<sup>28</sup>), -NR<sup>23</sup>SO<sub>2</sub>-, -SO<sub>2</sub>NR<sup>23</sup>-, -C(R<sup>27</sup>)-  
 (R<sup>28</sup>)NH-, -CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, -CF<sub>2</sub>CF<sub>2</sub>-,

35



40



45

ist,  
 Y O oder S ist,  
 Z O, NR<sup>11</sup> oder S ist,  
 m 1 bis 5 ist,  
 n 1 bis 10 ist,  
 p 0 bis 3 ist,  
 q 2 bis 3 ist,  
 r 0 bis 2 ist,  
 s 0 bis 5 ist,  
 t 0 oder 1 ist,

50


55

und pharmazeutisch annehmbare Salze dieser Verbindungen,

vorausgesetzt, daß

- (1) die Gruppe  $R^1$  nicht in ortho-Stellung ist,
- (2) wenn  $R^1$

5




10

15

ist,  $X$  eine Einfachbindung ist und  $R^{13}$   $CO_2H$  oder


20



25

ist,  $R^{13}$  sich in ortho- oder meta-Stellung befinden muß, oder wenn  $R^1$  und  $X$  wie vorstehend sind und  $R^{13}$   $NHSO_2CF_3$  oder  $NHSO_2CH_3$  ist,  $R^{13}$  ortho sein muß,

30




35

ist und  $X$  etwas anderes als eine Einfachbindung ist,  $R^{13}$  ortho sein muß, außer wenn  $X = NR^{23}CO$  und  $R^{13}$   $NHSO_2CF_3$  oder  $NHSO_2CH_3$  ist,  $R^{13}$  ortho oder meta sein muß,

40

- (4) wenn  $R^1$  4- $CO_2H$  oder ein Salz desselben ist,  $R^6$  nicht S-Alkyl sein kann,
- (5) wenn  $R^1$  4- $CO_2H$  oder ein Salz desselben ist, der Substituent in der 4-Stellung des Imidazols nicht  $CH_2OH$ ,  $CH_2OCOCH_3$  oder  $CH_2CO_2H$  sein kann,
- (6) wenn  $R^1$

45




50


55

ist,  $X$  - $OCH_2$ - ist,  $R^{13}$  2- $CO_2H$  ist und  $R^7$  H ist,  $R^6$  nicht  $C_2H_5S$  ist,


(7) wenn R<sup>1</sup>



ist und R<sup>6</sup> n-Hexyl ist, R<sup>7</sup> und R<sup>8</sup> nicht beide Wasserstoff sind,  
(8) wenn R<sup>1</sup>




ist, R<sup>6</sup> nicht Methoxybenzyl ist,  
(9) die Gruppe R<sup>6</sup> nicht



oder CH<sub>2</sub>OH ist.


2. Verbindung von Anspruch 1 mit der Formel



(II)

worin

R<sup>1</sup> -CO<sub>2</sub>H, -NHSO<sub>2</sub>CF<sub>3</sub>,





ist,


R<sup>6</sup> Alkyl mit 3 bis 10 Kohlenstoffatomen, Alkenyl mit 3 bis 10 Kohlenstoffatomen, Alkynyl mit 3 bis 10 Kohlenstoffatomen, Cycloalkyl mit 3 bis 8 Kohlenstoffatomen, am Phenylring mit bis zu zwei, aus Alkoxy mit 1 bis 4 Kohlenstoffatomen, Halogen, Alkyl mit 1 bis 4 Kohlenstoffato-






men und Nitro ausgewählten Gruppen substituiertes Benzyl ist,  
 $R^8$  Phenylalkenyl, worin der aliphatische Teil 2 bis 4 Kohlenstoffatome ist,  $-(CH_2)_m$ -imidazol-1-yl,  
 $-(CH_2)_m$ -1,2,3-triazol-1-yl, das gegebenenfalls mit einer oder zwei aus  $CO_2CH_3$  oder Alkyl mit  
 1 bis 4 Kohlenstoffatomen ausgewählten Gruppen substituiert ist,  $(CH_2)_m$ -tetrazolyl,  $-(CH_2)_nOR^{11}$ ,



ist,  
 $R^{13}$   $-CO_2H$ ,  $-CO_2R^9$ ,  $NHSO_2CF_3$  und



ist,  
 $R^{16}$  H, Alkyl mit 1 bis 5 Kohlenstoffatomen,  $OR^{17}$  oder  $NR^{18}R^{19}$  ist,  
 35 X eine Kohlenstoff-Kohlenstoff-Einfachbindung,  $-CO-$ ,




50  $-OCH_2-$ ,  $-CH_2O-$ ,  $-O-$ ,  $-SCH_2-$ ,  $-CH_2S-$ ,  $-NHCH_2-$ ,  $-CH_2NH-$  oder  $-CH=CH-$  ist, und pharmazeutisch annehmbare Salze dieser Verbindungen.

3. Verbindung von Anspruch 2, worin

$R^2$  H, Alkyl mit 1 bis 4 Kohlenstoffatomen, Halogen oder Alkoxy mit 1 bis 4 Kohlenstoffatomen ist,

55  $R^6$  Alkyl, Alkenyl oder Alkynyl mit 3 bis 7 Kohlenstoffatomen ist,


$R^7$  H, Cl, Br, I oder  $CF_3$  ist,



R<sup>8</sup>  $-(\text{CH}_2)_m \text{OR}^{11},$



$-(\text{CH}_2)_m \text{NHSO}_2 \text{R}^{10},$



oder  $-\text{COR}^{16}$  ist,

R<sup>10</sup>  $\text{CF}_3$ , Alkyl mit 1 bis 6 Kohlenstoffatomen oder Phenyl ist,

R<sup>11</sup> H oder Alkyl mit 1 bis 4 Kohlenstoffatomen ist,

R<sup>13</sup>  $\text{CO}_2\text{H}$ ,  $\text{CO}_2\text{CH}_2\text{OCO}(\text{CH}_3)_3$ ,  $\text{NHSO}_2\text{CF}_3$  und



ist,

R<sup>14</sup> H oder Alkyl mit 1 bis 4 Kohlenstoffatomen ist,

R<sup>15</sup> H, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Acyl mit 1 bis 4 Kohlenstoffatomen ist,

R<sup>16</sup> H, Alkyl mit 1 bis 5 Kohlenstoffatomen,  $\text{OR}^{17}$  oder



ist,

m 1 bis 5 ist,

X = Einfachbindung,  $-\text{O}-$ ,  $-\text{CO}-$ ,  $-\text{NHCO}-$  oder  $-\text{OCH}_2-$ , und pharmazeutisch annehmbare Salze.

4. Verbindungen der Ansprüche 1 bis 3, die aus 2-Butyl-4-chlor-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)-methyl]-5-(hydroxymethyl)imidazol oder einem pharmazeutisch annehmbaren Salz desselben, 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder einem pharmazeutisch annehmbaren Salz desselben, 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(methoxycarbonyl)aminomethyl]imidazol oder einem pharmazeutisch annehmbaren Salz desselben, 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(propoxycarbonyl)aminomethyl]imidazol oder einem pharmazeutisch annehmbaren Salz desselben, 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch annehmbaren Salz desselben, 2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch an-

nehmbaren Salz desselben,

2-(1E-Butenyl)-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder einem pharmazeutisch annehmbaren Salz desselben,

2-(1E-Butenyl)-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch annehmbaren Salz desselben,

2-Propyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder einem pharmazeutisch annehmbaren Salz desselben,

2-Propyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch annehmbaren Salz desselben,

2-Butyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch annehmbaren Salz desselben,


2-(1E-Butenyl)-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder einem pharmazeutisch annehmbaren Salz desselben und

2-(1E-Butenyl)-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxaldehyd oder einem pharmazeutisch annehmbaren Salz desselben ausgewählt sind.

5. Pharmazeutische Zusammensetzung umfassend einen pharmazeutisch geeigneten Träger und wenigstens eine Verbindung von Anspruch 1 bis 4.

6. Verfahren zur Herstellung einer Verbindung von Anspruch 1 bis 4, in welcher r 1 ist, welches das In-Berührung-Bringen eines Imidazolderivats der Formel 1 mit einem Benzylderivat der Formel 2 in einem Lösungsmittel in Anwesenheit einer Base während etwa 1 bis etwa 10 Stunden bei einer Temperatur im Bereich von etwa 20 °C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden eines Benzylimidazols der Formel 3

25



35

worin R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>6</sup>, R<sup>7</sup> und R<sup>8</sup> jeweils unter den Reaktionsbedingungen stabil ist und eine in Anspruch 1 definierte Gruppe oder ein Zwischenprodukt oder eine geschützte Form derselben ist, die in eine derartige Gruppe umgewandelt werden kann, und worin X<sup>1</sup> Halogen, p-Toluolsulfonyloxy oder Methylsulfonyloxy ist, und danach bei Bedarf das Umwandeln des Zwischenprodukts oder der geschützten Formen der Gruppen R in in Anspruch 1 definierte Gruppen R umfaßt.


40

7. Verfahren von Anspruch 6, bei welchem die Verbindungen 1 und 2 in Anwesenheit einer Base, welche aus der Gruppe ausgewählt ist, die aus einem Metallhydrid, MH, einem Metallalkoxid, MOR, Natriumcarbonat, Kaliumcarbonat, Triethylamin und Pyridin besteht, in einem dipolaren, aprotischen Lösungsmittel oder, wenn die Base MOR ist, das Lösungsmittel ein Alkohol, ROH, sein kann, worin M Lithium, Natrium oder Kalium ist und R Methyl, Ethyl oder t-Butyl ist, in Berührung gebracht werden.

50

55

8. Verfahren von Anspruch 6, bei welchem R<sup>1</sup>




ist,

- |    |                                     |  |
|----|-------------------------------------|--|
| 15 | X                                   | eine Kohlenstoff-Kohlenstoff-Einfachbindung, -CO-, -O-, -S- oder -NH- ist,   |
|    | R <sup>2</sup> und R <sup>3</sup>   | jeweils unabhängig H, Cl, Br, I, CO <sub>2</sub> R <sup>14</sup> , F, NO <sub>2</sub> , Alkyl mit 1 bis 4 Kohlenstoffatomen, Alkoxy mit 1 bis 4 Kohlenstoffatomen, Aryl oder Furyl sind, |
|    | R <sup>6</sup> und R <sup>7</sup>   | wie in Anspruch 1 definiert sind,  |
|    | R <sup>8</sup>                      | Alkyl mit 1 bis 10 Kohlenstoffatomen oder Alkenyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F substituierten Gruppen, Phenylalkenyl, worin der aliphatische                     |
| 20 |                                     | Teil 2 bis 6 Kohlenstoffatome ist, -(CH <sub>2</sub> ) <sub>n</sub> OR <sup>11</sup> , -(CH <sub>2</sub> ) <sub>n</sub> SR <sup>15</sup> oder -(CH <sub>2</sub> ) <sub>n</sub> CN ist,   |
|    | R <sup>11</sup>                     | wie in Anspruch 20 definiert ist,  |
|    | R <sup>13</sup>                     | CO <sub>2</sub> R <sup>14</sup> , CN, NO <sub>2</sub> , Trialkylzinntetrazol oder Trityltetrazol ist und   |
|    | R <sup>14</sup> und R <sup>15</sup> | wie in Anspruch 1 definiert sind.  |

- 25 9. Verfahren von Anspruch 8, bei welchem R<sup>13</sup> -CO<sub>2</sub>R<sup>14</sup> ist und das Produkt der Formel 3 mit einer Alkalie in einem wäßrigen alkoholischen Lösungsmittel oder mit CF<sub>3</sub>CO<sub>2</sub>H etwa 1-24 Stunden bei einer Temperatur im Bereich von etwa 20 °C bis zur Rückflußtemperatur des Lösungsmittels in Berührung gebracht wird, gefolgt von der Einstellung des pH des Gemisches auf einen Wert im Bereich von 3 bis 7 unter Überführen des Produkts in das entsprechende Produkt, worin R<sup>13</sup> -CO<sub>2</sub>H ist.
- 30 10. Verfahren von Anspruch 9, bei welchem wenigstens eines von R<sup>2</sup>, R<sup>3</sup> oder R<sup>13</sup> in Formel 1 -CO<sub>2</sub>R<sup>14</sup> ist und in -CO<sub>2</sub>H überführt wird.
- 35 11. Verfahren von Anspruch 9, bei welchem R<sup>14</sup> t-Butyl ist und die Reaktion in CF<sub>3</sub>CO<sub>2</sub>H ausgeführt wird.
- 40 12. Verfahren von Anspruch 8, bei welchem R<sup>13</sup> -CN ist und das Produkt der Formel 3 mit (i) einer starken Säure 2-96 Stunden bei Rückflußtemperatur des Lösungsmittels oder (ii) einer starken Alkalie in einem Alkohollösungsmittel bei einer Temperatur im Bereich von etwa 20 °C und der Rückflußtemperatur des Lösungsmittels etwa 2-96 Stunden, gefolgt von der Einstellung des pH auf etwa 3-7 oder (iii) Schwefelsäure gefolgt von Säure oder Alkali unter Überführen des Produkts in die entsprechende Verbindung, worin R<sup>13</sup> -CO<sub>2</sub>H ist, in Berührung gebracht wird.
- 45 13. Verfahren von Anspruch 12, bei welchem wenigstens eines von R<sup>2</sup>, R<sup>3</sup> oder R<sup>13</sup> -CO<sub>2</sub>R<sup>14</sup> ist und in -CO<sub>2</sub>H überführt wird.
- 50 14. Verfahren von Anspruch 12, bei welchem R<sup>8</sup> -(CH<sub>2</sub>)<sub>n</sub>CN ist und in -(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>H überführt wird oder -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup> ist und in (CH<sub>2</sub>)<sub>n</sub>OH überführt wird, wenn R<sup>13</sup> in -CO<sub>2</sub>H überführt wird.
- 55 15. Verfahren von Anspruch 8, bei welchem R<sup>13</sup> -CN ist und das Produkt der Formel 3 mit einem Gemisch äquimolarer Mengen Natriumazid und Ammoniumchlorid in einem polaren aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 30 °C bis zur Rückflußtemperatur des Lösungsmittels etwa 1 Stunde bis 10 Tage unter Überführen des Produkts in die entsprechende Verbindung, worin R<sup>13</sup> 5-Tetrazolyl ist, in Berührung gebracht wird.
16. Verfahren von Anspruch 15, bei welchem R<sup>8</sup> -(CH<sub>2</sub>)<sub>m</sub>CN ist und in -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl überführt wird, wenn R<sup>13</sup> in 5-Tetrazolyl überführt wird.

17. Verfahren von Anspruch 8, bei welchem  $R^{13}$  -CN ist und das Produkt der Formel 3 mit Trialkylzinnazid oder Triarylzinnazid umgesetzt wird, gefolgt von der sauren oder basischen Hydrolyse unter Überführen des Produkts in die entsprechende Verbindung, worin  $R^{13}$  5-Tetrazolyl ist.
18. Verfahren von Anspruch 17, bei welchem  $R^8$   $-(CH_2)_nCN$  ist und in  $-(CH_2)_m$ -tetrazolyl überführt wird, wenn  $R^{13}$  in 5-Tetrazolyl überführt wird.
19. Verfahren von Anspruch 8, bei welchem  $R^{13}$   $-NO_2$  ist und das Produkt der Formel 3 mit einem Reduktionsmittel unter Bilden eines zweiten Zwischenprodukts der Formel 3, in welcher  $R^{13}$   $NH_2$  ist, in Berührung gebracht wird und letzteres mit einem Anhydrid  $(CH_3SO_2)_2O$  oder  $(CF_3SO_2)_2O$  oder einem Chlorid  $CH_3SO_2Cl$  oder  $CF_3SO_2Cl$  einer Sulfonsäure in einem Lösungsmittel unter Herstellen einer Verbindung, in welcher  $R^{13}$   $-NHSO_2CH_3$  oder  $-NHSO_2CF_3$  ist, in Berührung gebracht wird.
20. Verfahren von Anspruch 19, bei welchem wenigstens eines von  $R^2$ ,  $R^3$  oder  $R^{13}$   $-NO_2$  ist und in  $-NHSO_2CH_3$  oder  $-NHSO_2CF_3$  überführt wird.
21. Verfahren von Anspruch 9 oder 12, bei welchem zum Liefern einer Verbindung, in der  $R^{13}$   $CONHOR^{12}$  ist, die Verbindung der Formel 3 mit  $R^{13} = CO_2H$  entweder
- (a) mit etwa 1-4 Äquivalenten Thionylchlorid in überschüssigem Thionylchlorid oder einem anderen Lösungsmittel bei einer Temperatur im Bereich von etwa  $20^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels über einen Zeitraum von etwa 5 Minuten bis etwa 2 Stunden unter Bilden eines Zwischenprodukts der Formel 3, worin  $R^{13}$   $COCl$  ist, in Berührung gebracht wird, und letzteres mit etwa 2-10 Äquivalenten des Hydroxylaminderivats  $H_2NOR^{12}$  in überschüssigem Hydroxylaminderivat  $H_2NOR^{12}$  oder einem anderen Lösungsmittel etwa 2-18 Stunden bei einer Temperatur im Bereich von etwa  $25-80^\circ C$  in Berührung gebracht wird, oder
- (b) mit dem Hydroxylaminderivat  $H_2NOR^{12}$ , Dicyclohexylcarbodiimid und 1-Hydroxybenzotriazol etwa 1-24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa  $0-30^\circ C$  in Berührung gebracht wird.
22. Verfahren von Anspruch 6, bei welchem  $R^1$



ist,

$X$  eine Kohlenstoff-Kohlenstoff-Einfachbindung,  $-CO-$ ,  $-O-$ ,  $-S-$  oder  $-NH-$  ist,  
 $R^2$ ,  $R^3$ ,  $R^6$  und  $R^7$  wie in Anspruch 1 definiert sind und  
 $R^8$   $(CH_2)_nOR^{11}$ ,  $(CH_2)_nOCOR^{14}$ ,  $(CH_2)_nCH(OH)R^{15}$ ,  $(CH_2)_nCOR^{15}$ ,  $(CH_2)_nCl$ ,  $(CH_2)_nCN$ ,  $CHO$  ist.

23. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOH$  ist und das Produkt der Formel 3 mit einem Alkohol  $R^{11}OH$  in wasserfreiem Zustand in Anwesenheit einer starken Säure oder einer Lewissäure in Berührung gebracht wird, gefolgt von der Verseifung irgendwelcher gleichzeitig gebildeter oder im Zwischenprodukt 3 vorhandener Gruppen  $CO_2R^{14}$  unter Bilden der entsprechenden Verbindung der Formel 3, worin  $R^8$   $(CH_2)_nOR^{11}$  ist und  $R^{11}$  nicht H ist.
24. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOR^{11}$  ist und  $R^{11}$  nicht H ist und das Produkt der Formel 3 mit einem wäßrigen sauren Medium bei einer Temperatur im Bereich von etwa  $25^\circ C$  und der Rückflußtemperatur des Lösungsmittels während eines Zeitraums von etwa 0,5-24 Stunden unter Bilden der entsprechenden Verbindung der Formel 3 in Berührung gebracht wird, worin  $R^8$   $(CH_2)_nOH$  ist.


25. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOH$  ist und das Produkt der Formel 3 mit  
 (a) einem Carbonsäureanhydrid  $(R^{14}CO)_2O$  oder -chlorid  $R^{14}COCl$  in einem Lösungsmittel in  
 Anwesenheit einer Base etwa 0,5-24 Stunden bei einer Temperatur im Bereich von etwa  $0^\circ C$  und  
 der Rückflußtemperatur des Lösungsmittels oder  
 5 (b) einer Carbonsäure  $R^{14}CO_2H$  unter wasserfreien Bedingungen in Anwesenheit einer starken Säure  
 oder Lewisäure etwa 0,5 bis 24 Stunden bei etwa  $0^\circ$ - $100^\circ C$  unter Bilden der entsprechenden  
 Verbindung, in welcher  $R^8$   $(CH_2)_nOCOR^{14}$  ist, in Berührung gebracht wird.
26. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOCOR^{14}$  ist und das Produkt der Formel 3 mit  
 10 wäßriger Säure oder Alkali unter Bilden der entsprechenden Verbindung, worin  $R^8$   $(CH_2)_nOH$  ist, in  
 Berührung gebracht wird.
27. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOH$  ist und das Produkt der Formel 3 mit einem  
 Oxidationsmittel etwa 1-200 Stunden bei einer Temperatur von etwa  $25-45^\circ C$  unter Herstellen einer  
 15 entsprechenden Verbindung der Formel 3, in welcher  $R^8$   $(CH_2)_{n-1}COR^{16}$  ist und  $R^{16}$  H ist, in Berührung  
 gebracht wird.
28. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nCOR^{16}$  ist und  $R^{16}$  H ist und das Produkt der  
 Formel 3 mit einer metallorganischen Verbindung  $R^{16}P$ , in welcher P MgBr oder Li ist, etwa 0,5-24  
 20 Stunden in einem Lösungsmittel bei einer Temperatur von etwa  $-78^\circ C$  bis  $100^\circ C$  unter Bilden einer  
 Verbindung der Formel 3 in Berührung gebracht wird, in der  $R^8$   $(CH_2)_nCH(OH)R^{16}$  ist und  $R^{16}$  nicht H  
 ist.
29. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nCH(OH)R^{16}$  ist und  $R^{16}$  nicht H ist und das Produkt  
 25 der Formel 3 mit einem Oxidationsmittel in einem Lösungsmittel unter Bilden einer entsprechenden  
 Verbindung der Formel 3 in Berührung gebracht wird, in der  $R^8$   $(CH_2)_nCOR^{16}$  ist und  $R^{16}$  nicht H ist.
30. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nCOR^{16}$  ist und  $R^{16}$  H ist und die Verbindung der  
 Formel 3 mit einem Oxidationsmittel in einem Lösungsmittel unter Bilden einer entsprechenden  
 30 Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8$   $(CH_2)_nCOR^{16}$  ist und  $R^{16}$  OH ist.
31. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nCOR^{16}$  ist und  $R^{16}$  OH ist und die Verbindung der  
 Formel 3 etwa 5 Minuten bis etwa 24 Stunden mit Thionylchlorid im Überschuß oder in einem anderen  
 Lösungsmittel bei einer Temperatur im Bereich von etwa  $0^\circ C$  bis zur Rückflußtemperatur des  
 35 Lösungsmittels unter Bilden einer entsprechenden Verbindung der Formel 3 in Berührung gebracht  
 wird, in welcher  $R^8$   $(CH_2)_nCOCl$  ist, gefolgt von der Berührung der letzteren mit einem Amin  $NHR^{18}R^{19}$   
 im Überschuß oder in einem Lösungsmittel bei Temperaturen im Bereich von etwa  $0^\circ C$  und der  
 Rückflußtemperatur des Lösungsmittels während etwa 5 Minuten bis 24 Stunden unter Bilden einer  
 entsprechenden Verbindung der Formel 3, in welcher  $R^8$   $(CH_2)_nCONR^{18}R^{19}$  ist.  
 40
32. Verfahren von Anspruch 22, bei welchem  $R^8$   $(CH_2)_nOR^{11}$  ist und  $R^{11}$  H ist und das Produkt der Formel  
 3 etwa 0,5-24 Stunden mit Thionylchlorid im Überschuß oder in einem Lösungsmittel bei einer  
 Temperatur im Bereich von etwa  $20^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels unter Bilden  
 einer Zwischenproduktverbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8$   $(CH_2)_nCl$  ist.  
 45
33. Verfahren von Anspruch 32, bei welchem die Verbindung der Formel 3, worin  $R^8$   $(CH_2)_mCl$  ist, mit  
 Imidazol, 1,2,3-Triazol, 1,2,4-Triazol, Tetrazol oder Phthalimid etwa 1-24 Stunden in Anwesenheit einer  
 Base in einem Lösungsmittel bei Temperaturen im Bereich von etwa  $55^\circ C$  bis zur Rückflußtemperatur  
 des Lösungsmittels unter Herstellen einer entsprechenden Verbindung der Formel 3 in Berührung  
 50 gebracht wird, in welcher  $R^8$   $(CH_2)_m$ -imidazol,  $(CH_2)_m$ -triazol,  $(CH_2)_m$ -tetrazol oder  $(CH_2)_m$ -phthalimid ist.
34. Verfahren von Anspruch 32, bei welchem die Verbindung der Formel 3, in welcher  $R^8$   $(CH_2)_nCl$  ist, mit  
 dem Natrium- oder Kaliumsalz eines Mercaptans  $R^{15}SH$  etwa 1-24 Stunden in einem Lösungsmittel bei  
 einer Temperatur im Bereich von etwa  $25-100^\circ C$  unter Bilden einer Verbindung der Formel 3 in  
 55 Berührung gebracht wird, in welcher  $R^8$   $(CH_2)_nSR^{15}$  ist.
35. Verfahren von Anspruch 32, bei welchem die Verbindung der Formel 3, in welcher  $R^8$   $(CH_2)_nCl$  ist, etwa  
 1-24 Stunden mit einem Alkalimetallcyanid in einem Lösungsmittel bei einer Temperatur im Bereich




von etwa 20-100 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_nCN$  ist, und die letztere Verbindung zu der entsprechenden Verbindung der Formel 3 hydrolysiert wird, in welcher  $R^8 (CH_2)_nCOR^{16}$  ist und  $R^{16} OH$  ist.

- 5 36. Verfahren von Anspruch 32, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_{n-1}Cl$  ist, etwa 0,5-24 Stunden mit dem Natrium- oder Kaliumsalz eines Malonsäuredialkylesters in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 20-100 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_nCH(CO_2Alkyl)_2$  ist, gefolgt von der Verseifung der letzteren mit wäßrigem Alkali bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels, gefolgt vom Ansäuern mit Mineralsäure unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nCH(C-O_2H)_2$  ist, gefolgt vom Erhitzen der letzteren auf etwa 120 °C oder in verdünnter Mineralsäure bei Rückflußtemperatur unter Bilden eines Produkts der Formel 3, in welcher  $R^8 (CH_2)_nCOR^{16}$  ist und  $R^{16} OH$  ist.
- 10 37. Verfahren von Beispiel 22, in welchem  $R^8 (CH_2)_nCN$  ist und die Verbindung der Formel 3 mit Natriumazid und Ammoniumchlorid etwa 1 Stunde bis etwa 10 Tage in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 30 °C und der Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Erfindung in Berührung gebracht wird, in welcher  $R^8 (CH_2)_n$ -tetrazol ist.
- 20 38. Verfahren von Anspruch 22, bei welchem  $R^8 -CHO$  ist und die Verbindung der Formel 3 mit einem Methylenphosphoran  $(C_6H_5)_3P=CH(CH_2)_sCHR^{14}OR^{15}$  oder  $(C_6H_5)_3P=CH(CH_2)_sCOR^{16}$  etwa 1-24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 -CH=CH(CH_2)_sCHR^{14}OR^{15}$  oder  $-CH=CH(CH_2)_sCOR^{16}$  ist, außer wo  $R^{15} H$  ist und  $R^{16} OH$  ist, und gegebenenfalls anschließend dem In-Berührung-Bringen der Verbindung der Formel 3, in welcher  $R^8 -CH=CH(CH_2)_sCOR^{16}$  ist, mit einem Reduktionsmittel in einem Lösungsmittel bei einer Temperatur von etwa 0 °-25 °C während etwa 0,5-24 Stunden unter Bilden eines Produkts der Formel 3, in welcher  $R^8 -CH=CH(CH_2)_sCHR^{14}OH$  ist.
- 25 39. Verfahren von Anspruch 22, bei welchem  $R^8 (CH_2)_mOH$  ist und die Verbindung der Formel 3 mit einem Fluorierungsmittel über einen Zeitraum von etwa 0,5-24 Stunden in einem Lösungsmittel bei einer Temperatur in dem Bereich von etwa -30 °C bis 25 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_mF$  ist.
- 30 40. Verfahren von Anspruch 22, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_mCl$  ist, etwa 1-24 Stunden mit Silbernitrat in einem dipolaren, aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 25-80 °C unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_mONO_2$  ist, in Berührung gebracht wird.
- 35 41. Verfahren von Anspruch 22, bei welchem  $R^8 (CH_2)_nOH$  ist und die Verbindung der Formel 3 mit einem Isocyanat der Formel  $R^{10}NCO$  in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels über einen Zeitraum von etwa 5 Minuten bis etwa 24 Stunden unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nOCONHR^{10}$  ist, in Berührung gebracht wird.
- 40 42. Verfahren von Anspruch 22, bei welchem die Verbindung, in welcher  $R^8 (CH_2)_nCl$  ist, mit einem Amin  $R^{11}NH_2$  in überschüssigem Amin oder in einem anderen Lösungsmittel über einen Zeitraum von etwa 1-24 Stunden bei einer Temperatur im Bereich von etwa 0 °C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden eines Zwischenprodukts der Formel 3, in welcher  $R^8 (CH_2)_nNHR^{11}$  ist, in Berührung gebracht wird.
- 45 43. Verfahren von Anspruch 22, bei welchem  $R^8 (CH_2)_nCl$  ist und die Verbindung der Formel 3 etwa 1-24 Stunden mit einem Alkalimetallazid in einem aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 25-80 °C unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nN_3$  ist, in Berührung gebracht wird und letztere mit einem Reduktionsmittel unter Bilden eines Zwischenproduktes der Formel 3, in welcher  $R^8 (CH_2)_nNH_2$  ist, in Berührung gebracht wird.
- 50 55


44. Verfahren von Anspruch 42 oder 43, bei welchem  $R^8$   $(CH_2)_nNHR^{11}$  oder  $(CH_2)_nNH_2$  ist und die Verbindung der Formel 3 mit einem Chlorameisensäureester der Formel  $R^{10}OCOCI$  oder einem Sulfonylderivat der Formel  $R^{10}SO_2Cl$  oder  $(R^{10}SO_2)O$  etwa 5 Minuten bis etwa 24 Stunden in einem Lösungsmittel in Anwesenheit einer Base bei einer Temperatur im Bereich von etwa  $0^\circ C$  bis zur Rückflußtemperatur eines Lösungsmittels unter Bilden einer Verbindung der Formel 3, in welcher  $R^8$  -  $(CH_2)_nNR^{11}CO_2R^{10}$  oder  $-(CH_2)_nNR^{11}SO_2R^{10}$  ist, in Berührung gebracht wird.
45. Verfahren von Anspruch 42 oder 43, bei welchem die Verbindung der Formel 3 mit  $R^8$  gleich  $-(CH_2)_nNHR^{11}$  oder  $(CH_2)_nNH_2$  mit einem Isocyanat oder Isothiocyanat  $R^{10}NCY$  etwa 5 Minuten bis etwa 24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa  $25^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Formel 3, in welcher  $R^8$   $-(CH_2)_nNR^{11}CYNHR^{10}$  ist, in Berührung gebracht wird.
46. Verfahren von Anspruch 6, bei welchem  $R^1$   $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$  und  $R^8$  wie in Anspruch 1 definiert sind, in welchem die Verbindung der Formel 3, worin  $R^1$   $NO_2$  ist, mittels Eisen und Essigsäure, Zinn(II)-chlorid oder Wasserstoff und Palladium zu einer Verbindung der Formel 3 reduziert wird, worin  $R^1$   $NH_2$  ist, und letztere mit einem geeigneten Säureanhydrid, wie Phthalsäureanhydrid oder ein substituiertes Phthalsäureanhydrid, in einem Lösungsmittel oder mit einem geeigneten Säurechlorid, wie etwa substituiertes Anthranilsäurechlorid, in Anwesenheit wäßrigen Alkalis oder einer Base oder mit einer geeignet substituierten Phthal- oder Anthranilsäure in Anwesenheit von Dicyclohexylcarbodiimid in einem Lösungsmittel unter Herstellen einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^1$



47. Verfahren von Anspruch 6, bei welchem  $R^1$   $OCH_2C_6H_5$  ist,  $R^2$  und  $R^3$  H sind und  $R^6$ ,  $R^7$  und  $R^8$  wie in Anspruch 1 definiert sind und die sich daraus ergebende Verbindung der Formel 3 mit  $R^1$  gleich  $OCH_2C_6H_5$  mit Trifluoressigsäure bei Rückflußtemperatur über einen Zeitraum von etwa 0,2-1 Stunde oder mit Wasserstoff und Palladium unter Bilden der entsprechenden Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^1$  OH ist, und letztere mit einer Base bei etwa  $25^\circ C$  und einem geeigneten Benzylhalogenid der Formel








ist, und X -OCH<sub>2</sub>- ist.

48. Verfahren von Anspruch 6, bei welchem R<sup>8</sup> -CHO ist, wobei das Benzylderivat der Formel 2 an das Imidazolderivat der Formel 1 vorzugsweise an das Stickstoffatom gebunden ist, welches dem Kohlenstoffatom des Imidazolrings, an welches R<sup>8</sup> gebunden ist, benachbart ist.


# Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung einer blutdrucksenkenden Verbindung der Formel




35 worin  
R<sup>1</sup>


-4-CO<sub>2</sub>H, -4-CO<sub>2</sub>R<sup>9</sup>,









45 -SO<sub>3</sub>H, -C(CF<sub>3</sub>)<sub>2</sub>OH,




55 -PO<sub>3</sub>H,






4-NHSO<sub>2</sub>CH<sub>3</sub>, -4-NHSO<sub>2</sub>CF<sub>3</sub>, -CONHOR<sup>12</sup>, -SO<sub>2</sub>NH<sub>2</sub>,




4-CONHNHSO<sub>2</sub>CF<sub>3</sub>,





ist,  
 $R^2$  H, Cl, Br, I, F,  $\text{NO}_2$ , Alkyl mit 1 bis 4 Kohlenstoffatomen, Acyloxy mit 1 bis 4 Kohlenstoffatomen, Alkoxy mit 1 bis 4 Kohlenstoffatomen,  $\text{CO}_2\text{H}$ ,  $\text{CO}_2\text{R}^9$ ,  $\text{NHSO}_2\text{CH}_3$ ,  $\text{NHSO}_2\text{CF}_3$ ,  $\text{CONHOR}^{12}$ ,  $\text{SO}_2\text{NH}_2$ ,



35

$R^3$  Aryl oder Furyl ist,  
H, Cl, Br, I oder F, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Alkoxy mit 1 bis 4 Kohlenstoffatomen ist,

$R^4$   $\text{CN}$ ,  $\text{NO}_2$  oder  $\text{CO}_2\text{R}^{11}$  ist,

$R^5$  H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Alkenyl oder Alkynyl mit 2 bis 4 Kohlenstoffatomen ist,

40


$R^6$  Alkyl mit 2 bis 10 Kohlenstoffatomen, Alkenyl oder Alkynyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F oder  $\text{CO}_2\text{R}^{14}$  substituierten Gruppen, Cycloalkyl mit 3 bis 8 Kohlenstoffatomen, Cycloalkylalkyl mit 4 bis 10 Kohlenstoffatomen, Cycloalkylalkenyl oder Cycloalkylalkynyl mit 5 bis 10 Kohlenstoffatomen, gegebenenfalls mit F oder  $\text{CO}_2\text{R}^{14}$  substituiertes  $(\text{CH}_2)_s\text{Z}(\text{CH}_2)_m\text{R}^5$ , Benzyl oder am Phenylring mit 1 oder 2 Halogenen, Alkoxy mit 1 bis 4 Kohlenstoffatomen, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Nitro substituiertes Benzyl ist,

$R^7$  H, F, Cl, Br, I,  $\text{NO}_2$ ,  $\text{CF}_3$  oder  $\text{CN}$  ist,


50

$R^8$  H,  $\text{CN}$ , Alkyl mit 1 bis 10 Kohlenstoffatomen, Alkenyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F substituierten Gruppen, Phenylalkenyl, in welchem der aliphatische Teil 2 bis 6 Kohlenstoffatome ist,  $-(\text{CH}_2)_m\text{-imidazol-1-yl}$ , gegebenenfalls mit einer oder zwei Gruppen, die aus  $\text{CO}_2\text{CH}_3$  oder Alkyl mit 1 bis 4 Kohlenstoffatomen ausgewählt sind, substituiertes  $-(\text{CH}_2)_m\text{-1,2,3-triazolyl}$ ,  $-(\text{CH}_2)_m\text{-tetrazolyl}$ ,  $-(\text{CH}_2)_n\text{OR}^{11}$ ,







5




10




15




20



25



30



35

R<sup>9</sup>

ist,

40



45

R<sup>10</sup>

ist,

Alkyl mit 1 bis 6 Kohlenstoffatomen oder Perfluoralkyl mit 1 bis 6 Kohlenstoffatomen, 1-Adamantyl, 1-Naphthyl, 1-(1-Naphthyl)ethyl oder (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub> ist,

R<sup>11</sup>

H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl oder Benzyl ist,


R<sup>12</sup>

H, Methyl oder Benzyl ist,

R<sup>13</sup>


-CO<sub>2</sub>H, -CO<sub>2</sub>R<sup>9</sup>, -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>R<sup>9</sup>,

50





55





$-\text{PO}_3\text{H}$ ,  $-\text{C}(\text{CF}_3)_2\text{OH}$ ,  $-\text{NHSO}_2\text{CH}_3$ ,  $-\text{NHSO}_2\text{CF}_3$ ,  $-\text{NHCOCF}_3$ ,  $-\text{CONHOR}^{12}$ ,  $-\text{SO}_2\text{NH}_2$ ,



$-\text{CONHNHSO}_2\text{CF}_3$ ,



ist,

$\text{R}^{14}$  H, Alkyl oder Perfluoralkyl mit 1 bis 8 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl oder Benzyl ist,

$\text{R}^{15}$  H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl, Benzyl, Acyl mit 1 bis 4 Kohlenstoffatomen, Phenacyl ist,

$\text{R}^{16}$  H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen,  $(\text{CH}_2)_p\text{C}_6\text{H}_5$ ,  $\text{OR}^{17}$  oder  $\text{NR}^{18}\text{R}^{19}$  ist,

$\text{R}^{17}$  H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Cycloalkyl mit 3 bis 6 Kohlenstoffatomen, Phenyl oder Benzyl ist,

$\text{R}^{18}$  und  $\text{R}^{19}$  unabhängig H, Alkyl mit 1 bis 4 Kohlenstoffatomen, Phenyl, Benzyl,  $\alpha$ -Methylbenzyl sind oder zusammengekommen einen Ring der Formel




bilden,

$\text{Q}$   $\text{NR}^{20}$ , O oder  $\text{CH}_2$  ist,

$\text{R}^{20}$  H, Alkyl mit 1-4 Kohlenstoffatomen oder Phenyl ist,


$\text{R}^{21}$  Alkyl mit 1 bis 6 Kohlenstoffatomen,  $-\text{NR}^{22}\text{R}^{23}$  oder -



5

ist,  
 $R^{22}$  und  $R^{23}$  unabhängig H, Alkyl mit 1 bis 6 Kohlenstoffatomen, Benzyl oder zusammenge-  
 nommen  $(\text{CH}_2)_u$  sind, worin u 3-6 ist,  
 $R^{24}$  H,  $\text{CH}_3$  oder  $-\text{C}_6\text{H}_5$  ist,  
 $R^{25}$   $\text{NR}^{27}\text{R}^{28}$ ,  $\text{OR}^{28}$ ,  $\text{NHCONH}_2$ ,  $\text{NHCSNH}_2$ ,

10




15

ist,  
 $R^{26}$  Wasserstoff, Alkyl mit 1 bis 6 Kohlenstoffatomen, Benzyl oder Allyl ist,  
 $R^{27}$  und  $R^{28}$  unabhängig Wasserstoff, Alkyl mit 1 bis 5 Kohlenstoffatomen oder Phenyl sind,  
 $R^{29}$  und  $R^{30}$  unabhängig Alkyl mit 1-4 Kohlenstoffatomen oder zusammenge-  
 nommen  $(\text{CH}_2)_q$  sind,  
 $R^{31}$  H, Alkyl mit 1 bis 4 Kohlenstoffatomen,  $-\text{CH}_2\text{CH}=\text{CH}_2$  oder  $-\text{CH}_2\text{C}_6\text{H}_4\text{R}^{32}$  ist,  
 $R^{32}$  H,  $\text{NO}_2$ ,  $\text{NH}_2$ , OH oder  $\text{OCH}_3$  ist,  
 25 X eine Kohlenstoff-Kohlenstoff-Einfachbindung,  $-\text{CO}-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{NH}-$ ,

20


25




30

$-\text{OCH}_2-$ ,  $-\text{CH}_2\text{O}-$ ,  $-\text{SCH}_2-$ ,  $-\text{CH}_2\text{S}-$ ,  $-\text{NHC}(\text{R}^{27})(\text{R}^{28})$ ,  $-\text{NR}^{23}\text{SO}_2-$ ,  $-\text{SO}_2\text{NR}^{23}-$ ,  $-\text{C}(\text{R}^{27})-$ ,  
 $(\text{R}^{28})\text{NH}-$ ,  $-\text{CH}=\text{CH}-$ ,  $-\text{CF}=\text{CF}-$ ,  $-\text{CH}=\text{CF}-$ ,  $-\text{CF}=\text{CH}-$ ,  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CF}_2\text{CF}_2-$ ,

35



40




45

ist,  
 Y O oder S ist,  
 50 Z O,  $\text{NR}^{11}$  oder S ist,  
 m 1 bis 5 ist,  
 n 1 bis 10 ist,  
 p 0 bis 3 ist,  
 q 2 bis 3 ist,  
 55 r 0 bis 2 ist,  
 s 0 bis 5 ist,  
 t 0 oder 1 ist,

und pharmazeutisch annehmbarer Salze dieser Verbindungen, vorausgesetzt, daß

- (1) die Gruppe  $R^1$  nicht in ortho-Stellung ist,  
 (2) wenn  $R^1$


5



10

ist,  $X$  eine Einfachbindung ist und  $R^{13}$   $CO_2H$  oder


15



20

ist,  $R^{13}$  sich in ortho- oder meta-Stellung befinden muß, oder wenn  $R^1$  und  $X$  wie vorstehend sind  
 und  $R^{13}$   $NHSO_2CF_3$  oder  $NHSO_2CH_3$  ist,  $R^{13}$  ortho sein muß,  
 (3) wenn  $R^1$

25




30

35

ist und  $X$  etwas anderes als eine Einfachbindung ist,  $R^{13}$  ortho sein muß, außer wenn  $X = NR^{23}CO$   
 und  $R^{13}$   $NHSO_2CF_3$  oder  $NHSO_2CH_3$  ist,  $R^{13}$  ortho oder meta sein muß,  
 (4) wenn  $R^1$  4- $CO_2H$  oder ein Salz desselben ist,  $R^6$  nicht S-Alkyl sein kann,  
 (5) wenn  $R^1$  4- $CO_2H$  oder ein Salz desselben ist, der Substituent in der 4-Stellung des Imidazols  
 nicht  $CH_2OH$ ,  $CH_2OCOCH_3$  oder  $CH_2CO_2H$  sein kann,  
 (6) wenn  $R^1$

40

45




50


ist,  $X$  - $OCH_2$ - ist,  $R^{13}$  2- $CO_2H$  ist und  $R^7$  H ist,  $R^6$  nicht  $C_2H_5S$  ist,

55


(7) wenn R<sup>1</sup>



ist und R<sup>6</sup> n-Hexyl ist, R<sup>7</sup> und R<sup>8</sup> nicht beide Wasserstoff sind,  
(8) wenn R<sup>1</sup>




ist, R<sup>6</sup> nicht Methoxybenzyl ist,  
(9) die Gruppe R<sup>6</sup> nicht



oder CH<sub>2</sub>OH ist,


welches das In-Berührung-Bringen eines Imidazolderivats der Formel 1 mit einem Benzylderivat der Formel 2 in einem Lösungsmittel in Anwesenheit einer Base während etwa 1 bis etwa 10 Stunden bei einer Temperatur im Bereich von etwa 20°C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden eines Benzylimidazols der Formel 3



worin R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>6</sup>, R<sup>7</sup> und R<sup>8</sup> jeweils unter den Reaktionsbedingungen stabil ist und eine vorstehend definierte Gruppe oder ein Zwischenprodukt oder eine geschützte Form derselben ist, die in eine derartige Gruppe umgewandelt werden kann, und worin X<sup>1</sup> Halogen, p-Toluolsulfonyloxy oder Methylsulfonyloxy ist, und danach bei Bedarf das Umwandeln des Zwischenprodukts oder der geschützten Formen der Gruppen R in vorstehend definierte Gruppen R umfaßt.




2. Verfahren von Anspruch 1, bei welchem die hergestellten Verbindungen die Formel



besitzen, worin

R<sup>1</sup> -CO<sub>2</sub>H, -NHSO<sub>2</sub>CF<sub>3</sub>,





30


ist,

R<sup>6</sup> Alkyl mit 3 bis 10 Kohlenstoffatomen, Alkenyl mit 3 bis 10 Kohlenstoffatomen, Alkynyl mit 3 bis 10 Kohlenstoffatomen, Cycloalkyl mit 3 bis 8 Kohlenstoffatomen, am Phenylring mit bis zu zwei, aus Alkoxy mit 1 bis 4 Kohlenstoffatomen, Halogen, Alkyl mit 1 bis 4 Kohlenstoffatomen und Nitro ausgewählten Gruppen substituiertes Benzyl ist,

35

R<sup>8</sup> Phenylalkenyl, worin der aliphatische Teil 2 bis 4 Kohlenstoffatome ist, -(CH<sub>2</sub>)<sub>m</sub>-imidazol-1-yl, -(CH<sub>2</sub>)<sub>m</sub>-1,2,3-triazol-1-yl, das gegebenenfalls mit einer oder zwei aus CO<sub>2</sub>CH<sub>3</sub> oder Alkyl mit 1 bis 4 Kohlenstoffatomen ausgewählten Gruppen substituiert ist, (CH<sub>2</sub>)<sub>m</sub>-tetrazolyl, -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>,






5

10


ist,  
 $R^{16}$  H, Alkyl mit 1 bis 5 Kohlenstoffatomen,  $OR^{17}$  oder  $NR^{18}R^{19}$  ist,  
 $X$  eine Kohlenstoff-Kohlenstoff-Einfachbindung,  $-CO-$ ,



15

$-CH_2CH_2-$ ,

20



25


$-OCH_2-$ ,  $-CH_2O-$ ,  $-O-$ ,  $-SCH_2-$ ,  $-CH_2S-$ ,  $-NHCH_2-$ ,  $-CH_2NH-$  oder  $-CH=CH-$  ist, und pharmazeutisch annehmbare Salze dieser Verbindungen.

3. Verfahren von Anspruch 2, bei welchem

30

$R^2$  H, Alkyl mit 1 bis 4 Kohlenstoffatomen, Halogen oder Alkoxy mit 1 bis 4 Kohlenstoffatomen ist,  
 $R^6$  Alkyl, Alkenyl oder Alkynyl mit 3 bis 7 Kohlenstoffatomen ist,  
 $R^7$  H, Cl, Br, I oder  $CF_3$  ist,  
 $R^8$   $-(CH_2)_mOR^{11}$ ,

35




40



45

$-(CH_2)_mNHSO_2R^{10}$ ,

50



55

oder  $-COR^{16}$  ist,  
 $R^{10}$   $CF_3$ , Alkyl mit 1 bis 6 Kohlenstoffatomen oder Phenyl ist,  
 $R^{11}$  H oder Alkyl mit 1 bis 4 Kohlenstoffatomen ist,  
 $R^{13}$   $CO_2H$ ,  $CO_2CH_2OCO(CH_3)_3$ ,  $NHSO_2CF_3$  und



5

ist,

- R<sup>14</sup> H oder Alkyl mit 1 bis 4 Kohlenstoffatomen ist,  
 R<sup>15</sup> H, Alkyl mit 1 bis 4 Kohlenstoffatomen oder Acyl mit 1 bis 4 Kohlenstoffatomen ist,  
 R<sup>16</sup> H, Alkyl mit 1 bis 5 Kohlenstoffatomen, OR<sup>17</sup> oder

10



15

ist,

- m 1 bis 5 ist,  
 X = Einfachbindung, -O-, -CO-, -NHCO- oder -OCH<sub>2</sub>-, und pharmazeutisch annehmbare Salze.


20

4. Verfahren von Anspruch 1 bis 3, bei welchem die hergestellten Verbindungen
- 2-Butyl-4-chlor-1-[(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(methoxycarbonyl)aminomethyl]imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-[(propoxycarbonyl)aminomethyl]imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben,
- 2-(1E-Butenyl)-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-(1E-Butenyl)-4-chlor-1-[(2'-carboxybiphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Propyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Propyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben,
- 2-Butyl-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben,
- 2-(1E-Butenyl)-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]-5-(hydroxymethyl)imidazol oder ein pharmazeutisch annehmbares Salz desselben und
- 2-(1E-Butenyl)-4-chlor-1-[2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazol-5-carboxyaldehyd oder ein pharmazeutisch annehmbares Salz desselben sind.

50

5. Verfahren von Anspruch 1, bei welchem die Verbindungen 1 und 2 in Anwesenheit einer Base, welche aus der Gruppe ausgewählt ist, die aus einem Metallhydrid, MH, einem Metallalkoxid, MOR, Natriumcarbonat, Kaliumcarbonat, Triethylamin und Pyridin besteht, in einem dipolaren, aprotischen Lösungsmittel oder, wenn die Base MOR ist, das Lösungsmittel ein Alkohol, ROH, sein kann, worin M Lithium, Natrium oder Kalium ist und R Methyl, Ethyl oder t-Butyl ist, in Berührung gebracht werden.

55

6. Verfahren von Anspruch 1, bei welchem R<sup>1</sup>

ist,

- 15
- |                                     |   |
|-------------------------------------|---|
| X                                   | eine Kohlenstoff-Kohlenstoff-Einfachbindung, -CO-, -O-, -S- oder -NH- ist,  |
| R <sup>2</sup> und R <sup>3</sup>   | jeweils unabhängig H, Cl, Br, I, CO <sub>2</sub> R <sup>14</sup> , F, NO <sub>2</sub> , Alkyl mit 1 bis 4 Kohlenstoffatomen, Alkoxy mit 1 bis 4 Kohlenstoffatomen, Aryl oder Furyl sind,  |
| R <sup>6</sup> und R <sup>7</sup>   | wie vorstehend definiert sind,  |
| R <sup>8</sup>                      | Alkyl mit 1 bis 10 Kohlenstoffatomen oder Alkenyl mit 3 bis 10 Kohlenstoffatomen oder dieselben, mit F substituierten Gruppen, Phenylalkenyl, worin der aliphatische Teil 2 bis 6 Kohlenstoffatome ist, -(CH <sub>2</sub> ) <sub>n</sub> OR <sup>11</sup> , -(CH <sub>2</sub> ) <sub>n</sub> SR <sup>15</sup> oder -(CH <sub>2</sub> ) <sub>n</sub> CN ist, |
| R <sup>11</sup>                     | wie vorstehend definiert ist,   |
| R <sup>13</sup>                     | CO <sub>2</sub> R <sup>14</sup> , CN, NO <sub>2</sub> , Trialkylzinn-tetrazol oder Trityltetrazol ist und   |
| R <sup>14</sup> und R <sup>15</sup> | wie vorstehend definiert sind.  |
- 20

- 25
7. Verfahren von Anspruch 6, bei welchem R<sup>13</sup> -CO<sub>2</sub>R<sup>14</sup> ist und das Produkt der Formel 3 mit einer Alkalie in einem wäßrigen alkoholischen Lösungsmittel oder mit CF<sub>3</sub>CO<sub>2</sub>H etwa 1-24 Stunden bei einer Temperatur im Bereich von etwa 20 °C bis zur Rückflußtemperatur des Lösungsmittels in Berührung gebracht wird, gefolgt von der Einstellung des pH des Gemisches auf einen Wert im Bereich von 3 bis 7 unter Überführen des Produkts in das entsprechende Produkt, worin R<sup>13</sup> -CO<sub>2</sub>H ist.
- 30

8. Verfahren von Anspruch 7, bei welchem wenigstens eines von R<sup>2</sup>, R<sup>3</sup> oder R<sup>13</sup> in Formel 1 -CO<sub>2</sub>R<sup>14</sup> ist und in -CO<sub>2</sub>H überführt wird.

9. Verfahren von Anspruch 8, bei welchem R<sup>14</sup> t-Butyl ist und die Reaktion in CF<sub>3</sub>CO<sub>2</sub>H ausgeführt wird.
- 35

10. Verfahren von Anspruch 7, bei welchem R<sup>13</sup> -CN ist und das Produkt der Formel 3 mit (i) einer starken Säure 2-96 Stunden bei Rückflußtemperatur des Lösungsmittels oder (ii) einer starken Alkalie in einem Alkohollösungsmittel bei einer Temperatur im Bereich von etwa 20 °C und der Rückflußtemperatur des Lösungsmittels etwa 2-96 Stunden, gefolgt von der Einstellung des pH auf etwa 3-7 oder (iii) Schwefelsäure gefolgt von Säure oder Alkali unter Überführen des Produkts in die entsprechende Verbindung, worin R<sup>13</sup> -CO<sub>2</sub>H ist, in Berührung gebracht wird.
- 40


11. Verfahren von Anspruch 10, bei welchem wenigstens eines von R<sup>2</sup>, R<sup>3</sup> oder R<sup>13</sup> -CO<sub>2</sub>R<sup>14</sup> ist und in -CO<sub>2</sub>H überführt wird.
- 45

12. Verfahren von Anspruch 10, bei welchem R<sup>8</sup> -(CH<sub>2</sub>)<sub>n</sub>CN ist und in -(CH<sub>2</sub>)<sub>n</sub>CO<sub>2</sub>H überführt wird oder -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup> ist und in (CH<sub>2</sub>)<sub>n</sub>OH überführt wird, wenn R<sup>13</sup> in -CO<sub>2</sub>H überführt wird.

13. Verfahren von Anspruch 6, bei welchem R<sup>13</sup> -CN ist und das Produkt der Formel 3 mit einem Gemisch äquimolarer Mengen Natriumazid und Ammoniumchlorid in einem polaren aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 30 °C bis zur Rückflußtemperatur des Lösungsmittels etwa 1 Stunde bis 10 Tage unter Überführen des Produkts in die entsprechende Verbindung, worin R<sup>13</sup> 5-Tetrazolyl ist, in Berührung gebracht wird.
- 50

14. Verfahren von Anspruch 13, bei welchem R<sup>8</sup> -(CH<sub>2</sub>)<sub>n</sub>CN ist und in -(CH<sub>2</sub>)<sub>m</sub>-tetrazolyl überführt wird, wenn R<sup>13</sup> in 5-Tetrazolyl überführt wird.
- 55

15. Verfahren von Anspruch 6, bei welchem  $R^{13}$  -CN ist und das Produkt der Formel 3 mit Trialkylzinnazid oder Triarylzinnazid umgesetzt wird, gefolgt von der sauren oder basischen Hydrolyse unter Überführen des Produkts in die entsprechende Verbindung, worin  $R^{13}$  5-Tetrazolyl ist.
- 5 16. Verfahren von Anspruch 15, bei welchem  $R^8$   $-(CH_2)_nCN$  ist und in  $-(CH_2)_m$ -tetrazolyl überführt wird, wenn  $R^{13}$  in 5-Tetrazolyl überführt wird.
17. Verfahren von Anspruch 6, bei welchem  $R^{13}$   $-NO_2$  ist und das Produkt der Formel 3 mit einem Reduktionsmittel unter Bilden eines zweiten Zwischenprodukts der Formel 3, in welcher  $R^{13}$   $NH_2$  ist, in  
10 Berührung gebracht wird und letzteres mit einem Anhydrid  $(CH_3SO_2)_2O$  oder  $(CF_3SO_2)_2O$  oder einem Chlorid  $CH_3SO_2Cl$  oder  $CF_3SO_2Cl$  einer Sulfonsäure in einem Lösungsmittel unter Herstellen einer Verbindung, in welcher  $R^{13}$   $-NHSO_2CH_3$  oder  $-NHSO_2CF_3$  ist, in Berührung gebracht wird.
18. Verfahren von Anspruch 17, bei welchem wenigstens eines von  $R^2$ ,  $R^3$  oder  $R^{13}$   $-NO_2$  ist und in  
15  $-NHSO_2CH_3$  oder  $-NHSO_2CF_3$  überführt wird.
19. Verfahren von Anspruch 7 oder 10, bei welchem zum Liefern einer Verbindung, in der  $R^{13}$   $CONHOR^{12}$  ist, die Verbindung der Formel 3 mit  $R^{13} = CO_2H$  entweder  
20 (a) mit etwa 1-4 Äquivalenten Thionylchlorid in überschüssigem Thionylchlorid oder einem anderen Lösungsmittel bei einer Temperatur im Bereich von etwa  $20^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels über einen Zeitraum von etwa 5 Minuten bis etwa 2 Stunden unter Bilden eines Zwischenprodukts der Formel 3, worin  $R^{13}$   $COCl$  ist, in Berührung gebracht wird, und letzteres mit etwa 2-10 Äquivalenten des Hydroxylaminderivats  $H_2NOR^{12}$  in überschüssigem Hydroxylaminderivat  $H_2NOR^{12}$  oder einem anderen Lösungsmittel etwa 2-18 Stunden bei einer Temperatur im Bereich  
25 von etwa  $25-80^\circ C$  in Berührung gebracht wird, oder  
(b) mit dem Hydroxylaminderivat  $H_2NOR^{12}$ , Dicyclohexylcarbodiimid und 1-Hydroxybenzotriazol etwa 1-24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa  $0-30^\circ C$  in Berührung gebracht wird.
- 30 20. Verfahren von Anspruch 1, bei welchem  $R^1$



ist,

X eine Kohlenstoff-Kohlenstoff-Einfachbindung,  $-CO-$ ,  $-O-$ ,  $-S-$  oder  $-NH-$  ist,  
 $R^2$ ,  $R^3$ ,  $R^6$  und  $R^7$  wie in Anspruch 1 definiert sind und  
 $R^8$   $(CH_2)_nOR^{11}$ ,  $(CH_2)_nOCOR^{14}$ ,  $(CH_2)_nCH(OH)R^{16}$ ,  $(CH_2)_nCOR^{16}$ ,  $(CH_2)_nCl$ ,  $(CH_2)_nCN$ ,  $CHO$  ist.

45

21. Verfahren von Anspruch 20, bei welchem  $R^8$   $(CH_2)_nOH$  ist und das Produkt der Formel 3 mit einem Alkohol  $R^{11}OH$  in wasserfreiem Zustand in Anwesenheit einer starken Säure oder einer Lewissäure in Berührung gebracht wird, gefolgt von der Verseifung irgendwelcher gleichzeitig gebildeter oder im Zwischenprodukt 3 vorhandener Gruppen  $CO_2R^{14}$  unter Bilden der entsprechenden Verbindung der Formel 3, worin  $R^8$   $(CH_2)_nOR^{11}$  ist und  $R^{11}$  nicht H ist.
- 50
22. Verfahren von Anspruch 20, bei welchem  $R^8$   $(CH_2)_nOR^{11}$  ist und  $R^{11}$  nicht H ist und das Produkt der Formel 3 mit einem wäßrigen sauren Medium bei einer Temperatur im Bereich von etwa  $25^\circ C$  und der Rückflußtemperatur des Lösungsmittels während eines Zeitraums von etwa 0,5-24 Stunden unter Bilden der entsprechenden Verbindung der Formel 3 in Berührung gebracht wird, worin  $R^8$   $(CH_2)_nOH$  ist.
- 55


23. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n OH$  ist und das Produkt der Formel 3 mit  
 (a) einem Carbonsäureanhydrid  $(R^{14}CO)_2O$  oder -chlorid  $R^{14}COCl$  in einem Lösungsmittel in  
 Anwesenheit einer Base etwa 0,5-24 Stunden bei einer Temperatur im Bereich von etwa  $0^\circ C$  und  
 der Rückflußtemperatur des Lösungsmittels oder  
 5 (b) einer Carbonsäure  $R^{14}CO_2H$  unter wasserfreien Bedingungen in Anwesenheit einer starken Säure  
 oder Lewisäure etwa 0,5 bis 24 Stunden bei etwa  $0^\circ$ - $100^\circ C$  unter Bilden der entsprechenden  
 Verbindung, in welcher  $R^8 (CH_2)_n OCOR^{14}$  ist, in Berührung gebracht wird.
24. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n OCOR^{14}$  ist und das Produkt der Formel 3 mit  
 10 wäßriger Säure oder Alkali unter Bilden der entsprechenden Verbindung, worin  $R^8 (CH_2)_n OH$  ist, in  
 Berührung gebracht wird.
25. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n OH$  ist und das Produkt der Formel 3 mit einem  
 Oxidationsmittel etwa 1-200 Stunden bei einer Temperatur von etwa  $25-45^\circ C$  unter Herstellen einer  
 15 entsprechenden Verbindung der Formel 3, in welcher  $R^8 (CH_2)_{n-1} COR^{16}$  ist und  $R^{16} H$  ist, in Berührung  
 gebracht wird.
26. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n COR^{16}$  ist und  $R^{16} H$  ist und das Produkt der  
 Formel 3 mit einer metallorganischen Verbindung  $R^{16}P$ , in welcher  $P$   $MgBr$  oder  $Li$  ist, etwa 0,5-24  
 20 Stunden in einem Lösungsmittel bei einer Temperatur von etwa  $-78^\circ C$  bis  $100^\circ C$  unter Bilden einer  
 Verbindung der Formel 3 in Berührung gebracht wird, in der  $R^8 (CH_2)_n CH(OH)R^{16}$  ist und  $R^{16}$  nicht  $H$   
 ist.
27. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n CH(OH)R^{16}$  ist und  $R^{16}$  nicht  $H$  ist und das Produkt  
 25 der Formel 3 mit einem Oxidationsmittel in einem Lösungsmittel unter Bilden einer entsprechenden  
 Verbindung der Formel 3 in Berührung gebracht wird, in der  $R^8 (CH_2)_n COR^{16}$  ist und  $R^{16}$  nicht  $H$  ist.
28. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n COR^{16}$  ist und  $R^{16} H$  ist und die Verbindung der  
 Formel 3 mit einem Oxidationsmittel in einem Lösungsmittel unter Bilden einer entsprechenden  
 30 Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_n COR^{16}$  ist und  $R^{16} OH$  ist.
29. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n COR^{16}$  ist und  $R^{16} OH$  ist und die Verbindung der  
 Formel 3 etwa 5 Minuten bis etwa 24 Stunden mit Thionylchlorid im Überschuß oder in einem anderen  
 Lösungsmittel bei einer Temperatur im Bereich von etwa  $0^\circ C$  bis zur Rückflußtemperatur des  
 35 Lösungsmittels unter Bilden einer entsprechenden Verbindung der Formel 3 in Berührung gebracht  
 wird, in welcher  $R^8 (CH_2)_n COCl$  ist, gefolgt von der Berührung der letzteren mit einem Amin  $NHR^{18}R^{19}$   
 im Überschuß oder in einem Lösungsmittel bei Temperaturen im Bereich von etwa  $0^\circ C$  und der  
 Rückflußtemperatur des Lösungsmittels während etwa 5 Minuten bis 24 Stunden unter Bilden einer  
 entsprechenden Verbindung der Formel 3, in welcher  $R^8 (CH_2)_n CONR^{18}R^{19}$  ist.  
 40
30. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_n OR^{11}$  ist und  $R^{11} H$  ist und das Produkt der Formel  
 3 etwa 0,5-24 Stunden mit Thionylchlorid im Überschuß oder in einem Lösungsmittel bei einer  
 Temperatur im Bereich von etwa  $20^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels unter Bilden  
 einer Zwischenproduktverbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_n Cl$  ist.  
 45
31. Verfahren von Anspruch 30, bei welchem die Verbindung der Formel 3, worin  $R^8 (CH_2)_m Cl$  ist, mit  
 Imidazol, 1,2,3-Triazol, 1,2,4-Triazol, Tetrazol oder Phthalimid etwa 1-24 Stunden in Anwesenheit einer  
 Base in einem Lösungsmittel bei Temperaturen im Bereich von etwa  $55^\circ C$  bis zur Rückflußtemperatur  
 des Lösungsmittels unter Herstellen einer entsprechenden Verbindung der Formel 3 in Berührung  
 50 gebracht wird, in welcher  $R^8 (CH_2)_m$ -imidazol,  $(CH_2)_m$ -triazol,  $(CH_2)_m$ -tetrazol oder  $(CH_2)_m$ -phthalimid ist.
32. Verfahren von Anspruch 30, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_n Cl$  ist, mit  
 dem Natrium- oder Kaliumsalz eines Mercaptans  $R^{15}SH$  etwa 1-24 Stunden in einem Lösungsmittel bei  
 einer Temperatur im Bereich von etwa  $25-100^\circ C$  unter Bilden einer Verbindung der Formel 3 in  
 55 Berührung gebracht wird, in welcher  $R^8 (CH_2)_n SR^{15}$  ist.
33. Verfahren von Anspruch 30, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_n Cl$  ist, etwa  
 1-24 Stunden mit einem Alkalimetallcyanid in einem Lösungsmittel bei einer Temperatur im Bereich




von etwa 20-100 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_nCN$  ist, und die letztere Verbindung zu der entsprechenden Verbindung der Formel 3 hydrolysiert wird, in welcher  $R^8 (CH_2)_nCOR^{16}$  ist und  $R^{16} OH$  ist.

- 5    34. Verfahren von Anspruch 30, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_{n-1}Cl$  ist, etwa 0,5-24 Stunden mit dem Natrium- oder Kaliumsalz eines Malonsäuredialkylesters in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 20-100 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_nCH(CO_2Alkyl)_2$  ist, gefolgt von der Verseifung der letzteren mit wäßrigem Alkali bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels, gefolgt vom Ansäuern mit Mineralsäure unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nCH(CO_2H)_2$  ist, gefolgt vom Erhitzen der letzteren auf etwa 120 °C oder in verdünnter Mineralsäure bei Rückflußtemperatur unter Bilden eines Produkts der Formel 3, in welcher  $R^8 (CH_2)_nCOR^{16}$  ist und  $R^{16} OH$  ist.
- 10    35. Verfahren von Beispiel 20, in welchem  $R^8 (CH_2)_nCN$  ist und die Verbindung der Formel 3 mit Natriumazid und Ammoniumchlorid etwa 1 Stunde bis etwa 10 Tage in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 30 °C und der Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Erfindung in Berührung gebracht wird, in welcher  $R^8 (CH_2)_n$ -tetrazol ist.
- 15    36. Verfahren von Anspruch 20, bei welchem  $R^8 -CHO$  ist und die Verbindung der Formel 3 mit einem Methylenphosphoran  $(C_6H_5)_3P=CH(CH_2)_sCHR^{14}OR^{15}$  oder  $(C_6H_5)_3P=CH(CH_2)_sCOR^{16}$  etwa 1-24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 -CH=CH(CH_2)_sCHR^{14}OR^{15}$  oder  $-CH=CH(CH_2)_sCOR^{16}$  ist, außer wo  $R^{15} H$  ist und  $R^{16} OH$  ist, und gegebenenfalls anschließend dem In-Berührung-Bringen der Verbindung der Formel 3, in welcher  $R^8 -CH=CH(CH_2)_sCOR^{16}$  ist, mit einem Reduktionsmittel in einem Lösungsmittel bei einer Temperatur von etwa 0 °-25 °C während etwa 0,5-24 Stunden unter Bilden eines Produkts der Formel 3, in welcher  $R^8 -CH=CH(CH_2)_sCHR^{14}OH$  ist.
- 20    37. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_mOH$  ist und die Verbindung der Formel 3 mit einem Fluorierungsmittel über einen Zeitraum von etwa 0,5-24 Stunden in einem Lösungsmittel bei einer Temperatur in dem Bereich von etwa -30 °C bis 25 °C unter Bilden einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^8 (CH_2)_mF$  ist.
- 25    38. Verfahren von Anspruch 20, bei welchem die Verbindung der Formel 3, in welcher  $R^8 (CH_2)_mCl$  ist, etwa 1-24 Stunden mit Silbernitrat in einem dipolaren, aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 25-80 °C unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_mONO_2$  ist, in Berührung gebracht wird.
- 30    39. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_nOH$  ist und die Verbindung der Formel 3 mit einem Isocyanat der Formel  $R^{10}NCO$  in einem Lösungsmittel bei einer Temperatur im Bereich von etwa 25 °C bis zur Rückflußtemperatur des Lösungsmittels über einen Zeitraum von etwa 5 Minuten bis etwa 24 Stunden unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nOCONHR^{10}$  ist, in Berührung gebracht wird.
- 35    40. Verfahren von Anspruch 20, bei welchem die Verbindung, in welcher  $R^8 (CH_2)_nCl$  ist, mit einem Amin  $R^{11}NH_2$  in überschüssigem Amin oder in einem anderen Lösungsmittel über einen Zeitraum von etwa 1-24 Stunden bei einer Temperatur im Bereich von etwa 0 °C bis zur Rückflußtemperatur des Lösungsmittels unter Bilden eines Zwischenprodukts der Formel 3, in welcher  $R^8 (CH_2)_nNHR^{11}$  ist, in Berührung gebracht wird.
- 40    41. Verfahren von Anspruch 20, bei welchem  $R^8 (CH_2)_nCl$  ist und die Verbindung der Formel 3 etwa 1-24 Stunden mit einem Alkalimetallazid in einem aprotischen Lösungsmittel bei einer Temperatur im Bereich von etwa 25-80 °C unter Bilden einer Verbindung der Formel 3, in welcher  $R^8 (CH_2)_nN_3$  ist, in Berührung gebracht wird und letztere mit einem Reduktionsmittel unter Bilden eines Zwischenproduktes der Formel 3, in welcher  $R^8 (CH_2)_nNH_2$  ist, in Berührung gebracht wird.
- 45    50    55


42. Verfahren von Anspruch 40 oder 41, bei welchem  $R^8$   $(CH_2)_nNHR^{11}$  oder  $(CH_2)_nNH_2$  ist und die Verbindung der Formel 3 mit einem Chlorameisensäureester der Formel  $R^{10}OCOCI$  oder einem Sulfonylderivat der Formel  $R^{10}SO_2Cl$  oder  $(R^{10}SO_2)O$  etwa 5 Minuten bis etwa 24 Stunden in einem Lösungsmittel in Anwesenheit einer Base bei einer Temperatur im Bereich von etwa  $0^\circ C$  bis zur Rückflußtemperatur eines Lösungsmittels unter Bilden einer Verbindung der Formel 3, in welcher  $R^8$  -  $(CH_2)_nNR^{11}CO_2R^{10}$  oder  $-(CH_2)_nNR^{11}SO_2R^{10}$  ist, in Berührung gebracht wird.
43. Verfahren von Anspruch 40 oder 41, bei welchem die Verbindung der Formel 3 mit  $R^8$  gleich  $-(CH_2)_nNHR^{11}$  oder  $(CH_2)_nNH_2$  mit einem Isocyanat oder Isothiocyanat  $R^{10}NCY$  etwa 5 Minuten bis etwa 24 Stunden in einem Lösungsmittel bei einer Temperatur im Bereich von etwa  $25^\circ C$  bis zur Rückflußtemperatur des Lösungsmittels unter Bilden einer Verbindung der Formel 3, in welcher  $R^8$   $-(CH_2)_nNR^{11}CYNHR^{10}$  ist, in Berührung gebracht wird.
44. Verfahren von Anspruch 1, bei welchem  $R^1$   $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$  und  $R^8$  wie in Anspruch 1 definiert sind, in welchem die Verbindung der Formel 3, worin  $R^1$   $NO_2$  ist, mittels Eisen und Essigsäure, Zinn(II)-chlorid oder Wasserstoff und Palladium zu einer Verbindung der Formel 3 reduziert wird, worin  $R^1$   $NH_2$  ist, und letztere mit einem geeigneten Säureanhydrid, wie Phthalsäureanhydrid oder ein substituiertes Phthalsäureanhydrid, in einem Lösungsmittel oder mit einem geeigneten Säurechlorid, wie etwa substituiertes Anthranilsäurechlorid, in Anwesenheit wäßrigen Alkalis oder einer Base oder mit einer geeignet substituierten Phthal- oder Anthranilsäure in Anwesenheit von Dicyclohexylcarbodiimid in einem Lösungsmittel unter Herstellen einer Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^1$



45. Verfahren von Anspruch 1, bei welchem  $R^1$   $OCH_2C_6H_5$  ist,  $R^2$  und  $R^3$  H sind und  $R^6$ ,  $R^7$  und  $R^8$  wie in Anspruch 1 definiert sind und die sich daraus ergebende Verbindung der Formel 3 mit  $R^1$  gleich  $OCH_2C_6H_5$  mit Trifluoressigsäure bei Rückflußtemperatur über einen Zeitraum von etwa 0,2-1 Stunde oder mit Wasserstoff und Palladium unter Bilden der entsprechenden Verbindung der Formel 3 in Berührung gebracht wird, in welcher  $R^1$  OH ist, und letztere mit einer Base bei etwa  $25^\circ C$  und einem geeigneten Benzylhalogenid der Formel








ist, und X -OCH<sub>2</sub>- ist.

46. Verfahren von Anspruch 1, bei welchem R<sup>8</sup> -CHO ist, wobei das Benzylderivat der Formel 2 an das Imidazolderivat der Formel 1 vorzugsweise an das Stickstoffatom gebunden ist, welches dem Kohlenstoffatom des Imidazolrings, an welches R<sup>8</sup> gebunden ist, benachbart ist.

# Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE


20 1. Un dérivé anti-hypertenseur de formule:



(I)








dans laquelle:


45 R<sup>1</sup> est  
R<sup>1</sup> is -4-CO<sub>2</sub>H; -4-CO<sub>2</sub>R<sup>9</sup>;



-SO<sub>3</sub>H, -C(CF<sub>3</sub>)<sub>2</sub>OH;

55





5

10

15

20

25

30

35





40

45



50

55


(l-isomère);



ou

R<sup>2</sup>

est un atome d'hydrogène; de chlore; de brome; d'iode; de fluor; ou un groupe NO<sub>2</sub>; un radical alkyle comprenant de 1 à 4 atomes de carbone; acyloxy comprenant de 1 à 4 atomes de carbone, alcoxy comprenant de 1 à 4 atomes de carbone; CO<sub>2</sub>H; CO<sub>2</sub>R<sup>9</sup>; NHSO<sub>2</sub>CH<sub>3</sub>; NHSO<sub>2</sub>CF<sub>3</sub>; CONHOR<sup>12</sup>; SO<sub>2</sub>NH<sub>2</sub>;

R<sup>3</sup>

aryle; ou furyle;

est un atome d'hydrogène; de chlore, de brome, d'iode ou de fluor; un radical alkyle comprenant de 1 à 4 atomes de carbone ou alcoxy comprenant de 1 à 4 atomes de carbone;

R<sup>4</sup>est CN, NO<sub>2</sub> ou CO<sub>2</sub>R<sup>11</sup>;R<sup>5</sup>

est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, alkényle ou alkynyle comprenant de 2 à 4 atomes de carbone;










R<sup>6</sup>

est un radical alkyle comprenant de 2 à 10 atomes de carbone; alkényle ou alkynyle comprenant de 3 à 10 atomes de carbone ou les mêmes groupes substitués par F ou un groupe CO<sub>2</sub>R<sup>14</sup>; un radical cycloalkyle comprenant de 3 à 8 atomes de carbone, cycloalkylalkyle comprenant de 4 à 10 atomes de carbone; cycloalkylalkényle ou cycloalkylalkynyle comprenant de 5 à 10 atomes de carbone; (CH<sub>2</sub>)<sub>s</sub>Z(CH<sub>2</sub>)<sub>m</sub>R<sup>5</sup> éventuellement substitué par un atome de fluor ou le groupe CO<sub>2</sub>R<sup>14</sup>; benzyle ou benzyle substitué sur le cycle phényle avec 1 ou 2 atomes d'halogènes, un radical alcoxy comprenant de 1 à 4 atomes de carbone, alkyle comprenant de 1 à 4 atomes de carbone, ou nitro;

R<sup>7</sup>


est un atome d'hydrogène, de fluor, de chlore, de brome ou d'iode, ou un groupe

$\text{NO}_2$ ,  $\text{CF}_3$  ou  $\text{CN}$ ;  
 $\text{R}^8$  est un atome d'hydrogène, un groupe  $\text{CN}$ , un radical alkyle comprenant de 1 à 10 atomes de carbone, alkényle comprenant de 3 à 10 atomes de carbone, ou les mêmes groupes substitués par  $\text{F}$ ; phénylalkényle dans lequel la portion aliphatique comprend de 2 à 6 atomes de carbone;  $-\text{C}(\text{CH}_2)_m\text{-imidazole-1-yle}$ ;  $-(\text{CH}_2)_m\text{-1,2,3-triazolyle}$  éventuellement substitué par un ou deux groupes choisis parmi  $\text{CO}_2\text{CH}_3$  ou alkyle comprenant de 1 à 4 atomes de carbone;  $-(\text{CH}_2)_m\text{-tétrazolyle}$ ;  $-(\text{CH}_2)_n\text{OR}^{11}$ ;




$\text{R}^9$  est

5



- 10 R<sup>10</sup> est un radical alkyle comprenant de 1 à 6 atomes de carbone ou perfluoroalkyle comprenant de 1 à 6 atomes de carbone, 1-adamantyle, 1-naphtyle, 1-(1-naphtyl)-éthyle ou (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>;
- R<sup>11</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou benzyle;
- R<sup>12</sup> est un atome d'hydrogène, un radical méthyle ou benzyle;
- 15 R<sup>13</sup> est -CO<sub>2</sub>H; -CO<sub>2</sub>R<sup>9</sup>; -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>R<sup>9</sup>;


15



20

-SO<sub>3</sub>H;


25




30

-PO<sub>3</sub>H; -C(CF<sub>3</sub>)<sub>2</sub>OH; -NHSO<sub>2</sub>CH<sub>3</sub>; -NHSO<sub>2</sub>CF<sub>3</sub>; -NHCOCF<sub>3</sub>; -CONHOR<sup>12</sup>; -SO<sub>2</sub>NH<sub>2</sub>;

35




40



45

-CONHNHSO<sub>2</sub>CF<sub>3</sub>;


50




55

- R<sup>14</sup> est un atome d'hydrogène, un radical alkyle ou perfluoroalkyle comprenant de 1 à 8 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou


- benzyle;  
 R<sup>15</sup> est un atome d'hydrogène, un radical alkyle comprenant de 3 à 6 atomes de carbone, phényle, benzyle, acyle comprenant de 1 à 4 atomes de carbone, phénacyle;  
 5 R<sup>16</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>, OR<sup>17</sup>, ou NR<sup>18</sup>R<sup>19</sup>;  
 R<sup>17</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou benzyle;  
 10 R<sup>18</sup> et R<sup>19</sup>, indépendamment l'un de l'autre, sont un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, phényle, benzyle, α-méthylbenzyle ou pris ensemble forment un cycle de formule:




- 20 Q est NR<sup>20</sup>, O ou CH<sub>2</sub>;  
 R<sup>20</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou un radical phényle;  
 R<sup>21</sup> est un radical alkyle comprenant de 1 à 6 atomes de carbone, -NR<sup>22</sup>R<sup>23</sup>, ou



- 30 R<sup>22</sup> et R<sup>23</sup> sont, indépendamment l'un de l'autre, un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, benzyle, ou sont pris ensemble sous la forme de (CH<sub>2</sub>)<sub>u</sub> dans lequel u est varié de 3 à 6;  
 R<sup>24</sup> est un atome d'hydrogène, CH<sub>3</sub> ou -C<sub>6</sub>H<sub>5</sub>;  
 35 R<sup>25</sup> est NR<sup>27</sup>R<sup>28</sup>, OR<sup>28</sup>, NHCONH<sub>2</sub>, NHCSNH<sub>2</sub>;




- R<sup>26</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, un radical benzyle ou allyle;  
 45 R<sup>27</sup> et R<sup>28</sup> sont, indépendamment, un atome d'hydrogène, un radical alkyle comprenant de 1 à 5 atomes de carbone, ou un radical phényle;  
 R<sup>29</sup> et R<sup>30</sup> sont, indépendamment, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou pris ensemble sont -(CH<sub>2</sub>)<sub>q</sub>;  
 R<sup>31</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes d'hydrogène, -CH<sub>2</sub>CH=CH<sub>2</sub> ou -CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>R<sup>32</sup>;  
 50 R<sup>32</sup> est un atome d'hydrogène, NO<sub>2</sub>, NH<sub>2</sub>, OH ou OCH<sub>3</sub>;  
 X est une simple liaison carbone-carbone, -CO-, -O-, -S-, -NH-, -N-,



-CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHC(R<sup>27</sup>)(R<sup>28</sup>), -NR<sup>23</sup>SO<sub>2</sub>-, -SO<sub>2</sub>NR<sup>23</sup>-, -C(R<sup>27</sup>)(R<sup>28</sup>)NH-,  
-CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, -CF<sub>2</sub>CF<sub>2</sub>-,

A.

5



15 Y est O ou S;  
Z est O, NR<sup>11</sup> ou S;  
m est compris entre 1 et 5, bornes incluses;  
n est compris entre 1 et 10, bornes incluses;  
p est compris entre 0 et 3, bornes incluses;  
20 q est compris entre 2 et 3, bornes incluses;  
r est compris entre 0 et 2, bornes incluses;  
s est compris entre 0 et 5, bornes incluses;  
t est égal à 0 ou 1,


ainsi que les sels pharmaceutiquement acceptables de ces dérivés;

25

étant entendu que:

- (1) le groupe R<sup>1</sup> n'est pas en position ortho  
(2) lorsque R<sup>1</sup> est


30



35

X est une simple liaison, et R<sup>13</sup> est CO<sub>2</sub>H, ou

40




45

auquel cas R<sup>13</sup> doit être en position ortho ou méta; ou, lorsque R<sup>1</sup> et X sont tels qu'indiqués ci-dessus, et que R<sup>13</sup> est NHSO<sub>2</sub>CF<sub>3</sub> ou NHSO<sub>2</sub>CH<sub>3</sub>, le substituant R<sup>13</sup> doit être un groupe en position ortho;

- (2) lorsque R<sup>1</sup> est

50




55

et X autre qu'une simple liaison, alors le substituant  $R^{13}$  doit être en position ortho, sauf lorsque X est  $NR^{23}CO$  et que  $R^{13}$  est  $NHSO_2CF_3$  ou  $NHSO_2CH_3$ , auquel cas,  $R^{13}$  doit être en position ortho ou méta;

(4) lorsque  $R^1$  est 4- $CO_2H$  ou un sel en dérivant,  $R^6$  ne peut pas être un groupe S-alkyle;


(5) lorsque  $R^1$  est 4- $CO_2H$  ou un sel en dérivant, le substituant en position 4 de l'imidazole ne peut être un radical  $CH_2OH$ ,  $CH_2OCOCH_3$  ou  $CH_2CO_2H$ ;

(6) lorsque  $R^1$  est




X est  $-OCH_2-$ ,  $R^{13}$  est 2- $CO_2H$  et  $R^7$  est un atome d'hydrogène alors  $R^6$  ne peut pas être  $C_2H_5S$ ;

(7) lorsque  $R^1$  est




et  $R^6$  est un groupe n-hexyle, alors  $R^7$  et  $R^8$  ne peuvent être simultanément de l'hydrogène;

(8) lorsque  $R^1$  est




$R^6$  n'est pas un groupe méthoxybenzyle;

(9) le groupe  $R^6$  n'est pas





2. Un dérivé selon la revendication 1, présentant la formule:




dans laquelle:

R<sup>1</sup> est -CO<sub>2</sub>H; -NHSO<sub>2</sub>CF<sub>3</sub>;


20

25



30

ou



35



R<sup>6</sup> est un radical alkyle comprenant de 3 à 10 atomes de carbone, alkényle comprenant de 3 à 10 atomes de carbone, alkynyle comprenant de 3 à 10 atomes de carbone, cycloalkyle comprenant de 3 à 8 atomes de carbone, benzyle substitué sur le cycle phényle avec au plus deux groupes sélectionnés parmi les radicaux alcoxy comprenant de 1 à 4 atomes de carbone, halogène, alkyle comprenant de 1 à 4 atomes de carbone, et nitro;

40


R<sup>8</sup> est un radical phénylalkényle dans lequel la partie aliphatique comprend 2 à 4 atomes de carbone, -(CH<sub>2</sub>)<sub>m</sub>-imidazole-1-yle, -(CH<sub>2</sub>)<sub>m</sub>-1,2,3-triazolyle éventuellement substitué par un ou deux groupes choisis parmi les composés CO<sub>2</sub>CH<sub>3</sub> ou alkyle comprenant de 1 à 4 atomes de carbone, (CH<sub>2</sub>)<sub>m</sub>-tétrazolyle, -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>;

45

50




55






5



$R^{13}$  est  $-CO_2H$ ,  $-CO_2R^9$ ,  $NHSO_2CF_3$ ; et


10




15

$R^{16}$  H, un radical alkyle comprenant de 1 à 5 atomes de carbone,  $OR^{17}$  ou  $NR^{18}R^{19}$ ;  
X est une simple liaison carbone-carbone,  $-CO-$ ,


20



25



30



$-OCH_2-$ ,  $-CH_2O-$ ,  $-O-$ ,  $-SCH_2-$ ,  $-CH_2S-$ ,  $-NHCH_2-$ ,  $-CH_2NH-$  ou  $-CH=CH-$ ;  
ainsi que les sels pharmaceutiquement acceptables en dérivant.

35

3. Un composé selon la revendication 2, dans lequel:

$R^2$  est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, un atome d'halogène ou un radical alcoxy comprenant de 1 à 4 atomes de carbone;


$R^6$  est un radical alkyle, alkényle ou alkynyle comprenant de 3 à 7 atomes de carbone;

40

$R^7$  est un atome d'hydrogène, de chlore, de brome, d'iode ou  $CF_3$ ;

$R^8$  est  $-(CH_2)_mOR^{11}$ ;


45




50




55



- or -COR<sup>16</sup>; ou - COR<sup>16</sup>;  
 R<sup>10</sup> est CF<sub>3</sub>, un radical alkyle comprenant de 1 à 6 atomes de carbone ou un radical phényle;  
 R<sup>11</sup> est un atome d'hydrogène ou un radical alkyle comprenant de 1 à 4 atomes de carbone;  
 R<sup>13</sup> est CO<sub>2</sub>H; CO<sub>2</sub>CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>; NHSO<sub>2</sub>CF<sub>3</sub> et



- R<sup>14</sup> est un atome d'hydrogène, ou un radical alkyle comprenant de 1 à 4 atomes de carbone;  
 R<sup>15</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou acyle comprenant de 1 à 4 atomes de carbone;  
 R<sup>16</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 5 atomes de carbone; OR<sup>17</sup>; ou



- m est compris entre 1 et 5, bornes incluses;  
 X est une simple liaison, -O-, -CO-, -NHCO-, ou -OCH<sub>2</sub>-;  
 et les sels pharmaceutiquement acceptables en dérivant.

4. Les dérivés selon les revendications 1 à 3, choisis dans le groupe comprenant:

2-butyl-4-chloro-[(2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-[(méthoxycarbonyl)aminométhyl]imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-[(propoxycarbonyl)-aminométhyl]imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-1-[(2'-carboxybiphényl-4-yl)méthyl]-imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-(1E-butényl)-4-chloro-1-[(2'-carboxybiphényl-4-yl) méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-(1E-butényl)-4-chloro-1-[(2'-carboxybiphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-propyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-propyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;


2-butyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-(1E-butényl)-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant; et

2-(1E-butényl)-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant.

5. Une composition pharmaceutique comprenant un support pharmaceutiquement acceptable et au moins un des composés selon les revendications 1 à 4.


6. Un procédé de préparation d'un composé selon les revendications 1 à 4, dans lequel  $r$  est égal à 1, qui consiste à mettre en contact un dérivé d'imidazole de formule 1 avec un dérivé benzylique de formule 2 dans un solvant, en présence d'une base, pendant environ 1 à 10 heures, à une température comprise dans un intervalle allant d'environ 20 °C à la température du reflux du solvant, pour former un benzylimidazole de formule 3:



dans lesquelles chacun des radicaux  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$  et  $R^8$  est stable dans les conditions réactionnelles et est un groupe tel que défini dans la revendication 1, ou une forme intermédiaire ou protégée en dérivant qui peut être transformée en un tel groupe, et dans lesquelles  $X^1$  est un halogène, un radical p-toluènesulfonyloxy; ou méthylsulfonyloxy et ensuite, le cas échéant, on transforme lesdites formes intermédiaires ou protégées des groupes  $R$  en des groupes  $R$  tels que définis dans la revendication 1.

7. Procédé selon la revendication 6, dans lequel les dérivés 1 et 2 sont mis en contact, en présence d'une base choisie dans le groupe comprenant = un hydrure métallique,  $MH$ , un alcoxyde métallique,  $MOR$ , le carbonate de sodium, le carbonate de potassium, la triéthylamine et la pyridine, dans un solvant aprotique dipolaire ou, lorsque la base est  $MOR$ , le solvant peut être un alcool,  $ROH$ , dans lequel  $M$  est le lithium, le sodium ou le potassium et  $R$  est un radical méthyle, éthyle ou t-butyle.

8. Procédé selon la revendication 6, dans lequel:  $R^1$  est:



$X$  est une simple liaison carbone-carbone,  $-CO-$ ,  $-O-$ ,  $-S-$  ou  $-NH-$ ;

$R^2$  et  $R^3$ , indépendamment l'un de l'autre, sont un atome d'hydrogène, de chlore, de brome, d'iode, le groupe  $CO_2R^{14}$ ,  $F$ ,  $NO_2$ , un radical alkyle comprenant de 1 à 4 atomes de carbone, alcoxy comprenant 1 à 4 atomes de carbone, aryle ou furyle;

$R^6$  et  $R^7$  sont tels que définis dans la revendication 1

$R^8$  est un radical alkyle comprenant 1 à 10 atomes de carbone ou alkényle comprenant 3 à 10 atomes de carbone, ou les mêmes groupes substitués par un atome de fluor; phénylalkényle dans lequel la partie aliphatique comprend 2 à 6 atomes de carbone;  $-(CH_2)_nOR^{11}$ ;  $-(CH_2)_nSR^{15}$ ; ou  $(CH_2)_nCN$ ;

$R^{11}$  est tel que défini dans la revendication 1

$R^{13}$  est un groupe  $CO_2R^{14}$ ,  $CN$ ,  $NO_2$ , trityltétrazole ou tétrazole de trialkylétain; et

$R^{14}$  et  $R^{15}$  sont tels que définis dans la revendication 20.

9. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CO_2R^{14}$  et le produit de formule 3 est mis au contact d'un dérivé alcalin dans un solvant hydroalcoolique ou mis au contact de  $CF_3CO_2H$  à une température d'environ 20 °C à la température du reflux du solvant, pendant environ 1 à 24 heures, cette étape étant suivie de l'amenée du pH du mélange à une valeur de 3 à 7, pour convertir le produit


en un produit correspondant dans lequel  $R^{13}$  est  $-CO_2H$ .

10. Procédé selon la revendication 9, dans lequel au moins l'un des substituants  $R^2$ ,  $R^3$  ou  $R^{13}$  dans la formule 1 consiste en  $-CO_2R^{14}$ , lequel est converti en  $-CO_2H$ .
- 5 11. Procédé selon la revendication 9, dans lequel  $R^{14}$  est le radical t-butyle et la réaction est effectuée dans  $CF_3CO_2H$ .
12. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est mis au contact de:
  - (i) un acide fort à la température du reflux du solvant durant environ 2 à 96 heures; ou
  - (ii) un dérivé alcalin fort dans un solvant formé d'alcool à une température d'environ  $20^\circ C$  à la température de reflux du solvant pendant environ 2 à 96 heures, étape que l'on fait suivre de l'amenée du pH à une valeur d'environ 3 - 7; ou
  - 15 (iii) de l'acide sulfurique suivi d'un traitement acide ou alcalin pour convertir le produit en le composé correspondant dans lequel  $R^{13}$  est du  $-CO_2H$ .
13. Procédé selon la revendication 12, dans lequel au moins un des substituants  $R^2$ ,  $R^3$  ou  $R^{13}$  consiste en  $-CO_2R^{14}$  et est converti en  $-CO_2H$ .
- 20 14. Procédé selon la revendication 12, dans lequel  $R^8$  est  $-(CH_2)_nCN$  et est converti en  $-(CH_2)_nCO_2H$ , ou est  $-(CH_2)_n-OR^{11}$  et est converti en  $(CH_2)_nOH$  lorsque  $R^{13}$  est converti en  $-CO_2H$ .
15. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est mis au contact d'un mélange formé de proportions équimolaires de l'azidure de sodium et du chlorure d'ammonium dans un solvant aprotique polaire à une température d'environ  $30^\circ C$  à la température de reflux du solvant durant environ 1 heure à 10 jours, aux fins de convertir le produit en le composé correspondant dans lequel  $R^{13}$  est le radical 5-tétrazolyle.
- 30 16. Procédé selon la revendication 15, dans lequel  $R^8$  est  $-(CH_2)_mCN$  et est converti en radical  $-(CH_2)_m$ -tétrazolyle lorsque  $R^{13}$  est converti en radical 5-tétrazolyle.
17. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est amené à réagir avec l'azidure de trialkylétain ou l'azidure de triarylétaïn, étape que l'on fait suivre d'une hydrolyse acide ou basique pour convertir le produit en un composé correspondant dans lequel  $R^{13}$  est le radical 5-tétrazolyle.
- 35 18. Procédé selon la revendication 17, dans lequel  $R^8$  est  $-(CH_2)_mCN$  et est converti en  $-(CH_2)_m$ -tétrazolyle lorsque  $R^{13}$  est converti en radical 5-tétrazolyle.
- 40 19. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-NO_2$  et le produit de formule 3 est mis au contact d'un agent réducteur pour former un second intermédiaire de formule 3, dans lequel  $R^{13}$  est  $NH_2$ , ce dernier étant mis au contact d'un anhydride  $(CH_3SO_2)_2O$  ou  $(CF_3SO_2)_2O$  ou d'un chlorure  $CH_3SO_2Cl$  ou  $CF_3SO_2Cl$  de l'acide sulfonique dans un solvant pour produire un composé dans lequel  $R^{13}$  est  $-NHSO_2CH_3$  ou  $-NHSO_2CF_3$ .
- 45 20. Procédé selon la revendication 19, dans lequel au moins un des substituants  $R^2$ ,  $R^3$  ou  $R^{13}$  est  $-NO_2$  et est converti en  $-NHSO_2CH_3$  ou  $-NHSO_2CF_3$ .
- 50 21. Procédé selon la revendication 9 ou 12, dans lequel le composé de formule 3, dans lequel  $R^{13}$  est  $CO_2H$ , est soit:
  - (a) mis au contact d'environ 1 à 4 équivalents de chlorure de thionyle dans un excès de chlorure de thionyle ou d'un autre solvant à une température d'environ  $20^\circ C$  à la température du reflux du solvant pendant une période d'environ 5 minutes à environ 2 heures pour former un intermédiaire de formule 3 dans lequel  $R^{13}$  est  $COCl$ , et ce dernier est mis au contact d'environ 2 à 10 équivalents d'un dérivé d'hydroxylamine  $H_2NOR^{12}$  dans un excès de dérivé d'hydroxylamine  $H_2NOR^{12}$  ou d'un autre solvant, à une température d'environ  $25$  à  $80^\circ C$  pendant environ 2 à 18 heures, soit
- 55

(b) mis au contact du dérivé d'hydroxylamine  $H_2NOR^{12}$ , de dicyclohexylcarbodiimide et de 1-hydroxybenzotriazole dans un solvant à une température d'environ 0 à 30 °C pendant environ 1 à 24 heures;

afin d'obtenir un composé dans lequel  $R^{13}$  est  $CONHOR^{12}$ .

22. Procédé selon la revendication 6, dans lequel:  $R^1$  est:



X est une simple liaison carbone-carbone, -CO-, -O-, -S- ou -NH-;

$R^2$ ,  $R^3$ ,  $R^6$  et  $R^7$  sont tels que définis dans la revendication 1 et

$R^8$  est  $(CH_2)_nOR^{11}$ ,  $(CH_2)_nOCOR^{14}$ ,  $(CH_2)_nCH(OH)R^{16}$ ,  $(CH_2)_nCOR^{16}$ ,  $(CH_2)_nCl$ ,  $(CH_2)_nCN$ , CHO.

23. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis en contact avec un alcool  $R^{11}OH$  à l'état anhydre en présence d'un acide fort ou d'un acide de Lewis, étape que l'on fait suivre d'une saponification de tout groupe  $CO_2R^{14}$  concomitamment formé ou présent sur l'intermédiaire 3, pour former le dérivé correspondant de formule 3, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  n'est pas H.

24. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  n'est pas H et dans lequel le produit de formule 3 est mis au contact d'un milieu aqueux acide à des températures d'environ 25 °C à la température de reflux du solvant pendant une période d'environ 0,5 à 24 heures pour former le dérivé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nOH$ .

25. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis au contact de:

(a) un anhydride d'acide carboxylique  $(R^{14}CO)_2O$  ou un chlorure  $R^{14}COCl$  dans un solvant en présence d'une base à une température d'environ 0 °C à la température de reflux du solvant pendant environ 0,5 à 24 heures; ou

(b) un acide carboxylique  $R^{14}CO_2H$  dans des conditions anhydres en présence d'un acide fort ou d'un acide de Lewis à environ 0 °C à 100 °C pendant environ 0,5 à 24 heures pour former le dérivé correspondant dans lequel  $R^8$  est  $(CH_2)_nOCOR^{14}$ .

26. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOCOR^{14}$  et le produit de formule 3 est mis au contact d'une solution aqueuse acide ou alcaline pour former le dérivé correspondant dans lequel  $R^8$  est  $(CH_2)_nOH$ .

27. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis au contact d'un agent oxydant à une température d'environ 25 °C à 45 °C pendant environ 1 à 200 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_{n-1}COR^{16}$  et  $R^{16}$  est H.

28. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est H et le produit de formule 3 est mis au contact d'un composé organométallique  $R^{16}P$  dans lequel P est MgBr ou Li dans un solvant, à une température d'environ -78 °C à 100 °C pendant environ 0,5 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(OH)R^{16}$  et  $R^{16}$  n'est pas H.

29. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCH(OH)R^{16}$  et  $R^{16}$  n'est pas H, et le produit de formule 3 est mis au contact d'un agent oxydant dans un solvant pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  n'est pas de l'hydrogène.



30. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est de l'hydrogène et le produit de formule 3 est mis au contact d'un agent oxydant dans un solvant pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est OH.
- 5 31. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est OH et le produit de formule 3 est mis au contact de chlorure de thionyle en excès ou dans un autre solvant à une température d'environ  $0^\circ C$  à la température de reflux du solvant pendant environ 5 minutes à environ 24 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOCl$ , étape que l'on fait suivre d'un contact de ce dernier composé avec une amine  $NHR^{18}R^{19}$  en excès  
10 dans un solvant à une température d'environ  $0^\circ C$  à la température de reflux du solvant pendant environ 5 minutes à environ 24 heures pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCONR^{18}R^{19}$ .
- 15 32. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  est un atome d'hydrogène et le produit de formule 3 est mis au contact de chlorure de thionyle en excès ou dans un solvant à une température d'environ  $20^\circ C$  à la température de reflux du solvant pendant environ 0,5 à 24 heures pour produire un composé intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$ .
- 20 33. Procédé selon la revendication 32, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mCl$  est mis au contact d'un imidazole, de 1,2,3-triazole, de 1,2,4-triazole ou de phthalimide en présence d'une base dans un solvant à une température d'environ  $55^\circ C$  à la température de reflux du solvant pendant environ 1 à 24 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_m$ -imidazole,  $(CH_2)_m$ -triazole,  $(CH_2)_m$ -tétrazole ou  $(CH_2)_m$ -phthalimide.
- 25 34. Procédé selon la revendication 32, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$  est mis au contact d'un sel de sodium ou de potassium d'un mercaptan  $R^{15}SH$ , dans un solvant, à une température d'environ  $25^\circ C$  à  $100^\circ C$ , pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_n-SR^{15}$ .
- 30 35. Procédé selon la revendication 32, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$  est mis au contact d'un cyanure de métal alcalin dans un solvant à une température d'environ  $20^\circ C$  à  $100^\circ C$ , pendant 1 à 24 heures, pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCN$  et ce dernier composé est hydrolysé en un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^6$  et  $R^{16}$  est OH.
- 35 36. Procédé selon la revendication 32, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_{n-1}Cl$  est mis au contact d'un sel de sodium ou de potassium d'un malonate de dialkyle, dans un solvant, à une température de  $20^\circ C$  à  $100^\circ C$ , pendant environ 0,5 à 24 heures, pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(CO_2alkyl)_2$ , étape que l'on fait suivre d'une saponification de ce  
40 dernier composé par action d'une solution alcaline aqueuse à une température d'environ  $25^\circ C$  à la température de reflux du solvant, ce que l'on fait suivre d'une acidification à l'aide d'un acide minéral pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(CO_2H)_2$  suivie d'un chauffage de ce dernier composé à environ  $120^\circ C$  dans une solution diluée d'un acide minéral à la température du reflux pour former un produit de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est OH.
- 45 37. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCN$  et le composé de formule 3 est mis au contact d'azidure de sodium et de chlorure d'ammonium dans un solvant à une température d'environ  $30^\circ C$  à la température de reflux du solvant pendant environ 1 heure à environ 10 jours pour former un composé selon la présente invention dans lequel  $R^8$  est  $(CH_2)_n$ -tétrazole.
- 50 38. Procédé selon la revendication 22, dans lequel  $R^8$  est  $-CHO$  et le composé de formule 3 est mis au contact d'un méthylène phosphorane  $(C_6H_5)_3P=CH(CH_2)_sCHR^{14}OR^{15}$  ou  $(C_6H_5)_3P=CH(CH_2)_sCOR^{16}$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCHR^{14}OR^{15}$  ou  $-CH=CH(CH_2)_sCOR^{16}$ , excepté dans le cas où  $R^{15}$  est H et  $R^{16}$  est OH, et éventuellement mise en contact du composé de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCOR^{16}$   
55 avec un agent réducteur dans un solvant à une température d'environ  $0^\circ C$  à  $25^\circ C$  pendant environ 0,5 à 24 heures, pour former un produit de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCHR^{14}OH$ .

39. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_mOH$  et le composé de formule 3 est mis au contact d'un agent de fluoration dans un solvant à une température d'environ  $-30^\circ C$  à  $25^\circ C$  pendant une période d'environ 0,5 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mF$ .

40. Procédé selon la revendication 22, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mCl$ , est mis au contact de nitrate d'argent dans un solvant aprotique dipolaire à une température d'environ  $25^\circ C$  à  $80^\circ C$  pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mONO_2$ .

41. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le composé de formule 3 est mis au contact d'un isocyanate de formule  $R^{10}NCO$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant une période d'environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nOCONHR^{10}$ .

42. Procédé selon la revendication 22, dans lequel le composé où  $R^8$  est  $(CH_2)_nCl$  est mis au contact d'une amine  $R^{11}NH_2$  dans un excès d'amine ou dans un autre solvant pendant une période d'environ 1 à 24 heures à une température d'environ  $0^\circ C$  à la température de reflux du solvant pour former un intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_nNHR^{11}$ .


43. Procédé selon la revendication 22, dans lequel  $R^8$  est  $(CH_2)_nCl$  et le composé de formule 3 est mis au contact d'un azidure de métal alcalin dans un solvant aprotique à une température d'environ  $25^\circ C$  à  $80^\circ C$  pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nN_3$  et ce dernier est alors mis au contact d'un agent réducteur pour former un intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_nNH_2$ .

44. Procédé selon la revendication 42 ou 43, dans lequel  $R^8$  est  $(CH_2)_nNHR^{11}$  ou  $(CH_2)_nNH_2$  et le composé de formule 3 est mis au contact d'un chloroformiate de formule  $R^{10}OCOCl$  ou d'un dérivé sulfonyl de formule  $R^{10}SO_2Cl$  ou  $(R^{10}SO_2)O$  dans un solvant en présence d'une base à une température d'environ  $0^\circ C$  à la température du reflux dans le solvant pendant environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-(CH_2)_nNR^{11}CO_2R^{10}$  ou  $-(CH_2)_nNR^{11}SO_2R^{10}$ .

45. Procédé selon la revendication 42 ou 43, dans lequel le composé de formule 3 où  $R^8$  est  $(CH_2)_nNHR^{11}$  ou  $(CH_2)_nNH_2$  est mis en contact d'un isocyanate ou d'un isothiocyanate  $R^{10}NCY$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-(CH_2)_n-NR^{11}CYNHR^{10}$ .

46. Procédé selon la revendication 6, dans lequel  $R^1$  est  $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$  et  $R^8$  sont tels que définis dans la revendication 1 et dans lequel le composé de formule 3 où  $R^1$  est  $NO_2$  est réduit au moyen de fer et d'acide acétique, de chlorure stanneux ou d'hydrogène et de palladium en un composé de formule 3 dans lequel  $R^1$  est  $NH_2$ , le composé ainsi obtenu étant amené à réagir avec un anhydride d'acide convenable tel qu'un anhydride phtalique ou un anhydride phtalique substitué, dans un solvant, ou avec un chlorure d'acide approprié tel que le chlorure d'acide anthranilique substitué, en présence d'une solution aqueuse alcaline ou d'une base, ou avec un acide anthranilique ou phtalique convenablement substitué, en présence de dicyclohexylcarbodiimide dans un solvant pour produire un composé de formule 3 dans laquelle:

$R^1$  est




et




X est NHCO.

47. Procédé selon la revendication 6, dans lequel  $R^1$  est  $OCH_2C_6H_5$ ,  $R^2$  et  $R^3$  sont H, et  $R^6$ ,  $R^7$  et  $R^8$  sont tels que définis dans la revendication 1 et dans lequel le composé résultant de formule 3 où  $R^1$  est le radical  $OCH_2C_6H_5$  est mis au contact d'acide trifluoroacétique à la température du reflux pendant une période d'environ 0,2 à 1 heure, ou d'hydrogène et de palladium pour former le composé correspondant de formule 3 dans lequel  $R^1$  est OH et ce dernier est mis au contact d'une base à environ 25 °C et d'un halogénure benzylique convenable de formule:



pour produire le composé correspondant de formule 3 dans lequel:

$R^1$  est




X est  
et  
est  $-OCH_2-$ .

48. Procédé selon la revendication 6, dans lequel  $R^8$  est  $-CHO$ , de façon à ce que le dérivé benzylique de formule 2 se lie au dérivé imidazole de formule 1 de préférence sur l'atome d'azote adjacent à l'atome de carbone du cycle imidazole auquel  $R^8$  est lié.

#### Revendications pour l'Etat contractant suivant : ES


1. Un procédé de préparation d'un dérivé anti-hypertenseur de formule:



dans laquelle:

$R^1$  est  $-4-CO_2H$ ;  $-4-CO_2R^9$ ;


5



10

$-SO_3H$ ,  $-C(CF_3)_2OH$ ;


15



20


$-PO_3H$ ;

25




$4-NHSO_2CH_3$ ;  $4-NHSO_2CF_3$ ;  $-CONHOR^{12}$ ;  $-SO_2NH_2$ ;


30




35



40




45






50

$4-CONHNHSO_2CF_3$ ;

55





25  
30

$R^2$  est un atome d'hydrogène; de chlore; de brome; d'iode; de fluor; ou un groupe  $\text{NO}_2$ ; un radical alkyle comprenant de 1 à 4 atomes de carbone; acyloxy comprenant de 1 à 4 atomes de carbone;  $\text{CO}_2\text{H}$ ;  $\text{CO}_2\text{R}^9$ ;  $\text{NHSO}_2\text{CH}_3$ ;  $\text{NHSO}_2\text{CF}_3$ ;  $\text{CONHOR}^{12}$ ;  $\text{SO}_2\text{NH}_2$ ;



40

$R^3$  aryle; ou furyle; est un atome d'hydrogène; de chlore, de brome, d'iode ou de fluor; un radical alkyle comprenant de 1 à 4 atomes de carbone ou alcoxy comprenant de 1 à 4 atomes de carbone;

$R^4$  est  $\text{CN}$ ,  $\text{NO}_2$  ou  $\text{CO}_2\text{R}^{11}$ ;

45

$R^5$  est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, alkényle ou alkynyle comprenant de 2 à 4 atomes de carbone;

50










$R^6$  est un radical alkyle comprenant de 2 à 10 atomes de carbone; alkényle ou alkynyle comprenant de 3 à 10 atomes de carbone ou les mêmes groupes substitués par F ou un groupe  $\text{CO}_2\text{R}^{14}$ ; un radical cycloalkyle comprenant de 3 à 8 atomes de carbone, cycloalkylalkyle comprenant de 4 à 10 atomes de carbone; cycloalkylalkényle ou cycloalkylalkynyle comprenant de 5 à 10 atomes de carbone;  $(\text{CH}_2)_s\text{Z}(\text{CH}_2)_m\text{R}^5$  éventuellement substitué par un atome de fluor ou le groupe  $\text{CO}_2\text{R}^{14}$ ; benzyle ou benzyle substitué sur le cycle phényle avec 1 ou 2 atomes d'halogènes, un radical alcoxy comprenant de 1 à 4 atomes de carbone, alkyle comprenant de 1 à 4 atomes de carbone, ou nitro;

55


$R^7$  est un atome d'hydrogène, de fluor, de chlore, de brome ou d'iode, ou un groupe  $\text{NO}_2$ ,  $\text{CF}_3$  ou  $\text{CN}$ ;

$R^8$  est un atome d'hydrogène, un groupe  $\text{CN}$ , un radical alkyle comprenant de 1 à 10

atomes de carbone, alkényle comprenant de 3 à 10 atomes de carbone, ou les  
 mêmes groupes substitués par F; phénylalkényle dans lequel la portion aliphatique  
 comprend de 2 à 6 atomes de carbone;  $-C(CH_2)_m$ -imidazole-1-yle;  $-(CH_2)_m$ -1,2,3-  
 triazolyle éventuellement substitué par un ou deux groupes choisis parmi  $CO_2CH_3$  ou  
 alkyle comprenant de 1 à 4 atomes de carbone;  $-(CH_2)_m$ -tétrazolyle;  
 $-(CH_2)_nOR^{11}$ ;



$R^9$  est



5

R<sup>10</sup> est un radical alkyle comprenant de 1 à 6 atomes de carbone ou perfluoroalkyle comprenant de 1 à 6 atomes de carbone, 1-adamantyle, 1-naphtyle, 1-(1-naphtyl)-éthyle ou (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>;


10

R<sup>11</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou benzyle;

R<sup>12</sup> est un atome d'hydrogène, un radical méthyle ou benzyle;

R<sup>13</sup> est -CO<sub>2</sub>H; -CO<sub>2</sub>R<sup>9</sup>; -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>R<sup>9</sup>;


15



20



-SO<sub>3</sub>H;

25




-PO<sub>3</sub>H; -C(CF<sub>3</sub>)<sub>2</sub>OH; -NHSO<sub>2</sub>CH<sub>3</sub>; -NHSO<sub>2</sub>CF<sub>3</sub>; -NHCOCF<sub>3</sub>; -CONHOR<sup>12</sup>; -SO<sub>2</sub>NH<sub>2</sub>;

30




35



40

-CONHNHSO<sub>2</sub>CF<sub>3</sub>;

45



50

R<sup>14</sup> est un atome d'hydrogène, un radical alkyle ou perfluoroalkyle comprenant de 1 à 8 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou benzyle;


55

R<sup>15</sup> est un atome d'hydrogène, un radical alkyle comprenant de 3 à 6 atomes de carbone, phényle, benzyle, acyle comprenant de 1 à 4 atomes de carbone, phénacyle;

R<sup>16</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, (CH<sub>2</sub>)<sub>p</sub>C<sub>6</sub>H<sub>5</sub>, OR<sup>17</sup>, ou NR<sup>18</sup>R<sup>19</sup>;

R<sup>17</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, cycloalkyle comprenant de 3 à 6 atomes de carbone, phényle ou benzyle;


R<sup>18</sup> et R<sup>19</sup>, indépendamment l'un de l'autre, sont un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, phényle, benzyle, α-méthylbenzyle ou pris ensemble forment un cycle de formule:



Q est NR<sup>20</sup>, O ou CH<sub>2</sub>;

R<sup>20</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou un radical phényle;


R<sup>21</sup> est un radical alkyle comprenant de 1 à 6 atomes de carbone, -NR<sup>22</sup>R<sup>23</sup>, ou



R<sup>22</sup> et R<sup>23</sup> sont, indépendamment l'un de l'autre, un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, benzyle, ou sont pris ensemble sous la forme de (CH<sub>2</sub>)<sub>u</sub> dans lequel u est varié de 3 à 6;

R<sup>24</sup> est un atome d'hydrogène, CH<sub>3</sub> ou -C<sub>6</sub>H<sub>5</sub>;

R<sup>25</sup> est NR<sup>27</sup>R<sup>28</sup>, OR<sup>28</sup>, NHCONH<sub>2</sub>, NHCSNH<sub>2</sub>;



R<sup>26</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 6 atomes de carbone, un radical benzyle ou allyle;


R<sup>27</sup> et R<sup>28</sup> sont, indépendamment, un atome d'hydrogène, un radical alkyle comprenant de 1 à 5 atomes de carbone, ou un radical phényle;

R<sup>29</sup> et R<sup>30</sup> sont, indépendamment, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou pris ensemble sont -(CH<sub>2</sub>)<sub>q</sub>;


R<sup>31</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes d'hydrogène, -CH<sub>2</sub>CH=CH<sub>2</sub> ou -CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>R<sup>32</sup>;

R<sup>32</sup> est un atome d'hydrogène, NO<sub>2</sub>, NH<sub>2</sub>, OH ou OCH<sub>3</sub>;

X est une simple liaison carbone-carbone, -CO-, -O-, -S-, -NH-, -N-,



-CH<sub>2</sub>O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHC(R<sup>27</sup>)(R<sup>28</sup>), -NR<sup>23</sup>SO<sub>2</sub>-, -SO<sub>2</sub>NR<sup>23</sup>-, -C(R<sup>27</sup>)(R<sup>28</sup>)NH-,  
-CH=CH-, -CF=CF-, -CH=CF-, -CF=CH-, -CH<sub>2</sub>CH<sub>2</sub>-, -CF<sub>2</sub>CF<sub>2</sub>-;




- Y est O ou S;  
 Z est O, NR<sup>11</sup> ou S;  
 m est compris entre 1 et 5, bornes incluses;  
 n est compris entre 1 et 10, bornes incluses;  
 p est compris entre 0 et 3, bornes incluses;  
 q est compris entre 2 et 3, bornes incluses;  
 r est compris entre 0 et 2, bornes incluses;  
 s est compris entre 0 et 5, bornes incluses;  
 t est égal à 0 ou 1,


ainsi que les sels pharmaceutiquement acceptables de ces dérivés;

étant entendu que:

- (1) le groupe R<sup>1</sup> n'est pas en position ortho  
 (2) lorsque R<sup>1</sup> est




X est une simple liaison, et R<sup>13</sup> est CO<sub>2</sub>H, ou



auquel cas R<sup>13</sup> doit être en position ortho ou méta; ou, lorsque R<sup>1</sup> et X sont tels qu'indiqués ci-dessus, et que R<sup>13</sup> est NHSO<sub>2</sub>CF<sub>3</sub> ou NHSO<sub>2</sub>CH<sub>3</sub>, le substituant R<sup>13</sup> doit être un groupe en position ortho;

- (3) lorsque R<sup>1</sup> est




et X autre qu'une simple liaison, alors le substituant R<sup>13</sup> doit être en position ortho, sauf lorsque X est NR<sup>23</sup>CO et que R<sup>13</sup> est NHSO<sub>2</sub>CF<sub>3</sub> ou NHSO<sub>2</sub>CH<sub>3</sub>, auquel cas, R<sup>13</sup> doit être en position ortho ou méta;


- (4) lorsque R<sup>1</sup> est 4-CO<sub>2</sub>H ou un sel en dérivant, R<sup>6</sup> ne peut pas être un groupe S-alkyle;

- (5) lorsque R<sup>1</sup> est 4-CO<sub>2</sub>H ou un sel en dérivant, le substituant en position 4 de l'imidazole ne peut être un radical CH<sub>2</sub>OH, CH<sub>2</sub>OCOCH<sub>3</sub> ou CH<sub>2</sub>CO<sub>2</sub>H;


(6) lorsque R<sup>1</sup> est




X est -OCH<sub>2</sub>-, R<sup>13</sup> est 2-CO<sub>2</sub>H et R<sup>7</sup> est un atome d'hydrogène alors R<sup>6</sup> ne peut pas être C<sub>2</sub>H<sub>5</sub>S;  
(7) lorsque R<sup>1</sup> est



et R<sup>6</sup> est un groupe n-hexyle, alors R<sup>7</sup> et R<sup>8</sup> ne peuvent être simultanément de l'hydrogène;  
(8) lorsque R<sup>1</sup> est




R<sup>6</sup> n'est pas un groupe méthoxybenzyle;  
(9) le groupe R<sup>6</sup> n'est pas



ou CH<sub>2</sub>OH.

lequel procédé consiste à mettre en contact avec un dérivé d'imidazole de formule 1 avec un dérivé benzylique de formule 2 dans un solvant, en présence d'une base, pendant environ 1 à 10 heures, et à une température d'environ 20 °C à la température du reflux du solvant, pour former un benzylimidazole de formule 3:




dans lesquelles chacun des radicaux R<sup>1</sup>, R<sup>2</sup>, RY<sub>3</sub>H, R<sup>6</sup>, R<sup>7</sup>, et R<sup>8</sup> est stable dans les conditions réactionnelles et est un groupe tel que défini ci-dessus, ou une forme intermédiaire ou protégée en dérivant qui peut être transformée en un tel groupe, et dans lesquelles X<sup>1</sup> est un halogène, un radical p-toluènesulfonyloxy ou méthylsulfonyloxy; et ensuite, le cas échéant, on transforme lesdites formes



intermédiaires ou protégées des groupes R en des groupes R tels que définis ci-dessus.

2. Un procédé selon la revendication 1, dans lequel les composés obtenus présentent la formule:



dans laquelle:

20  $R^1$  est  $-CO_2H$ ;  $-NHSO_2CF_3$ ;







40  $R^6$  est un radical alkyle comprenant de 3 à 10 atomes de carbone, alkényle comprenant de 3 à 10 atomes de carbone, alkynyle comprenant de 3 à 10 atomes de carbone, cycloalkyle comprenant de 3 à 8 atomes de carbone, benzyle substitué sur le cycle phényle avec au plus deux groupes sélectionnés parmi les radicaux alcoxy comprenant de 1 à 4 atomes de carbone, halogène, alkyle comprenant de 1 à 4 atomes de carbone, et nitro;


45  $R^8$  est un radical phénylalkényle dans lequel la partie aliphatique comprend 2 à 4 atomes de carbone,  $-(CH_2)_m$ -imidazole-1-yle,  $-(CH_2)_m$ -1,2,3-triazolyle éventuellement substitué par un ou deux groupes choisis parmi les composés  $CO_2CH_3$  ou alkyle comprenant de 1 à 4 atomes de carbone,  $(CH_2)_m$ -tétrazolyle,  $-(CH_2)_nOR^{11}$ ;

50




55



R<sup>13</sup> est -CO<sub>2</sub>H, -CO<sub>2</sub>R<sup>9</sup>, NHSO<sub>2</sub>CF<sub>3</sub>; et ;






R<sup>16</sup> H, un radical alkyle comprenant de 1 à 5 atomes de carbone, OR<sup>17</sup> ou NR<sup>18</sup>R<sup>19</sup>;  
X est une simple liaison carbone-carbone, -CO-,



-OCH<sub>2</sub>-, -CH<sub>2</sub>O-, -O-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -NHCH<sub>2</sub>-, -CH<sub>2</sub>NH- ou -CH=CH-;  
ainsi que les sels pharmaceutiquement acceptables en dérivant.

3. Un procédé selon la revendication 2, dans lequel:

- R<sup>2</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, un atome d'halogène ou un radical alcoxy comprenant de 1 à 4 atomes de carbone;  
R<sup>6</sup> est un radical alkyle, alkényle ou alkynyle comprenant de 3 à 7 atomes de carbone;  
R<sup>7</sup> est un atome d'hydrogène, de chlore, de brome, d'iode ou CF<sub>3</sub>;  
R<sup>8</sup> est -(CH<sub>2</sub>)<sub>m</sub>OR<sup>11</sup>;



- 20
- ou - COR<sup>16</sup>;
- R<sup>10</sup> est CF<sub>3</sub>, un radical alkyle comprenant de 1 à 6 atomes de carbone ou un radical phényle;
- R<sup>11</sup> est un atome d'hydrogène ou un radical alkyle comprenant de 1 à 4 atomes de carbone;
- R<sup>13</sup> est CO<sub>2</sub>H; CO<sub>2</sub>CH<sub>2</sub>OCOC(CH<sub>3</sub>)<sub>3</sub>; N H S O<sub>2</sub> C F<sub>3</sub> et



- 30
- R<sup>14</sup> est un atome d'hydrogène, ou un radical alkyle comprenant de 1 à 4 atomes de carbone;
- R<sup>15</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 4 atomes de carbone, ou acyle comprenant de 1 à 4 atomes de carbone;
- R<sup>16</sup> est un atome d'hydrogène, un radical alkyle comprenant de 1 à 5 atomes de carbone; O R<sup>17</sup>;
- ou



- 40
- m est compris entre 1 et 5, bornes incluses;
- X est une simple liaison, -O-, -CO-, -NHCO-, ou -OCH<sub>2</sub>-;
- et les sels pharmaceutiquement acceptables en dérivant.

4. Un procédé selon les revendications 1 à 3, dans lequel les composés obtenus sont choisis dans le
- groupe comprenant:
- 2-butyl-4-chloro-[(2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-[(méthoxycarbonyl)aminométhyl]imidazole, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]-5-[(propoxycarbonyl)-aminométhyl]imidazole, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-butyl-4-chloro-1-[(2'-carboxybiphényl-4-yl)-méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-butyl-1-[(2'-carboxybiphényl-4-yl)méthyl]-imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;
- 2-(1E-butényl)-4-chloro-1-[(2'-carboxybiphényl-4-yl)méthyl]-5-(hydroxyméthyl)imidazole, ou un sel

pharmaceutiquement acceptable en dérivant;

2-(1E-butényl)-4-chloro-1-[(2'-carboxybiphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-propyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-(hydroxyméthyl)imidazole, ou un sel pharmaceutiquement acceptable en dérivant;

2-propyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;

2-butyl-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant;


2-(1E-butényl)-4-chloro-1-[2'-(1H-tétrazole-5-yl)biphényl-4-yl)méthyl]-5-hydroxyméthyl]imidazole, ou un sel pharmaceutiquement acceptable en dérivant; et

2-(1E-butényl)-4-chloro-1-[2'-(1H-tétrazole-5-yl)-biphényl-4-yl)méthyl]-imidazole-5-carboxaldéhyde, ou un sel pharmaceutiquement acceptable en dérivant.

5. Procédé selon la revendication 1, dans lequel les dérivés 1 et 2 sont mis en contact, en présence d'une base choisie dans le groupe comprenant = un hydruure métallique, MH, un alcoxyde métallique, MOR, le carbonate de sodium, le carbonate de potassium, la triéthylamine et la pyridine, dans un solvant aprotique dipolaire ou, lorsque la base est MOR, le solvant peut être un alcool, ROH, dans lequel M est le lithium, le sodium ou le potassium et R est un radical méthyle, éthyle ou t-butyle.

6. Procédé selon la revendication 1, dans lequel:

R<sup>1</sup> est:



X est une simple liaison carbone-carbone, -CO-, -O-, -S-ou -NH-;  
R<sup>2</sup> et R<sup>3</sup>, indépendamment l'un de l'autre, sont un atome d'hydrogène, de chlore, de brome, d'iode, le groupe CO<sub>2</sub>R<sup>14</sup>, F, NO<sub>2</sub>, un radical alkyle comprenant de 1 à 4 atomes de carbone, alcoxy comprenant 1 à 4 atomes de carbone, aryle ou furyle;

R<sup>6</sup> et R<sup>7</sup> sont tels que définis au-dessus  
R<sup>8</sup> est un radical alkyle comprenant 1 à 10 atomes de carbone ou alkényle comprenant 3 à 10 atomes de carbone, ou les mêmes groupes substitués par un atome de fluor; phénylalkényle dans lequel la partie aliphatique comprend 2 à 6 atomes de carbone; -(CH<sub>2</sub>)<sub>n</sub>OR<sup>11</sup>; -(CH<sub>2</sub>)<sub>n</sub>SR<sup>15</sup>; ou (CH<sub>2</sub>)<sub>n</sub>CN;


R<sup>11</sup> est tel que défini au-dessus  
R<sup>13</sup> est un groupe CO<sub>2</sub>R<sup>14</sup>, CN, NO<sub>2</sub>, trityltétrazole ou tétrazole de trialkylétain; et  
R<sup>14</sup> et R<sup>15</sup> sont tels que définis au-dessus

7. Procédé selon la revendication 6, dans lequel R<sup>13</sup> est -CO<sub>2</sub>R<sup>14</sup> et le produit de formule 3 est mis au contact d'un dérivé alcalin dans un solvant hydroalcoolique ou mis au contact de CF<sub>3</sub>CO<sub>2</sub>H à une température d'environ 20 °C à la température du reflux du solvant, pendant environ 1 à 24 heures, cette étape étant suivie de l'amenée du pH du mélange à une valeur de 3 à 7, pour convertir le produit en un produit correspondant dans lequel R<sup>13</sup> est -CO<sub>2</sub>H.

8. Procédé selon la revendication 7, dans lequel au moins l'un des substituants R<sup>2</sup>, R<sup>3</sup> ou R<sup>13</sup> dans la formule 1 consiste en -CO<sub>2</sub>R<sup>14</sup>, lequel est converti en -CO<sub>2</sub>H.

9. Procédé selon la revendication 8, dans lequel R<sup>14</sup> est le radical t-butyle et la réaction est effectuée dans CF<sub>3</sub>CO<sub>2</sub>H.

10. Procédé selon la revendication 7, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est mis au contact de:
- (i) un acide fort à la température du reflux du solvant durant environ 2 à 96 heures; ou
  - (ii) un dérivé alcalin fort dans un solvant formé d'alcool à une température d'environ  $20^{\circ}C$  à la température de reflux du solvant pendant environ 2 à 96 heures, étape que l'on fait suivre de l'amenée du pH à une valeur d'environ 3 - 7; ou
  - (iii) de l'acide sulfurique suivi d'un traitement acide ou alcalin pour convertir le produit en le composé correspondant dans lequel  $R^{13}$  est du  $-CO_2H$ .
11. Procédé selon la revendication 12, dans lequel au moins un des substituants  $R^2$ ,  $R^3$  ou  $R^{13}$  consiste en  $-CO_2R^{14}$  et est converti en  $-CO_2H$ .
12. Procédé selon la revendication 12, dans lequel  $R^8$  est  $-(CH_2)_nCN$  et est converti en  $-(CH_2)_nCO_2H$ , ou est  $-(CH_2)_n-OR^{11}$  et est converti en  $(CH_2)_nOH$  lorsque  $R^{13}$  est converti en  $-CO_2H$ .
13. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est mis au contact d'un mélange formé de proportions équimolaires de l'azidure de sodium et du chlorure d'ammonium dans un solvant aprotique polaire à une température d'environ  $30^{\circ}C$  à la température de reflux du solvant durant environ 1 heure à 10 jours, aux fins de convertir le produit en le composé correspondant dans lequel  $R^{13}$  est le radical 5-tétrazolylo.
14. Procédé selon la revendication 15, dans lequel  $R^8$  est  $-(CH_2)_mCN$  et est converti en radical  $-(CH_2)_m$ -tétrazolylo lorsque  $R^{13}$  est converti en radical 5-tétrazolylo.
15. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-CN$  et le produit de formule 3 est amené à réagir avec l'azidure de trialkylétain ou l'azidure de triarylétaïn, étape que l'on fait suivre d'une hydrolyse acide ou basique pour convertir le produit en un composé correspondant dans lequel  $R^{13}$  est le radical 5-tétrazolylo.
16. Procédé selon la revendication 17, dans lequel  $R^8$  est  $-(CH_2)_nCN$  et est converti en  $-(CH_2)_m$ -tétrazolylo lorsque  $R^{13}$  est converti en radical 5-tétrazolylo.
17. Procédé selon la revendication 8, dans lequel  $R^{13}$  est  $-NO_2$  et le produit de formule 3 est mis au contact d'un agent réducteur pour former un second intermédiaire de formule 3, dans lequel  $R^{13}$  est  $NH_2$ , ce dernier étant mis au contact d'un anhydride  $(CH_3SO_2)_2O$  ou  $(CF_3SO_2)_2O$  ou d'un chlorure  $CH_3SO_2Cl$  ou  $CF_3SO_2Cl$  de l'acide sulfonique dans un solvant pour produire un composé dans lequel  $R^{13}$  est  $-NHSO_2CH_3$  ou  $-NHSO_2CF_3$ .
18. Procédé selon la revendication 17, dans lequel au moins un des substituants  $R^2$ ,  $R^3$  ou  $R^{13}$  est  $-NO_2$  et est converti en  $-NHSO_2CH_3$  ou  $-NHSO_2CF_3$ .
19. Procédé selon la revendication 7 ou 10, dans lequel le composé de formule 3, dans lequel  $R^{13}$  est  $CO_2H$ , est soit:
- (a) mis au contact d'environ 1 à 4 équivalents de chlorure de thionyle dans un excès de chlorure de thionyle ou d'un autre solvant à une température d'environ  $20^{\circ}C$  à la température du reflux du solvant pendant une période d'environ 5 minutes à environ 2 heures pour former un intermédiaire de formule 3 dans lequel  $R^{13}$  est  $COCl$ , et ce dernier est mis au contact d'environ 2 à 10 équivalents d'un dérivé d'hydroxylamine  $H_2NOR^{12}$  dans un excès de dérivé d'hydroxylamine  $H_2NOR^{12}$  ou d'un autre solvant, à une température d'environ  $25$  à  $80^{\circ}C$  pendant environ 2 à 18 heures, soit
  - (b) mis au contact du dérivé d'hydroxylamine  $H_2NOR^{12}$ , de dicyclohexylcarbodiimide et de 1-hydroxybenzotriazole dans un solvant à une température d'environ  $0$  à  $30^{\circ}C$  pendant environ 1 à 24 heures;
- afin d'obtenir un composé dans lequel  $R^{13}$  est  $CONHOR^{12}$ .
20. Procédé selon la revendication 6, dans lequel:
- $R^1$  est:



X est une simple liaison carbone-carbone, -CO-, -O-, -S- ou -NH-;  
 $R^2$ ,  $R^3$ ,  $R^6$  et  $R^7$  sont tels que définis dans la revendication 1 et  
 $R^8$  est  $(CH_2)_nOR^{11}$ ,  $(CH_2)_nOCOR^{14}$ ,  $(CH_2)_nCH(OH)R^{16}$ ,  $(CH_2)_nCOR^{16}-(CH_2)_nCl$ ,  
 $(CH_2)_nCN$ , CHO.

15 21. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis en contact avec un alcool  $R^{11}OH$  à l'état anhydre en présence d'un acide fort ou d'un acide de Lewis, étape que l'on fait suivre d'une saponification de tout groupe  $CO_2R^{14}$  concomitamment formé ou présent sur l'intermédiaire 3, pour former le dérivé correspondant de formule 3, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  n'est pas H.

20 22. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  n'est pas H et dans lequel le produit de formule 3 est mis au contact d'un milieu aqueux acide à des températures d'environ  $25^\circ C$  à la température de reflux du solvant pendant une période d'environ 0,5 à 24 heures pour former le dérivé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nOH$ .

25 23. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis au contact de:  
 (a) un anhydride d'acide carboxylique  $(R^{14}CO)_2O$  ou un chlorure  $R^{14}COCl$  dans un solvant en présence d'une base à une température d'environ  $0^\circ C$  à la température de reflux du solvant pendant environ 0,5 à 24 heures; ou  
 (b) un acide carboxylique  $R^{14}CO_2H$  dans des conditions anhydres en présence d'un acide fort ou d'un acide de Lewis à environ  $0^\circ C$  à  $100^\circ C$  pendant environ 0,5 à 24 heures pour former le dérivé correspondant dans lequel  $R^8$  est  $(CH_2)_nOCOR^{14}$ .

35 24. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOCOR^{14}$  et le produit de formule 3 est mis au contact d'une solution aqueuse acide ou alcaline pour former le dérivé correspondant dans lequel  $R^8$  est  $(CH_2)_nOH$ .

40 25. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOH$  et le produit de formule 3 est mis au contact d'un agent oxydant à une température d'environ  $25^\circ C$  à  $45^\circ C$  pendant environ 1 à 200 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_{n-1}COR^{16}$  et  $R^{16}$  est H.

45 26. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est H et le produit de formule 3 est mis au contact d'un composé organométallique  $R^{16}P$  dans lequel P est MgBr ou Li dans un solvant, à une température d'environ  $-78^\circ C$  à  $100^\circ C$  pendant environ 0,5 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(OH)R^{16}$  et  $R^{16}$  n'est pas H.

50 27. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nCH(OH)R^{16}$  et  $R^{16}$  n'est pas H, et le produit de formule 3 est mis au contact d'un agent oxydant dans un solvant pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  n'est pas de l'hydrogène.

55 28. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est de l'hydrogène et le produit de formule 3 est mis au contact d'un agent oxydant dans un solvant pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est OH.

29. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est OH et le produit de formule 3 est mis au contact de chlorure de thionyle en excès ou dans un autre solvant à une température d'environ  $0^\circ C$  à la température de reflux du solvant pendant environ 5 minutes à environ



- 24 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOCl$ , étape que l'on fait suivre d'un contact de ce dernier composé avec une amine  $NHR^{18}R^{19}$  en excès dans un solvant à une température d'environ  $0^\circ C$  à la température du reflux du solvant pendant environ 5 minutes à environ 24 heures pour former un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCONR^{18}R^{19}$ .
- 5
30. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nOR^{11}$  et  $R^{11}$  est un atome d'hydrogène et le produit de formule 3 est mis au contact de chlorure de thionyle en excès ou dans un solvant à une température d'environ  $20^\circ C$  à la température de reflux du solvant pendant environ 0,5 à 24 heures pour produire un composé intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$ .
- 10
31. Procédé selon la revendication 30, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mCl$  est mis au contact d'un imidazole, de 1,2,3-triazole, de 1,2,4-triazole ou de phthalimide en présence d'une base dans un solvant à une température d'environ  $55^\circ C$  à la température de reflux du solvant pendant environ 1 à 24 heures pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_m$ -imidazole,  $(CH_2)_m$ -triazole,  $(CH_2)_m$ -tétrazole ou  $(CH_2)_m$ -phthalimide.
- 15
32. Procédé selon la revendication 30, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$  est mis au contact d'un sel de sodium ou de potassium d'un mercaptan  $R^{15}SH$ , dans un solvant, à une température d'environ  $25^\circ C$  à  $100^\circ C$ , pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_n-SR^{15}$ .
- 20
33. Procédé selon la revendication 30, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCl$  est mis au contact d'un cyanure de métal alcalin dans un solvant à une température d'environ  $20^\circ C$  à  $100^\circ C$ , pendant 1 à 24 heures, pour produire un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCN$  et ce dernier composé est hydrolysé en un composé correspondant de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^6$  ET  $R^{16}$  est  $OH$ .
- 25
34. Procédé selon la revendication 30, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_{n-1}Cl$  est mis au contact d'un sel de sodium ou de potassium d'un malonate de dialkyle, dans un solvant, à une température de  $20^\circ C$  à  $100^\circ C$ , pendant environ 0,5 à 24 heures, pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(CO_2alkyl)_2$ , étape que l'on fait suivre d'une saponification de ce dernier composé par action d'une solution alcaline aqueuse à une température d'environ  $25^\circ C$  à la température du reflux du solvant, ce que l'on fait suivre d'une acidification à l'aide d'un acide minéral pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCH(CO_2H)_2$  suivie d'un chauffage de ce dernier composé à environ  $120^\circ C$  dans une solution diluée d'un acide minéral à la température du reflux pour former un produit de formule 3 dans lequel  $R^8$  est  $(CH_2)_nCOR^{16}$  et  $R^{16}$  est  $OH$ .
- 30
35. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_nCN$  et le composé de formule 3 est mis au contact d'azide de sodium et de chlorure d'ammonium dans un solvant à une température d'environ  $30^\circ C$  à la température de reflux du solvant pendant environ 1 heure à environ 10 jours pour former un composé selon la présente invention dans lequel  $R^8$  est  $(CH_2)_n$ -tétrazole.
- 40
36. Procédé selon la revendication 20, dans lequel  $R^8$  est  $-CHO$  et le composé de formule 3 est mis au contact d'un méthylène phosphorane  $(C_6H_5)_3P=CH(CH_2)_sCHR^{14}OR^{15}$  ou  $(C_6H_5)_3P=CH(CH_2)_sCOR^{16}$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCHR^{14}OR^{15}$  ou  $-CH=CH(CH_2)_sCOR^{16}$ , excepté dans le cas où  $R^{15}$  est  $H$  et  $R^{16}$  est  $OH$ , et éventuellement mise en contact du composé de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCOR^{16}$  avec un agent réducteur dans un solvant à une température d'environ  $0^\circ C$  à  $25^\circ C$  pendant environ 0,5 à 24 heures, pour former un produit de formule 3 dans lequel  $R^8$  est  $-CH=CH(CH_2)_sCHR^{14}OH$ .
- 45
37. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_mOH$  et le composé de formule 3 est mis au contact d'un agent de fluoration dans un solvant à une température d'environ  $-30^\circ C$  à  $25^\circ C$  pendant une période d'environ 0,5 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_mF$ .
- 50
- 55

38. Procédé selon la revendication 20, dans lequel le composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_m Cl$ , est mis au contact de nitrate d'argent dans un solvant aprotique dipolaire à une température d'environ  $25^\circ C$  à  $80^\circ C$  pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_m ONO_2$ .

39. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_n OH$  et le composé de formule 3 est mis au contact d'un isocyanate de formule  $R^{10} NCO$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant une période d'environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_n OCONHR^{10}$ .

40. Procédé selon la revendication 20, dans lequel le composé où  $R^8$  est  $(CH_2)_n Cl$  est mis au contact d'une amine  $R^{11} NH_2$  dans un excès d'amine ou dans un autre solvant pendant une période d'environ 1 à 24 heures à une température d'environ  $0^\circ C$  à la température de reflux du solvant pour former un intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_n NHR^{11}$ .


41. Procédé selon la revendication 20, dans lequel  $R^8$  est  $(CH_2)_n Cl$  et le composé de formule 3 est mis au contact d'un azidure de métal alcalin dans un solvant aprotique à une température d'environ  $25^\circ C$  à  $80^\circ C$  pendant environ 1 à 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $(CH_2)_n N_3$  et ce dernier est alors mis au contact d'un agent réducteur pour former un intermédiaire de formule 3 dans lequel  $R^8$  est  $(CH_2)_n NH_2$ .

42. Procédé selon la revendication 40 ou 41, dans lequel  $R^8$  est  $(CH_2)_n NHR^{11}$  ou  $(CH_2)_n NH_2$  et le composé de formule 3 est mis au contact d'un chloroformiate de formule  $R^{10} OCOCl$  ou d'un dérivé sulfonyl de formule  $R^{10} SO_2 Cl$  ou  $(R^{10} SO_2) O$  dans un solvant en présence d'une base à une température d'environ  $0^\circ C$  à la température du reflux dans le solvant pendant environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-(CH_2)_n NR^{11} CO_2 R^{10}$  ou  $-(CH_2)_n NR^{11} SO_2 R^{10}$ .

43. Procédé selon la revendication 40 ou 41, dans lequel le composé de formule 3 où  $R^8$  est  $(CH_2)_n NHR^{11}$  ou  $(CH_2)_n NH_2$  est mis en contact d'un isocyanate ou d'un isothiocyanate  $R^{10} NCY$  dans un solvant à une température d'environ  $25^\circ C$  à la température de reflux du solvant pendant environ 5 minutes à environ 24 heures pour former un composé de formule 3 dans lequel  $R^8$  est  $-(CH_2)_n -NR^{11} CYNHR^{10}$ .

44. Procédé selon la revendication 1, dans lequel  $R^1$  est  $NO_2$ ,  $R^2$ ,  $R^3$ ,  $R^6$ ,  $R^7$  et  $R^8$  sont tels que définis dans la revendication 1 et dans lequel le composé de formule 3 où  $R^1$  est  $NO_2$  est réduit au moyen de fer et d'acide acétique, de chlorure stanneux ou d'hydrogène et de palladium en un composé de formule 3 dans lequel  $R^1$  est  $NH_2$ , le composé ainsi obtenu étant amené à réagir avec un anhydride d'acide convenable tel qu'un anhydride phtalique ou un anhydride phtalique substitué, dans un solvant, ou avec un chlorure d'acide approprié tel que le chlorure d'acide anthranilique substitué, en présence d'une solution aqueuse alcaline ou d'une base, ou avec un acide anthranilique ou phtalique convenablement substitué, en présence de dicyclohexylcarbodiimide dans un solvant pour produire un composé de formule 3 dans laquelle:

$R^1$  est



et  
X est  $NHCO$ .


45. Procédé selon la revendication 1, dans lequel  $R^1$  est  $OCH_2 C_6 H_5$ ,  $R^2$  et  $R^3$  sont H, et  $R^6$ ,  $R^7$  et  $R^8$  sont tels que définis dans la revendication 1 et dans lequel le composé résultant de formule 3 où  $R^1$  est le radical  $OCH_2 C_6 H_5$  est mis au contact d'acide trifluoroacétique à la température du reflux pendant une



période d'environ 0,2 à 1 heure, ou d'hydrogène et de palladium pour former le composé correspondant de formule 3 dans lequel R<sup>1</sup> est OH et ce dernier est mis au contact d'une base à environ 25°C et d'un halogénure benzylique convenable de formule:

5

10




15

pour produire le composé correspondant de formule 3 dans lequel:  
R<sup>1</sup> est

20

25



30

46. Procédé selon la revendication 1, dans lequel R<sup>8</sup> est -CHO, de façon à ce que le dérivé benzylique de formule 2 se lie au dérivé imidazole de formule 1 de préférence sur l'atome d'azote adjacent à l'atome de carbone du cycle imidazole auquel R<sup>8</sup> est lié.

35

40

45

50

55

**THIS PAGE BLANK (USPTO)**